



Journal of Epidemiology and Global Health

ISSN (Online): 2210-6014

ISSN (Print): 2210-6006

Journal Home Page: <https://www.atlantis-press.com/journals/jegh>

Prevalence of self-reported diabetes, hypertension and heart disease in individuals seeking State funding in Trinidad and Tobago, West Indies

Derek Chadee, Terence Seemungal, Lexley M. Pinto Pereira, Mary Chadee, Rohan Maharaj, Surujpal Teelucksingh

To cite this article: Derek Chadee, Terence Seemungal, Lexley M. Pinto Pereira, Mary Chadee, Rohan Maharaj, Surujpal Teelucksingh (2013) Prevalence of self-reported diabetes, hypertension and heart disease in individuals seeking State funding in Trinidad and Tobago, West Indies, Journal of Epidemiology and Global Health 3:2, 95–103, DOI: <https://doi.org/10.1016/j.jegh.2013.02.002>

To link to this article: <https://doi.org/10.1016/j.jegh.2013.02.002>

Published online: 23 April 2019



Prevalence of self-reported diabetes, hypertension and heart disease in individuals seeking State funding in Trinidad and Tobago, West Indies

Derek Chadee ^a, Terence Seemungal ^b, Lexley M. Pinto Pereira ^{c,*},
Mary Chadee ^a, Rohan Maharaj ^c, Surujpal Teelucksingh ^b

^a *Departments of Behavioural Sciences, The University of the West Indies, St. Augustine, Trinidad and Tobago*

^b *Clinical Medical Sciences, The University of the West Indies, St. Augustine, Trinidad and Tobago*

^c *Para Clinical Sciences, The University of the West Indies, St. Augustine, Trinidad and Tobago*

Received 9 October 2012; received in revised form 4 February 2013; accepted 6 February 2013
Available online 16 March 2013

KEYWORDS

Hypertension; Diabetes;
Heart disease; Gender
specificity; Age;
Education

Abstract Objective: Diabetes, hypertension and heart disease inflict a heavy health burden on the Caribbean Republic of Trinidad and Tobago. This study assessed the prevalence of self-reported diabetes, hypertension and heart disease in lower socioeconomically placed individuals accessing welfare grants.

Method: Data collected between July 2008 and June 2009 were analyzed from 14,793 responses. The survey sought information on education, average monthly income, health, housing, and household facilities.

Results: Self-reported disease prevalence was 19.5% (95% CI: 18.9–20.2) for diabetes mellitus; 30.2% (95% CI: 29.5–30.9) for hypertension; and 8.2% (95% CI: 7.7–8.6) for cardiac disease. Diabetes and cardiac disease had equivalent gender frequency; hypertension was more prevalent in women ($p < .001$). Disease prevalence was highest in Indo-Trinidadians, married and divorced subjects, non-Christians and increased with age. Those with primary education alone were at greatest risk.

Conclusion: Trinidad and Tobago have a high prevalence of hypertension, diabetes and heart disease. Hypertension showed gender specificity in women. Prevalence was highest in Indo-Trinidadians, increased with age, and primary education

* Corresponding author. Address: Faculty of Medical Sciences, The University of the West Indies, St. Augustine, Trinidad and Tobago. Tel./fax: +1 868 663 8613.

E-mail addresses: dachadee@yahoo.com (D. Chadee), tseemungal@aol.com (T. Seemungal), lexleyp@gmail.com (L.M. Pinto Pereira), mchadee@yahoo.com (M. Chadee), Rohan.Maharaj@sta.uwi.edu (R. Maharaj), pteelucksingh@gmail.com (S. Teelucksingh).

alone was a risk factor. Interventions to arrest the high prevalence of chronic non-communicable diseases to promote wellness are needed in Trinidad and Tobago.

© 2013 Ministry of Health, Saudi Arabia. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Chronic non-communicable diseases (CNCDs) are viewed to be the leading causes of global death and disability in the not so far off 2030 and are predicted to cause three quarters of all deaths [1]. Cardiovascular disorders, chronic obstructive pulmonary disease, asthma, diabetes, obesity, some cancers and disabling mental illness are included under the umbrella of this acronym. CNCDs have dominated the global health burden in recent years, and poorer countries seem to have provided the model for this observation. Evidence clearly points to CNCDs as disproportionately affecting the poor [2]. In low- and middle-income countries, chronic disease is the current cause for an estimate of more than 80% of deaths [1,3]. Countries economically not on par with the developed nations bear the biggest brunt of these diseases, and Caribbean territories with less than enviable economies are significantly saddled with this health encumbrance. In 1990 non-communicable diseases surpassed maternal, peri-natal, communicable, and nutritional disorders in the Caribbean [4]. More than a decade later the health scenario suggests these diseases are still the leading causes of morbidity. The Caribbean has the highest prevalence of CNCDs in the region of the Americas [5]. In the twin-island republic of Trinidad and Tobago, the mortality rates from diabetes and cardiovascular disease are higher than in North America (United States and Canada) [5].

The epidemiologic transition to CNCDs has been well documented in Trinidad and Tobago [6,7]. The first survey of a stratified random sample of the entire population of Trinidad screened 24,069 persons in the 1960s and reported that 1.89% of the sample satisfied the criteria for diabetes [8], which was one of the highest reported prevalence for this disease in the region. In a second study conducted between 1977 and 1985 of 2491 persons in an urban setting, the incidence rates of hypertension in men were 37 per 1000 person-years and 29.5 per 1000 person-years in women. The incidence of diabetes (per 1000 person-years) in Indians was 23.5/1000 and was significantly higher than in Africans [9]. However, following these two early population-based epidemiologic studies, more recent data on the epidemic have been based on self-reporting of the CNCDs. Thus, the 2005 Survey of Living Condi-

tions reports that 10.4% of the poorest and 17.6% in the richest of 2086 community dwelling respondents had self-reported diabetes. In this survey no information was provided for hypertension [10]. These data provide an insight into the evolution of the diabetes epidemic in Trinidad and Tobago. The death rate from diabetes in the country is ten times higher than in the United States of America, and the age-adjusted, gender-specific mortality rates from heart disease and all cardiovascular diseases rank among the highest in the Caribbean [11].

Although population-based epidemiologic surveys are the gold standard, information derived from self-reporting of diabetes in a Taiwanese population has been found to be accurate in determining disease prevalence [12]. Further, in a USA study of 2037 persons self-reporting of diabetes when compared with a review of their medical records revealed a specificity of >90% but with lower (66%) sensitivity [13]. The self reported prevalence of diabetes, hypertension and heart disease in a large population of individuals from the lower socioeconomic group who applied for State-sponsored welfare grants was studied. The determined prevalence rates of these conditions, particularly diabetes and hypertension – the two most common chronic non-communicable diseases in Trinidad and Tobago – will contribute to the health care system planning over the next decade.

2. Materials and methods

2.1. Ethical permission

This is a retrospective database study. Data from patients were anonymized and permission was obtained from the Ministry of the People and Social Development for analysis of these data. It was not deemed necessary to obtain ethical permission from an Institutional Review Board as these data were entered on a database of the above-mentioned Ministry and were taken with the informed consent of respondents and were anonymously analyzed.

2.2. Population

Respondents came from a short-term financial assistance program which is offered to economically vulnerable persons and families within Trini-

dad and Tobago. Eligibility for this program is determined using a Means Test Score applied by the Ministry of the People and Social Development.

2.3. Data collection

Data were collected from persons who accessed the program from its inception in August 2006 to January 31, 2008. The exercise targeted 21,790 recipients of the program. A database of recipients was provided in May 2008 by the government agency responsible for the implementation of the program. Data were formally collected during the period July 2008 to June 2009. Correspondence was sent by the Ministry of the People and Social Development to all persons listed on a database that were accessing this program via mailings and advertisements inviting them to attend an interview session at specific centers across Trinidad and Tobago on a specific date, time and venue. For the purpose of administering the interview, Trinidad and Tobago was divided into five zones (East, Northeast, South, Central and Tobago) and 23 centers manned by supervisors and interviewers were set up within these zones. The invitation letters to recipients indicated the specific documents required to be provided at the interview. A household interview was subsequently conducted with each respondent who attended the initial interview. When respondents did not attend the first interview, attempts to obtain the data were made by house visits. A field verification exercise was conducted after the interview to verify living conditions.

2.3.1. Instrument of collection

Data were collected via interviewer administered application forms, using information provided by recipients. A 32-page manual was developed and utilized in the training of interviewers before the start of the exercise. The main instrument of data collection was a 27-page application form developed and prepared in consultation with a governmental agency and consisted of sections on characteristics of household members, levels of educational attainment, average monthly income, health/disease status, housing, and household facilities. The following demographic data were all subject-defined: ethnicity, income level, educational attainment, religious affiliation, marital status and residence.

3. Results

3.1. Description of population

Though attempts were made to interview all relevant recipients (21,790), only 15,649 persons were

interviewed. Data were incomplete for 856 participants; the remaining 14,793 interview responses were utilized for this paper. The final demographic configuration of recipients was: 86% females and 14% males with the mean age being 46.4 years (SD = 12.9).

3.2. Age and ethnicity

On a more detailed examination of age, 13% of recipients were between 17 and 30 years, 22% between 31 and 40 years, 26% between 41 and 50 years, 23% between 51 and 60 years, 14% between 61 and 70 years and 2% were over 71 year old; 65% of recipients were 41 years and over, 49% of respondents were of East Indian origin, 35% were of African origins, 15% were of mixed origin, and 1% belonged to other ethnic groups. The marital status showed that 29% of recipients were married, 29% were single, 14% were in a 'common law' relationship, 13% were widowed, 9% were separated, 4% were divorced and for 1% no information was given.

3.3. Religious affiliation

The distribution of religious backgrounds of recipients is as follows: 28% Hindu, 17% Pentecostal, 16% Roman Catholics, 11% Baptist, 6% Anglicans, 5% Islam, 5% Seven Day Adventists (SDAs), 3% Presbyterian, 1% Jehovah Witness, 1% Methodist, less than 1% Moravian, 6% other and in 1% none. The other religious groups included Assembly Church of God, Bahai, Ethiopian Orthodox, Open Bible, Rastafarian and Spiritual Baptist.

3.4. Education and Income

The vast majority of recipients (63%) had only a primary education, 34% had a secondary education, less than 1% were in a tertiary education at university or taking distance learning programs, less than 1% were enrolled in an adult literacy program and 2% had no education. The mean monthly income of recipients was TT\$ 430.36. The mean monthly income and monthly expenditure of each household were TT \$1,715.34 and TT\$ 1,050.38 respectively, so that the net income was TT\$ 664.98 (income and expenditure are based upon means test criteria). (1US\$ = 6.44 TT\$).

3.5. Geographic location

Respondents came from the following regional areas as determined by the regional offices: 7.7% from Port of Spain, 14.3% from San Juan, 12.3% from Sangre Grande, 9.3% from Chaguanas, 9.6%

from Couva, 7.7% from San Fernando, 15.7% from Princes Town, 11.6% from Penal, 5.2% from Point Fortin and 6.6% were from Tobago.

3.6. Marital status

Table 1 shows that single and married subjects within the studied group were approximately equal in prevalence. Only 3.7% were divorced.

3.7. Medical illness in the population interviewed

Diabetes mellitus was reported by 19.5% (95% CI: 18.9–20.2), hypertension by 30.2% (95% CI: 29.5–30.9) and cardiac disease was reported by 8.2% (95% CI 7.7–8.6) of the studied group. Both diabetes and cardiac disease were approximately equally prevalent among males and females, but hyperten-

Table 1 The composition of individuals seeking State-funded, short-term public financial assistance in Trinidad and Tobago (2008).

	Self-reported (N)	Self-reported (%)
<i>Gender</i>		
Male	1990	14.1
Female	12139	85.8
<i>Ethnicity</i>		
Afro-Trinidadian	4989	35.5
Indo-Trinidadian	6978	49.6
Mixed	2099	14.9
<i>Marital status</i>		
Single	4165	29.9
Married	4110	29.5
Separated	1267	9.1
Divorced	515	3.7
Common Law	1987	14.2
Widowed	1902	13.6
<i>Religion</i>		
Hinduism	3956	30.1
Islam	646	4.9
Roman Catholic	2197	16.7
SDA	709	5.4
Other christian groups	5616	42.8
<i>Age</i>		
17–30	1835	13.0
31–40	3108	22.0
41–50	3610	25.5
51–60	3312	23.4
61–70	2002	14.2
71+	267	1.9
<i>Education</i>		
Primary	9 213	65.5
Secondary	4 735	33.7
University	110	.8
<i>Diabetes mellitus</i>		
Yes	2 758	19.7
No	11 273	80.3
<i>Hypertension</i>		
Yes	4 272	30.4
No	9 764	69.6
<i>Cardiac disease</i>		
Yes	1 156	8.3
No	12 820	91.7

Table 2 Prevalence of NIDDM, HTN and Cardiac Disease in individuals seeking financial assistance in a State-funded, short-term support program (2008).

	Diabetes mellitus						Hypertension		Cardiac Disease			
	Self-reported	Yes					Self-reported	Yes	Self-reported	Yes		
	(%)	N	χ^2	p-value	(%)	N	χ^2	p-value	(%)	N	χ^2	p-value
<i>Gender</i>												
Male	18.6 ^a	367	1.60	.21	22.0	434	76.92	.000	10.7	211	18.27	.000
Female	19.8	2389			31.8	3832			7.9	944		
<i>Ethnicity</i>												
Afro-Trinidadian	13.0	644	435.03	.000	25.5	1266	173.33	.000	4.4	219	264.24	.000
Indo-Trinidadian	26.7	1850			35.6	2470			12.1	835		
mixed	12.1	251			25.0	52			4.8	99		
<i>Marital status</i>												
Single	14.9	614	423.29	.000	25.1	1036	546.13	.000	5.6	233	236.18	.000
Married	21.0	858			30.5	1249			8.7	353		
Separated	19.8	249			30.5	383			8.1	102		
Divorced	21.1	108			33.9	173			10.2	52		
Common law	12.0	237			20.9	412			4.8	94		
Widowed	35.1	663			51.7	977			16.4	308		
<i>Religion</i>												
Hinduism	27.0	1060	218.48	.000	36.3	1428	117.19	.000	13.3	519	199.26	.000
Islam	23.4	150			33.0	211			10.5	67		
Roman Catholic	14.9	326			25.1	546			5.3	114		
SDA	12.2	86			23.2	163			3.7	26		
Other Christian Groups	17.0	951			29.3	1633			6.5	362		
<i>Age</i>												
17–30	2.9	53	1326.13	.000	7.5	136	1794.90	.000	1.0	18	699.61	.000
31–40	7.7	239			15.0	462			2.2	69		
41–50	16.9	605			26.2	940			6.1	218		
51–60	32.4	1064			46.8	1537			13.7	446		
61–70	36.7	729			54.0	1072			18.2	360		
71+	25.6	68			46.2	123			16.5	44		
<i>Education</i>												
Primary	24.8	2267	438.34	.000	36.9	3379	521.38	.000	10.8	980	210.78	.000
Secondary	10.0	472			18.3	859			3.6	170		
University	7.3	8			15.5	17			2.8	3		

^a Column percent.

