

Journal of Epidemiology and Global Health

ISSN (Online): 2210-6014

ISSN (Print): 2210-6006

Journal Home Page: <https://www.atlantis-press.com/journals/jegh>


Burden of chronic kidney disease and its risk factors in Malaysia

Rubina Begum, Tahir Mehmood Khan, Long Chiau Ming

To cite this article: Rubina Begum, Tahir Mehmood Khan, Long Chiau Ming (2017)
Burden of chronic kidney disease and its risk factors in Malaysia, Journal of
Epidemiology and Global Health 6:4, 325–326, DOI:
<https://doi.org/10.1016/j.jegh.2016.03.004>

To link to this article: <https://doi.org/10.1016/j.jegh.2016.03.004>

Published online: 23 April 2019



Burden of chronic kidney disease and its risk factors in Malaysia



The dialysis acceptance and prevalence rates in Malaysia have almost doubled in recent years. For instance, the dialysis acceptance was 1733 per million population (pmp) in 2004 increasing to 3156 pmp in 2013. Based on the National Renal Registry 2014, the total number of patients on dialysis has increased sharply from 6702 in 2000 to 31,637 in 2013, causing a serious strain on Malaysian health resources.

This increase in kidney failure is mainly due to the increasing incidence of diabetic kidney disease, which is responsible for 58.0% of new end stage renal disease patients. In fact, Malaysia had the highest incident rate of patients with diabetes entering renal replacement therapy [1]. The National Health and Morbidity survey in Malaysia shows that 32.7% of adults aged ≥ 18 years have hypertension (12.8% are known to have hypertension while 19.8% have hypertension that is as yet undiagnosed). Meanwhile, 15.2% (2.6 million) of adults aged ≥ 18 years are diabetic (7.2% are known to have diabetes while 8.0% have diabetes that is as yet undiagnosed). In addition, about 11.3% of patients with dengue viral infection experience acute kidney injury [2]. Studies have also shown metabolic syndrome to be an independent risk factor for chronic kidney disease and the older population (aged >65 years) are also at increased risk of renal impairment. Autoimmune disease and smoking also increase the risk of kidney diseases [3]. Environmental factors such as groundwater contaminated with microbial and toxic heavy metals can damage the kidneys as well, according to a recent study [4].

Since kidney disease in its initial stage is asymptomatic, patients tend to present late and with more aggressive symptoms of chronic kidney disease. This means that, to reduce the load of this disease in Malaysia, targeted screening and timely intervention is necessary. In addition, the Ministry of Health, together with the National Diabetes Institute, the Malaysian Society of Hypertension, and the Malaysian Society of Nephrology should intervene to increase patient awareness. Under the National Strategic Plan for Non-Communicable Disease (2010–2014) seven strategies including primary prevention and promotion; clinical management; increasing patient compliance; action with non-governmental organisation, professional bodies, and other stakeholders; monitoring, research, and surveillance; capacity building; and policy and regulatory interventions were implemented. However, more drastic strategies are needed to ensure optimal diabetic control and to reduce the renal disease burden in Malaysia by 2025.

Conflicts of interest

We declare no competing interests.

Acknowledgment

No funding was received.

References

- [1] Huri HZ, Lim LP, Lim SK. Glycemic control and antidiabetic drugs in type 2 diabetes mellitus patients with renal complications. *Drug Des Devel Ther* 2015;9(Suppl 3): 4355–71.

Peer review under responsibility of Ministry of Health, Saudi Arabia.

<http://dx.doi.org/10.1016/j.jegh.2016.03.004>

2210-6006/Crown Copyright © 2016 Ministry of Health, Saudi Arabia, Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

- [2] Mallhi TH, Khan AH, Sarriif A, Syazril AA, Khan YH, Jummaat F. Predictors of the development of acute kidney injury in dengue viral infection: a retrospective study. *Nephrol Dial Transplant* 2015;30(Suppl 3):iii457.
- [3] Hazreen MA, Su TT, Jalaludin MY, Dahlui M, Chinna K, Ismail M, et al. An exploratory study on risk factors for chronic non-communicable diseases among adolescents in Malaysia: overview of the Malaysian Health and Adolescents Longitudinal Research Team study (The MyHeART study). *BMC Public Health* 2014;14(Suppl 3):S6.
- [4] Ambu S, Yong SFY, Lim YAL, Wah MJ, Chen DKF, Ooi SS, et al. The chemical, heavy metal and microbial quality of well water in an urbanised village in the Klang Valley. *Int e-J Sci Med Educ* 2014;8(Suppl 3):28–44.

Rubina Begum
Tahir Mehmood Khan
*School of Pharmacy, Monash University,
Bandar Sunway, Selangor 45700, Malaysia*

Long Chiau Ming*
*Unit for Medication Outcomes Research and
Education (UMORE), Pharmacy,
School of Medicine, University of Tasmania,
Private Bag 26,* Hobart 7001, Australia*
Corresponding author.
E-mail address: long.ming@utas.edu.au
(L.C. Ming)

Received 30 September 2015; revised 3 March 2016;
accepted for publication 14 March 2016
Available online 16 April 2016

Available online at www.sciencedirect.com

ScienceDirect