



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P1.05: EFFECT OF AGING ON THE ANNUAL CHANGES OF CENTRAL HEMODYNAMIC INDICES IN MIDDLE-AGED MEN: A PROSPECTIVE EVALUATION

H. Tomiyama, A. Yamashina

To cite this article: H. Tomiyama, A. Yamashina (2011) P1.05: EFFECT OF AGING ON THE ANNUAL CHANGES OF CENTRAL HEMODYNAMIC INDICES IN MIDDLE-AGED MEN: A PROSPECTIVE EVALUATION, Artery Research 5:4, 150–150, DOI: <https://doi.org/10.1016/j.artres.2011.10.011>

To link to this article: <https://doi.org/10.1016/j.artres.2011.10.011>

Published online: 14 December 2019

waveforms. We used it here to investigate whether systolic ventricular function is different in patients requiring chronic beta-blockade (BB) compared to controls.

Methods and Results: In 81 patients on BB and 87 age- (61 ± 7 yrs) and sex-matched untreated controls, we obtained untreated STI and blood pressures. BB patients discontinued medication 1 day prior to measurements. Intra-session coefficients of variation for T_{ej} and T_{ic} were 0.8% and 6.8%; and inter-session 2.7% and 8.4% (i.e. $100\% \cdot SD_{intra/inter}/mean$).

T_{ej} and pulse pressure (PP), thus the $PP \cdot T_{ej}$ product reflecting ventricular stroke work, were significantly greater in (untreated) BB patients (all $p < 0.002$). T_{ic} was not different between the groups. The T_{ic}/T_{ej} ratio was smaller for the BB group ($p = 0.04$), indicating increased systolic ventricular function, likely matching the greater stroke work required.

	T_{ic} ms	T_{ej} ms	R-R int. ms	T_{ic}/T_{ej} %	DBP mmHg	SBP mmHg	PP mmHg	$PP \cdot T_{ej}$ mmHg · s
BB	34 ± 7	286 ± 27	908 ± 148	12 ± 3	75 ± 10	137 ± 19	61 ± 13	18 ± 5
Ctrls	35 ± 7	274 ± 22	885 ± 145	13 ± 3	77 ± 10	133 ± 17	56 ± 11	15 ± 3
*p-val.	0.36	0.0017	0.32	0.04	0.41	0.15	0.006	0.0005

*Paired t-test. T_{ic}/T_{ej} : approximate Tei index; $PP \cdot T_{ej}$: stroke work (pressure-time product).

Conclusions: Patients requiring chronic beta-blockade have longer ventricular ejection periods combined with greater pulse pressures, indicating greater ventricular stroke work and thus enhanced systolic function. Our findings readily demonstrate the accessibility and reproducibility of non-invasive STI measurement in a clinical population.

P1.05

EFFECT OF AGING ON THE ANNUAL CHANGES OF CENTRAL HEMODYNAMIC INDICES IN MIDDLE-AGED MEN: A PROSPECTIVE EVALUATION

H. Tomiyama, A. Yamashina
Tokyo Medical University, Tokyo, Japan

Objective: To examine the effects of aging on the annual changes of the pressure wave reflection (PWR) and central blood pressures (CBPs), prospectively. Background: Such prospective study has not been conducted.

Methods: In 1291 middle-aged men, the first and second peaks of the radial systolic pressure waveform (SBP1 and SBP2), the radial augmentation index (rAI), reflected wave transit time (RWTT) and the brachial-ankle pulse wave velocity (baPWV), were measured twice in a 3-years' interval.

Results: Subjects were divided into three groups {aged <40 years old (aged 30); <40 aged <50 years old (aged 40); and aged >50 years old (aged 50)}. The change of rAI was higher and that of RWTT was lower in aged 30 group than in aged 50 group ($p < 0.01$), but that of SBP2 was similar among the groups. The change of RWTT, but not that of baPWV, had a significant inverse relationship with that of rAI in all groups ($p < 0.01$). Stepwise multivariate analysis demonstrated that the accounting rate of change of SBP1, but not that of rAI, for the variance in that of SBP2 was increased in a phased manner along with age.

Conclusion: In middle-aged men, aging may differently affect PWR and CBPs. Aging may prolong the time of PWR and blunt that arterial stiffness increases PWR. These phenomena may contribute to the aging-related attenuation of annual increase of PWR. However, aging may not affect the annual elevations of CBPs, because the incident pressure wave may compensate the aging-related attenuation of their elevations by PWR.

P1.06

BLOOD PRESSURE VARIABILITY IN RELATION TO AUTONOMIC NERVOUS SYSTEM DYSREGULATION: THE X-CELLENT STUDY

Y. Zhang^{1,2}, D. Agnoletti¹, J. Blacher¹, M. E. Safar¹
¹Paris Descartes University; AP-HP; Diagnosis and Therapeutic Center, Hôtel-Dieu, Paris, France
²Centre for Epidemiological Studies and Clinical Trials, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

To investigate the association of autonomic nervous system dysregulation with blood pressure variability. Of 2370 participants in the X-CELLENT study, 577 patients (59.0 ± 10.2 years) were randomly selected to participate in an ambulatory blood pressure monitoring ancillary study. We proposed a novel autonomic nervous system regulation index termed dSBP/dHR, which was defined as the steepness of the slope of the relationship between 24h systolic blood pressure and heart rate for each participant. Within-subject standard deviation of systolic blood pressure, weighted for the time interval between consecutive validated readings from 24h ambulatory blood pressure monitoring, was used to evaluate blood pressure variability. When dSBP/dHR

was divided into tertiles, from tertile 1 to tertile 3, we observed a progressive increase in daytime systolic blood pressure, a progressive decrease in nighttime systolic blood pressure, and consequently a progressive increase in day-night systolic blood pressure gradient ($P < 0.001$). On the contrary, standard deviation of both daytime and nighttime systolic blood pressure were consistently and significantly increased from tertile 1 to tertile 3 ($P < 0.01$). Both before and after adjustment for age, gender and 24h mean blood pressure, all of these increasing or decreasing trends reached statistical significance ($P < 0.01$). Furthermore, in our sensitivity analysis, when men and women were considered separately, the present finding remained unaltered. In summary, autonomic nervous system dysfunction was associated with the enlarged day-night systolic blood pressure gradient and more variable systolic blood pressure in 24 hours in patients with essential hypertension.

P1.07

CAROTID FUNCTION AND BARORECEPTOR SENSITIVITY IN MODERATE CHRONIC KIDNEY DISEASE: THE EPP3 STUDY

L. Zanolì^{1,3}, M. Alivon¹, N. Estrugo¹, G. Ecriou¹, H. Ketthab¹, J. F. Prunty¹, P. Castellino³, S. Yanes¹, D. Laude¹, K. Bean¹, F. Thomas², J. P. Empana¹, X. Jouven¹, S. Laurent¹, P. Boutouyrie¹
¹Université Paris Descartes, Department of Pharmacology, Hôpital Européen Georges Pompidou, APHP, Paris, France
²Institut Prévention cardiovasculaire, Paris, France
³University of Catania, Department of Internal Medicine, Catania, Italy

Introduction: Short-term variation of blood pressure (BP) is largely controlled by autonomic function though the baroreflex. Carotid distension rate was recently introduced instead of BP to evaluate the carotid function and neural component of the baroreflex in small populations. Autonomic dysfunction and arterial stiffness occurs in patients with severe chronic kidney disease (CKD) but little is known in moderate CKD.

Aims: To study the baroreflex and to analyse the link with the carotid function in moderate CKD.

Methods: From the EPP3 cohort, 123 patients with moderate CKD (Stage 3A) and 615 controls with $GFR \geq 60 \text{ ml/min/1.73m}^2$, matched for age, gender and body surface area were enrolled (age 64 ± 6 years). Carotid measurements were performed by a high-resolution echotracking device. Spontaneous BRS was calculated with the fast Fourier transform of carotid distension rate and R-R interval in the low-frequency (LF) range (0.04-0.15 Hz).

Results: Internal diastolic diameter, intima-media thickness, circumferential wall stress, carotid pulse pressure, R-R interval and BRS were comparable between the two groups. Carotid strain and distensibility were significantly reduced, elastic incremental modulus and carotid stiffness were significantly increased in CKD.

Neural baroreflex appeared very sensitive to vascular component since carotid strain was the strongest determinants of BRS in both groups (Fig-1). The explained BRS variability was higher in patients with CKD ($R^2 = 0.31$) than in controls ($R^2 = 0.14$).

Conclusions: In moderate CKD we detected carotid dysfunction and no changes in neural baroreflex. The role of carotid strain, as a determinant of neural baroreflex, is confirmed in moderate CKD and in controls.




Fig -1