



## Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

---

### **P7.10: RENAL FUNCTION DOES NOT MODIFY PREDICTIVE VALUE OF CENTRAL PULSE PRESSURE AND PULSATILITY IN PATIENTS WITH CAD**

P. Jankowski, A. Bednarek, M. Kloch-Badętek, J. Wiliński, L. Bryniarski, D. Dudek, D. Czarnecka, K. Kawecka-Jaszcz

**To cite this article:** P. Jankowski, A. Bednarek, M. Kloch-Badętek, J. Wiliński, L. Bryniarski, D. Dudek, D. Czarnecka, K. Kawecka-Jaszcz (2011) P7.10: RENAL FUNCTION DOES NOT MODIFY PREDICTIVE VALUE OF CENTRAL PULSE PRESSURE AND PULSATILITY IN PATIENTS WITH CAD, Artery Research 5:4, 181–181, DOI: <https://doi.org/10.1016/j.artres.2011.10.117>

**To link to this article:** <https://doi.org/10.1016/j.artres.2011.10.117>

Published online: 14 December 2019

moving the patients to the high risk group was detection of carotid artery plaques (100%). The contribution of ultrasound parameters to risk stratification has been compared: plaques were detected in 59% patients, IMT >0,9 mm was found in 5%. Moreover, the IMT >0,9 mm occurred only in 7% patients with plaques and only 0,8% without plaques.

**Conclusion:** The assessment of CV risk should include carotid duplex ultrasound studies to reveal plaques. The level of CV risk was affected by carotid plaques presence to substantially greater extent than by IMT.

#### P7.09

##### PULSE WAVE ANALYSIS REVEALS THAT MYOCARDIAL ISCHAEMIA IS NOT LIKELY TO EXPLAIN THE 'J-CURVE' ASSOCIATION BETWEEN DIASTOLIC BLOOD PRESSURE AND MORTALITY

M. G. Schultz<sup>1</sup>, W. P. Abhayaratna<sup>2</sup>, J. E. Sharman<sup>1</sup>

<sup>1</sup>Menzies Research Institute Tasmania, University of Tasmania., Hobart, Australia

<sup>2</sup>Australia National University, Canberra Hospital, Canberra, Australia

**Methods:** The study group consisted of 755 patients (214 women and 545 men; mean age: 57.7±10.0 years) with preserved left ventricular function (EF>40%) undergoing coronary angiography. Demographic and clinical information as well as invasive ascending aortic BP were obtained at baseline. The follow-up was 53.1±18.7 months. The primary end point was: cardiovascular death, myocardial infarction, stroke, cardiac arrest or myocardial revascularization. The Cox proportional hazard regression analysis was used to assess the relation between BP and primary end point.

**Results:** The primary end point occurred in 152 (20.1%) patients whereas CV death, myocardial infarction (MI) or stroke in 79 (10.5%) subjects. Both ascending aortic PP (increase per 10 mmHg: HR 1.12 [95% CI 1.01-1.24]) and pulsatility (increase per 0.1: 1.18 [1.04-1.34]) predicted the risk of primary end point as well as of CV death, MI, or stroke (1.14 [1.00-1.33] and 1.30 [1.10-1.54], resp.). HRs according to the stage of chronic renal disease are presented in the table.

**Conclusion:** Renal function does not modify predictive value of central pulse pressure and pulsatility in patients with CAD.

	GFR<60 ml/min/1.73m <sup>2</sup>	GFR 60-90 ml/min/1.73m <sup>2</sup>	GFR≥90 ml/min/1.73m <sup>2</sup>	P for interaction
Primary end point				
Central PP	1.09	1.19	1.09	NS
Central pulsatility	1.07	1.30	1.12	NS
CV death, MI or stroke				
Central PP	1.11	1.11	1.24	NS
Central pulsatility	1.37	1.32	1.32	NS

Values are hazard ratios for 10 mmHg increase in PP and 0.1 in pulsatility

**Background:** There is a well-established 'J-curve' relationship between brachial DBP and mortality. A purported, although unconfirmed mechanism for the "J-curve" is reduced myocardial perfusion due to low DBP. However, we hypothesised this would be unlikely because DBP may be a poor marker of myocardial perfusion. This study aimed to determine the relationship between DBP and subendocardial perfusion in patients with and without coronary artery disease (CAD).

**Methods:** 134 patients with CAD (aged 76±7 years; 69% male) and 134 matched healthy controls (HC) (aged 77±2 years; 69% male) underwent measurement of brachial DBP and radial tonometry to derive subendocardial viability ratio (SEVR), a marker of subendocardial perfusion. These measures were additionally undertaken in 47 patients (aged 63±10 years) at baseline and during peak dobutamine stress echocardiography in presence or absence of myocardial ischaemia.

**Results:** There was no difference in DBP or SEVR between HC and CAD patients ( $P>0.05$ ), nor was there a difference in SEVR across quartiles of DBP in CAD ( $P=0.07$ ) or HC ( $P=0.14$ ) patients. Associations between DBP and SEVR in HC ( $r=0.185$ ,  $P=0.03$ ) and CAD patients ( $r=0.204$ ,  $P=0.02$ ) were non significant after adjustment for age and height ( $p=0.07$  and  $p=0.11$ , respectively). At peak dobutamine stress, SEVR was significantly reduced in patients with ischaemia versus those without inducible ischaemia (84±17 vs. 101±22 mmHg.s.min<sup>-1</sup>,  $P=0.01$ ). However, DBP was not significantly different (65±14 vs. 67±15 mmHg,  $P=0.32$ ).

**Conclusion:** Brachial DBP is a poor marker of subendocardial perfusion, suggesting the 'J-curve' relationship between DBP and mortality is unlikely attributable to reduced myocardial perfusion.

#### P7.10

##### RENAL FUNCTION DOES NOT MODIFY PREDICTIVE VALUE OF CENTRAL PULSE PRESSURE AND PULSATILITY IN PATIENTS WITH CAD

P. Jankowski<sup>1</sup>, A. Bednarek<sup>1</sup>, M. Kloch-Badelek<sup>1</sup>, J. Wiliński<sup>1</sup>, L. Bryniarski<sup>1</sup>, D. Dudek<sup>2</sup>, D. Czarnecka<sup>1</sup>, K. Kawecka-Jaszcz<sup>1</sup>

<sup>1</sup>Department of Cardiology and Hypertension, Jagiellonian University Medical College, Cracow, Poland

<sup>2</sup>Department of Cardiology, Jagiellonian University Medical College, Cracow, Poland

**Background:** The differences between central and peripheral blood pressure (BP) values have been known for decades. Although the predictive value of central BP in coronary patients with impaired renal function has not been studied so far. Therefore, the aim of the study was to assess the influence of renal function on the predictive value of ascending aortic pulse pressure (PP) and pulsatility (the ratio of PP to mean BP) in patients with coronary artery disease.

#### P7.11

##### NORMAL VALUES OF PULSE WAVE VELOCITY AND AUGMENTATION INDEX AMONG OMANI VOLUNTEERS; PRELIMINARY REPORT

K. M. S. AlHashmi, M. AlHooti, A. H. Al-Ghafri, Z. N. Al-Rashdi, M. O. Hassan  
Department of Physiology, College of Medicine & Health Sciences, Sultan Qaboos University, Muscat, Oman

**Background:** Stiffness of large arteries has been found to be an independent predictor of adverse cardiovascular events in the general population, in patients with essential hypertension, diabetes mellitus and end-stage renal disease. Of the several indices used to reflect arterial stiffness, aortic pulse wave velocity (AoPWV) is considered to be the gold standard. Determining the normal distribution of AoPWV and Alx of a population is important to apply them clinically. This study therefore aims at determining normal values for arterial stiffness indices in normal Omani subjects.

**Method:** Augmentation index (Alx) and aortic pulse wave velocity (AoPWV) were measured using applanation tonometry (SphygmoCor®; Atcor medical) in 43 (23 women and 20 men) healthy Omani volunteers.

**Result:** The mean age of women was 30 ± 9 years and for men was 36 ± 6 years. Reference values were estimated using 97.5 and 2.5 percentiles. The estimated values for Alx corrected for heart rate was -16 to 38 in women and 0 to 26 in men. The AoPWV were 4.6 to 7.1 m/s and 5.2 to 9.6 m/s in women and in men respectively. Men had significantly higher AoPWV compared to women (6.9 ± 0.9 Vs 5.7 ± 0.7,  $P = 0.001$ ) but there was no significant gender difference in the Alx (11.1 ± 12.2 Vs 10.1 ± 8.3,  $P = 0.77$ ).

**Conclusion:** Preliminary data of this study show that men had significantly higher AoPWV than women with no gender difference in the Alx. Recruitment of more subjects is needed to confirm the above findings.

#### P7.12

##### CENTRAL HAEMODYNAMICS COULD EXPLAIN THE INVERSE ASSOCIATION BETWEEN HEIGHT AND CARDIOVASCULAR MORTALITY

J. C. Reeve<sup>1</sup>, W. P. Abhayaratna<sup>2</sup>, J. E. Sharman<sup>1</sup>

<sup>1</sup>Menzies Research Institute of Tasmania, Hobart, Australia

<sup>2</sup>Clinical Trials Unit of the Canberra Hospital, Canberra, Australia

**Introduction:** Mechanisms underlying the inverse relationship between height and cardiovascular mortality are unknown, but could be related to central haemodynamics. This study aimed to determine the relation of height to central and peripheral haemodynamics.

**Methods:** Study population comprised 1161 randomly selected community-dwelling adults (aged 67.7±12.3; 48% male). Brachial BP was recorded by