



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

12.14: ADJUSTMENT FOR BLOOD PRESSURE WHEN MEASURING PULSE WAVE VELOCITY IN NEWLY DIAGNOSED UNTREATED HYPERTENSIVE PATIENTS

C. Strandhave, M. Svensson, K. Holdensen, C.M. Skov, H. Krarup, J.H. Christensen

To cite this article: C. Strandhave, M. Svensson, K. Holdensen, C.M. Skov, H. Krarup, J.H. Christensen (2011) 12.14: ADJUSTMENT FOR BLOOD PRESSURE WHEN MEASURING PULSE WAVE VELOCITY IN NEWLY DIAGNOSED UNTREATED HYPERTENSIVE PATIENTS, Artery Research 5:4, 204–204, DOI: <https://doi.org/10.1016/j.artres.2011.10.197>

To link to this article: <https://doi.org/10.1016/j.artres.2011.10.197>

Published online: 14 December 2019

P12.14
ADJUSTMENT FOR BLOOD PRESSURE WHEN MEASURING PULSE WAVE VELOCITY IN NEWLY DIAGNOSED UNTREATED HYPERTENSIVE PATIENTS

C. Strandhave¹, M. Svensson², K. Holdensen¹, C. M. Skov¹, H. Krarup³, J. H. Christensen¹

¹Dept. of Nephrology, Aalborg Hospital, Aarhus University Hospital, Aalborg, Denmark

²Dept. of Nephrology, Skejby Hospital, Aarhus University Hospital, Aarhus, Denmark

³Dept. of Clinical Biochemistry, Aalborg Hospital, Aarhus University Hospital, Aalborg, Denmark

Pulse wave velocity (PWV) is a valid and well-documented predictor of cardiovascular risk in hypertensive patients. It is essential to adjust for potential confounders, and especially age and blood pressure (BP) modulate the level of PWV. Both systolic BP and mean arterial pressure (MAP) have been used when adjusting in statistical models. However, in recent years MAP has been favoured for BP adjustment.

The aim was to investigate if MAP was superiorly correlated to PWV compared to systolic BP in newly diagnosed untreated hypertension.

We included 126 newly diagnosed untreated hypertensive patients. Twenty-four hour ambulatory BP measurement was performed and PWV was measured using the SphygmoCor device.

The cohort consisted of 61 men and 73 women with an average age of 50 ± 12 years. The mean systolic BP was 146 ± 12 mmHg and the mean MAP was 108 ± 9 mmHg with no significant difference between genders. The median PWV was 8.4 m/s (range 5.3 to 16.8) with no gender difference.

In a linear regression model age was highly positively correlated with PWV ($\beta = .004$, $p < .001$). In an age-adjusted model, MAP was positively correlated with PWV ($\beta = .004$, $p = .02$) but systolic BP showed a higher positive correlation with PWV ($\beta = .003$, $p = .008$).

PWV measurements must be corrected for BP levels as well as age. Adjusting for systolic BP may be superior to adjusting for MAP in newly diagnosed untreated hypertensive patients.

P12.15
ACCURACY OF OSCILLOMETRIC DETERMINATION OF THE ANKLE-BRACHIAL INDEX AS SCREENING METHOD FOR PERIPHERAL ARTERY DISEASE

L. Campens¹, T. De Backer¹, S. Simoens¹, F. Vermassen², M. Coeman¹, M. De Pauw¹, D. De Bacquer¹

¹Department of Cardiology, University Hospital Ghent, Ghent, Belgium

²Department of Thoracic and Vascular Surgery, University Hospital Ghent, Ghent, Belgium

Aim: The ankle brachial index (ABI) is a non-invasive measure for the assessment of peripheral artery disease (PAD). Currently, its use is not sufficiently implemented in general practice, because it is time-consuming. An automated test may facilitate screening for PAD. In this study we assessed the accuracy and reliability of oscillometric ABI determination compared with Doppler.

Method: A total of 99 subjects, including 27 PAD patients, with various cardiovascular risk profiles (hypertension 45,9%; cardiovascular disease 29,6%; diabetes 24,5%; dyslipidemia 38,8%; smoking 21,4%), were recruited between January 2010 and February 2011 at the Department of Cardiology. Doppler with a handheld probe (Hadedo Bidop ES-100V3) and oscillometric measurements with a validated automated blood pressure system (Data-scope Acutorr Plus) were performed in a randomized order. PAD was defined as ABI below 0.9.

Results: The Spearman correlation coefficient between the Doppler and oscillometric method was 0.71 ($P < 0.001$). The sensitivity and specificity for the oscillometric device for detecting PAD were resp. 80% and 97%, the positive and negative predictive values were resp. 89% and 95%. The average time for Doppler and oscillometric ABI determination significantly differed between both methods (resp. 11 minutes (min) 40 seconds (s) and 4min 29s; $p < 0.001$). The mean ABI measured with the oscillometric device was slightly higher than with Doppler (resp. 1.03 ± 0.16 and 1.01 ± 0.19 ; $P < 0.05$ by paired t-test).

Conclusion: These results show that automated ABI determination is an accurate and reliable screening method for the detection of PAD.

P12.16
ATHEROSCLEROTIC PLAQUE EVALUATION BASED ON THE RESULTS OF ULTRASOUND DUPLEX SCANNING (DS), MULTISPECTRAL RESONANCE IMAGING (MRI), MULTISPIRAL COMPUTED TOMOGRAPHY (MSCT) AND

HISTOCHEMICAL ANALYSIS IN ASYMPTOMATIC PATIENTS WITH MARKED CAROTID ATHEROSCLEROSIS

L. R. Tokloueva, T. V. Balakhonova, E. Y. U. Strazden, M. M. Loukyanov, S. Boytsov

Russian Cardiology Research Center, Moscow, Russian Federation

Aim: To evaluate morphological and functional characteristics of an atherosclerotic plaque (ASP) in patients with marked carotid atherosclerosis.

Materials and methods: Forty patients (25 males, age 68.9 ± 6.6 years) with marked carotid atherosclerosis (stenosis $\geq 70\%$, based on ultrasound DS, MSCT and MRI data) were examined before carotid endarterectomy (16 cases with history of cerebral stroke, CS – Group 1; 24 cases without – Group2). All the patients were asymptomatic (no acute cerebrovascular events in the last 6 months). Histochemical examination of the excised carotid ASP was performed in 21 cases (9 cases in Group 1 and 12 cases in Group 2).

Results: Incidence of unstable carotid ASP in groups with and without history of CS was respectively: 50% vs. 23%, $p = 0.02$ (DS data); 25% and 25%, $p > 0.5$ (MRI). Patients with and without history of CS have respectively: ischemic heart disease, IHD (43.5% vs. 68.8% of cases, $p = 0.03$); history of myocardial infarction, MI (25.1% vs. 8.7%; $p = 0.01$). Carotid ASP was unstable in 71.4% of cases ($n = 16$) and stable – in 28.6% ($n = 6$) based on histochemical examination data. As compared to histochemical examination data (conventional "gold standard") sensitivity and specificity were respectively: 67% and 83% for DS; 93% and 67% for MRI.

Conclusions: Asymptomatic patients with marked carotid atherosclerosis have unstable ASP (detected by DS), IHD and history of MI significantly more often in cases with history of CS than without it. Compared with histochemical examination data MRI was the most sensitive method of unstable carotid plaque detection, while DS was the most specific one.

P12.17
SYSTOLIC AMBULATORY BLOOD PRESSURE IS MORE CLOSELY RELATED TO LEFT VENTRICULAR HYPERTROPHY THAN CENTRAL AORTIC AND BRACHIAL BLOOD PRESSURE IN NEVER TREATED HYPERTENSIVE PATIENTS

E. Rodilla Sala¹, J. M. Pascual Izuel^{1,2}, J. A. Costa Muñoz¹, F. Pérez Lahiguera¹, J. Cardona¹, S. Tejero¹

¹Hospital de Sagunto, Puerto de Sagunto, Spain

²Universidad de Valencia, Valencia, Spain

Background: Our purpose was to assess the relationship between left ventricular mass (LVM) and three different ways to measure blood pressure in hypertensive patients.

Methods: Cross-sectional study that included non-diabetic, untreated hypertensive patients. Central aortic pressure (CAP), derived from pulse wave analysis (SphygmoCor-System), office brachial blood pressure (oBBP) and ambulatory 24-hour blood pressure (ABP) were obtained. LVM was determined by standard echocardiography (cut-off values for left ventricular hypertrophy (LVH) were 125 g/m² for men and 110 g/m² for women).

Results: The study population comprised 418 (229 men, with mean age, 49 ± 13 years). Of the participants, 126 (30%) had LVH. Patients with LVH were older [52(13) vs. 47(12) years; $p < 0.001$], had higher values of systolic oBBP [151(18) vs. 143 (16) mm Hg; $p = 0.001$], CAP [141(18) vs. 132 (16) mm Hg; $p = 0.03$], and systolic ABP [139(14) vs. 131 (9) mm Hg; $p = 0.001$].

Stepwise regression analysis corrected by age, gender and presence of metabolic syndrome, revealed that systolic BP ($\beta = 0,0001$, $p < 0,001$), systolic CAP ($\beta = 0,002$, $p < 0,001$), and systolic ABP ($\beta = 0,001$, $p < 0,001$) were all significantly related to LVH.

In a logistic multivariate regression analysis, dividing LV mass in tertiles, systolic CAP (OR: 2,1;95% CI,1,2-3,7; $p = 0,01$), and systolic ABPM-values (OR:3,21;95% CI,1,8-5,7; $p = 0,001$), but not brachial BP, were associated with LVH.

Conclusions: In untreated hypertensives, LVH is closer related to CAP than to office brachial BP, but systolic ABP-values show the best association to the prevalence of LVH.

Key Words: arterial stiffness, central aortic pressure, left ventricular hypertrophy.

P12.18
COMPARISON OF THE CENTRAL PRESSURE MEASURED WITH A BRACHIAL CUFF AND INVASIVELY MEASURED AORTIC PRESSURE

O. Korolkova¹, A. Lowe², J. E. Davies¹, A. D. Hughes¹, K. H. Parker¹, J. H. Siggers¹