



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P1.12: COMPARISON OF ARTERIAL RELAXATION TIME IN NORMOTENSIVE AND HYPERTENSIVE SUBJECTS

L.W. Winchester, C.C. Winchester, N.Y. Chou

To cite this article: L.W. Winchester, C.C. Winchester, N.Y. Chou (2008) P1.12: COMPARISON OF ARTERIAL RELAXATION TIME IN NORMOTENSIVE AND HYPERTENSIVE SUBJECTS, Artery Research 2:3, 95–95, DOI: <https://doi.org/10.1016/j.artres.2008.08.320>

To link to this article: <https://doi.org/10.1016/j.artres.2008.08.320>

Published online: 21 December 2019

What aspect contributed the most to these results (diet, exercise or possession of a personal vascular image) needs further larger early detection study.

doi:10.1016/j.artres.2008.08.318

P1.11

DIURNAL AND GENDER VARIATION OF ARTERIAL STIFFNESS IN YOUNG HEALTHY VOLUNTEERS

É. Komonyi¹, Sz. Lengyel¹, M. Juhász¹, É. Katona¹, Z. Jenei¹, Gy. Paragh¹, B. Fülesdi², D. Páll¹.

¹ First Department of Internal Medicine, Debrecen, Hungary

² Department of Anesthesiology and Intensive care, Debrecen, Hungary

Objective: The augmentation index (Alx) is a measure of arterial stiffness, and is an autonomous determinant independent of other cardiovascular risk factors.

Methods: We examined whether the peripheral and central blood pressure (BP), pulse rate, Alx of healthy volunteers, measured at three different times of the day (8 AM, 12 PM, 5 PM), would show any difference. The measurements were carried out using SphygmoCor device.

Results: 52 healthy volunteers were included into the study. The average age and BMI of the 23 males were 24.4±2.5 years and 24.0±2.9 kg/m², while it was 23.0±1.4 years and 20.5±3.6 kg/m² at the 29 females, respectively. During the three different times of measurement, the actual peripheral and central systolic and diastolic BP and pulse rate did not show any significant difference either for males, or females. In contrast, the Alx of the males was significantly reduced over the course of the day: 8 AM: 13.1±10.2%; 12 PM: 5.3±9.8%; 5PM: 3.4±8.2% (p<0,01). We observed a similar trend in the case of the females as well: 8 AM: 17.0±9.1%; 12 PM: 13.3±10.2%; 5 PM: 11.3±7.5% (p<0,01). The females Alx were higher than that of males (p<0,01).

Conclusion: We conclude that augmentation index shows a considerable diurnal variation and gender difference. There is a gradual decrease from morning values over the course of the day along. Our data suggest to take period of the day at which the measurement was taken and gender into account, while determining physiological and pathological values or interpreting test results.

doi:10.1016/j.artres.2008.08.319

P1.12

COMPARISON OF ARTERIAL RELAXATION TIME IN NORMOTENSIVE AND HYPERTENSIVE SUBJECTS

L.W. Winchester¹, C.C. Winchester², N.Y. Chou¹.

¹ CW Optics, Inc., Yorktown, Virginia, USA

² University of Mary Washington, Fredericksburg, Virginia, USA

Arterial stiffening is emerging as an important determinant of increased systolic blood pressure (SBP) and pulse pressure in the aging population. The relation of the brachial artery was studied in both normotensive and hypertensive volunteers. The test subjects include six healthy, normotensive (SBP/DBP < 120/80 mm Hg) subjects (control group, four in the age range 19 to 23 years, and two in the older subjects, 57 and 60 years) and four subjects (48 to 58 years) with elevated SBP (ranging from 130-168 mm Hg) and who are currently on hypertensive medications (hypertensive group). Temporal measurements of the pressure and volume waveforms were recorded in both hands with one arm at heart level and the other initially at heart level and then raised a distance of 35 cm above heart level. Upon raising the arm, a delay is observed in the pulse measurement of the raised arm relative to the hand at heart level. The delay has previously been shown to decay exponentially with time. For the normotensive subjects, the average values of the pressure and volume relaxation times τ_p and τ_v were 75 s and 41 s for the younger-age group and 86 s and 68 s for the older-age group. The delay times for subjects in the hypertensive group were assumed to approach the baseline asymptotically. The validity of this assumption has been demonstrated in the control group. The average value of τ_p in the hypertensive group was found to be 581 s, significantly larger than the control group.

doi:10.1016/j.artres.2008.08.320

P1.13

ENHANCED VASCULAR REACTIVITY TO COLD PRESSOR TEST IN AFRICAN NORMOTENSIVE SUBJECTS

D. Adamopoulos, W. Ngatsou, D. Lemogoum, B. Biholong, S. Beloka, O. Lheureux, P. Kayembe, J.P. Degaute, P. van de Borne.

Erasmé Hospital, Dept. of Cardiology, ULB, Brussels, Belgium

Background: Cold exposure increases arterial wave reflection to the aorta mainly through adrenergic vasoconstriction. Normotensive blacks exhibit

heightened sympathetic response to cold as compared to Whites controls. We therefore decided to test the hypothesis that cold exposure would elicit a greater increase in arterial wave reflection in normotensive Africans when compared to normotensive Caucasian subjects. In addition we sought to investigate whether this would be accompanied by a higher increase in aortic pulse wave velocity (PWV) and by a more pronounced vasoconstriction of the skin microcirculation.

Methods: 17 young normotensive Africans and 17 age and weight matched Caucasians were recruited for the study. All underwent assessment of PWV and Augmentation Index corrected for heart rate (Alx) at rest, during and after hand immersion in ice water (cold pressor test, [CPT]). Concomitantly, skin microvascular blood flow response to cold was continuously monitored by laser Doppler flowmetry method.

Results: At rest, Africans exhibited higher values of PWV than Caucasians (7.3±0.3 vs 6.4±0.2 m/sec respectively, p=0.04). During CPT the magnitudes of increases in SBP and PWV were greater in Africans than Caucasians (26±3 mmHg vs 15±3 mmHg and 0.8±0.2 m/sec vs 0.4±0.2 m/sec respectively, p<0.05 for all). Additionally, CPT induced a more pronounced skin microvascular vasoconstriction in Africans as compared to Caucasians (-45±7 % vs -25±7 %, p=0.01). Finally, Africans exhibited higher Alx values during CPT (12.6±2.4 vs 5.5±2.4 %, p=0.04) when compared to Caucasian subjects.

Conclusions: Normotensive Africans exhibit intensified wave reflection to the aorta as response to CPT when compared to Caucasians. This is accompanied by a greater increase in PWV and a more pronounced vasoconstriction of the microcirculation.

doi:10.1016/j.artres.2008.08.321

P1.14

CORRELATION BETWEEN AORTIC PULSE WAVE VELOCITY AND ASYMPTOMATIC CAROTID ATHEROSCLEROSIS

R. Bocskei¹, B. Benczur², F. Molnar³, A. Cziraki¹, M. Illyes¹.

¹ University of Pecs Heart Institute and 1st Department of Internal Medicine, Pecs, Hungary

² Hetenyi Geza County Hospital, Dept. of Cardiology, Szolnok, Hungary

³ TensioMed Arterial Stiffness Centre, Budapest, Hungary

Background: It seems that traditional risk factors are not sufficient predictors of the atherosclerosis and cardiovascular disease (CVD). Although the carotid ultrasonography is a gold standard of the detection of asymptomatic atherosclerosis, the correlation between aortic pulse wave velocity (PWVao) and the presence of preclinical carotid plaques was poorly investigated. The aim of this study was to examine this correlation and determine the most sensitive cut-off value of the aortic PWV for carotid atherosclerosis by using ROC (Receiving Operating Characteristic) analysis.

Methods: 557 asymptomatic subjects were included without known CVD. Arterial stiffness parameters (PWVao) were measured with non-invasive oscillometric device (Arteriograph) and carotid scan was performed with carotid ultrasonography in both side by a "blinded" investigator who was unaware of the stiffness-parameters in all subjects as well.

Results: We have found 283 carotid negative subjects and 274 carotid positive subjects. The carotid ultrasonography was positive when the IMT > 1.3mm or calcificated plaque ≥ 1mm was detected. The mean values were: age 57, BP 134/82 mmHg, HR 72/min, Aix -8,7%, PWVao 9,9 m/s. In data analysis the SPSS software and the ROC curve were used. The most sensitive PWVao was 9.62m/s. The sensitivity of increased PWVao was 77.7%, the specificity 65.4%, the positive predictive value (PPV) 68,5%, the negative predictive value (NPV) 75,2%. The significance between these values was very good.

Conclusion: The increased (>9,62 m/s) PWVao shows a strong association with the asymptomatic carotid plaques and seems to be a suitable method to detect preclinical atherosclerosis.

doi:10.1016/j.artres.2008.08.322

P1.15

ASSOCIATION BETWEEN CAROTID AND FEMORAL ATHEROSCLEROTIC BURDEN AND VASCULOGENIC ERECTILE DYSFUNCTION

A. Tsokanis, C. Vlachopoulos, N. Ioakeimidis, K. Aznaouridis, K. Baou, D. Terentes-Printzios, K. Rokkas, C. Stefanadis.

1st Department of Cardiology, Athens Medical School, Athens, Greece

Background: Vasculogenic erectile dysfunction (ED) may be considered a clinical manifestation of a generalized arterial disease. We attempted to evaluate the association between penile vascular dysfunction and extent of