



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P2.52: ASSOCIATION BETWEEN PWV AND DIFFERENT DEFINITIONS OF THE METABOLIC SYNDROME

S. Holewijn, M. den Heijer, A.F.H. Stalenhoef, J. de Graaf

To cite this article: S. Holewijn, M. den Heijer, A.F.H. Stalenhoef, J. de Graaf (2008) P2.52: ASSOCIATION BETWEEN PWV AND DIFFERENT DEFINITIONS OF THE METABOLIC SYNDROME, Artery Research 2:3, 119–119, DOI: <https://doi.org/10.1016/j.artres.2008.08.418>

To link to this article: <https://doi.org/10.1016/j.artres.2008.08.418>

Published online: 21 December 2019

Parameter	C (n= 125)	MS (n= 138)	DBT (n= 29)	p
Age	54 ± 12	54 ± 11	51 ± 10	NS
Sex (% males)	70	64	52	NS
SBP (mmHg)	141 ± 18	141 ± 16	142 ± 10	NS
DBP (mmHg)	85 ± 11	86 ± 9	87 ± 10	NS
HR (bpm)	69 ± 9	71 ± 10	74 ± 10	NS
% abnormal IMT	19	33	41	.012
% Plaques	22	54	55	< .001
%Abnormal FMD	33	39	45	NS
%Abnormal PWV	30	38	65	.002
FS (mean)	10 ± 8	10 ± 6	25 ± 12	< .001
VS (mean)	2,1 ± 1,4	2,5 ± 1,4	3,0 ± 1,7	.004

Conclusions: 1- In a selected population of matched p. according to age, sex and BP, there seems to be an increase in the degree of VD when MS is present. 2- The pattern of VD in MS p. was close to that in DM p. consistent with the described increase in CV risk in MS. 3-A VS may be useful in the clinical practice.

doi:10.1016/j.artres.2008.08.417

P2.52 ASSOCIATION BETWEEN PWV AND DIFFERENT DEFINITIONS OF THE METABOLIC SYNDROME

S. Holeyijn, M. den Heijer, A.F.H. Stalenhoef, J. de Graaf.
Radboud University Nijmegen Medical Centre, Nijmegen, Netherlands

Objectives: To explore 1. the presence of arterial stiffness, as measured with pulse wave velocity (PWV) in subjects with the Metabolic Syndrome (MetS) and 2. the contribution of PWV and the MetS in the prediction of prevalent CVD in a 50-70-year-old Dutch population-based cohort.

Background: Inconsistent results have been published about 1. the relations between arterial stiffness and the MetS and 2. the role of both in the prediction of CVD. Most studies have been performed in selected high-risk populations.

Method: We measured arterial stiffness by PWV using tonometry in 1510 participants of the Nijmegen Biomedical Study and determined its associations with different definitions of the MetS (according to the NCEP, IDF and WHO). Multiple regression analysis was performed to determine the role of PWV and MetS in CVD risk prediction.

Results: Both men and women with the MetS had a significantly higher PWV than those without the MetS, independent of the definition used (for IDF: men: +MetS:9.5(2.3)vs -MetS11.1(3.1)m/s, women: +MetS:9.1(2)vs -MetS 10.9(3)m/s). Age and gender were the main contributors to PWV, explaining 18.7% of the variance in PWV. Adding any definition of the MetS to the model increased the percentage explained variance of PWV to only 24% whereas adding all traditional CV risk factors predicted 43.1%. In regression analysis of prevalent CVD, the CV risk factors predicted 26.3% and adding PWV or the MetS did not improve the prediction of prevalent CVD.

Conclusion: Our results question the additive value of 1. using the definition of the MetS in the prediction of arterial stiffness and 2. measuring PWV and using the MetS in the prediction of prevalent CVD. Prospective studies have to determine the additive contribution of PWV and the MetS in CVD risk-prediction in general population.

doi:10.1016/j.artres.2008.08.418

P2.54 INFLUENCE OF CENTRAL OBESITY ON EARLY CAROTID INTIMA-MEDIA THICKENING IS INDEPENDENT OF THAT FROM OTHER RISK FACTORS

V.M.G. Maher¹, M.O. Dowd², M. Carey¹, C. Markham¹, A. Byrne², E. Hand¹, D. McLnerney²

¹ Dept of Cardiology, Adelaide and Meath Hospital, Tallaght, Dublin, Ireland
² Dept of Radiology, Adelaide Meath Hospital, Tallaght, Dublin, Ireland

Its unknown if obesity or its associated risk factors influence early vascular change.

Methods: We investigated if anthropometric measurements, body mass index [BMI], waist/hip ratio [WHR], waist circumference [Wc] and Waist/height ratio WHTR, or metabolic parameters (glucose, insulin, lipid, uric acid and blood pressure {BP}) correlated more with Carotid intima-media thickness (IMT), vascular stiffness [Augmentation Index] and brachial artery reactivity). 100 Subjects (71F, 29M) without vascular events, BP <140/90, LDL < 4 mmol/l, glucose < 6.2 mmol/l participated.

Results: BMI, WHR, WC, WHTR correlated significantly with triglyceride, HDL, LDL, insulin, glucose, uric acid and BP levels (p<0.001). IMT correlated with WHTR, BMI, WC, Glucose (p<0.001), Homeostasis Index (HOMA) and Cholesterol levels (p<0.05). Only Age, WHTR or BMI were significant correlates of IMT in a multivariate analysis (MVA) (p<0.01) including WHTR or BMI, with age, sex, SBP, HDLc and HOMA. Augmentation Index correlated with age (p<0.0001), WHTR and WC (p<0.0005) but with age only in a MVA. Vascular reactivity did not correlate with any anthropometric or metabolic parameters. Anthropometric cut off points, (BMI <25, WC <102cm M, <88cm F, WHR <0.9 M, <0.8 F, and WHTR < 0.5) significantly differentiated normal from abnormal metabolic and vascular measurements. The WHTR ratio < 0.5 was as reliable as the BMI cut-off < 25 in determining metabolic and vascular abnormalities. BMI and WHTR were strongly associated with 89% agreement (p<0.0001). These results demonstrated that anthropometric and metabolic parameters correlated, but anthropometric parameters were the significant correlates of vascular change. A waist/height ratio > 0.5 predicts both early vascular and metabolic changes. These data support a risk factor independent vasculotrophic effect of obesity.

doi:10.1016/j.artres.2008.08.423

P2.55 CHANGES IN TRUNK FAT MASS AND PERIPHERAL LEAN MASS ARE ASSOCIATED WITH CHANGES IN CAROTID ARTERIAL STIFFNESS IN A HEALTHY POPULATION – A 6-YEAR FOLLOW-UP STUDY

F. Schouten¹, I. Ferreira³, M.R. de Boer¹, C.D.A. Stehouwer³, Y.M. Smulders², J.W. Twisk¹

¹ VU University, Amsterdam, Netherlands

² VU University Medical Centre, Amsterdam, Netherlands

³ University Hospital Maastricht, Maastricht, Netherlands

Introduction: In cross-sectional studies, total body fat mass and particularly central fat distribution are associated with greater arterial stiffness. How changes in body fat and body fat distribution impact on changes in large artery stiffness is unknown.

Methods: Data were derived from the Amsterdam Growth and Health Longitudinal Study (n=268; 126 male and 142 female). At age 36, body fat distribution was determined by dual-energy x-ray absorptiometry and arterial (carotid and femoral) properties were measured using ultrasound imaging. These measurements were repeated 6 years later. Data were analysed with multiple linear regression analyses.

Results: Increases in total body fat mass (standardized β : -0.14, 95%CI: -0.37; 0.08) and especially trunk fat mass (β : -0.25, 95%CI: -0.49; -0.02) were associated with decreases in the carotid distensibility coefficient. In contrast, increases in peripheral lean mass were inversely associated with carotid Young's elastic modulus (β : -0.27, 95%CI: -0.49; -0.05). These associations were independent of changes in other body composition compartments or riskfactors (i.e. gender, mean arterial pressure, total/HDL cholesterol ratio, triglycerides, glycated haemoglobin, heart rate). Changes in body fat distribution were not associated with changes in femoral artery stiffness.

Conclusion: Increases in trunk fat mass during adulthood have adverse effects on carotid stiffness, whereas increases in peripheral lean mass may counteract this.

doi:10.1016/j.artres.2008.08.419

P2.56 EFFECTS OF LEAN AND FAT MASS ON BONE MINERAL DENSITY AND ARTERIAL STIFFNESS IN ELDERLY MEN

A. Kearney-Schwartz², A. Zervoudaki¹, P. Salvi¹, C. Labat⁴, G. Weryha³, A. Benetos¹

¹ Department of Geriatrics, University Hospital, Nancy, France

² Clinical Investigation Center, University Hospital, Nancy, France

³ Department of Endocrinology, University Hospital, Nancy, France

⁴ INSERM U684, University of Nancy, Nancy, France

Introduction: The aim of this study was to evaluate the influence of fat and lean mass on both arterial stiffness and bone mass density (BMD) in elderly men.

Methods: This study was performed in 169 French males over 60 years. Aortic stiffness was assessed by carotid/femoral pulse wave velocity (PWV) using a validated automated device (PulsePen®). BMD and body composition were determined with a dual-energy X-ray absorptiometry (DEXA) device in lumbar spine L1-L4, femoral neck and total body.