



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P1.06: THE ARTERIOGRAPH: CORRELATED TO AORTIC STIFFNESS, BUT MEASURING AXILLO-BRACHIAL ARTERY STIFFNESS?

B. Trachet, P. Reymond, J. Kips, A. Swillens, M.L. De Buyzere, B. Suys, N. Stergiopulos, P. Segers

To cite this article: B. Trachet, P. Reymond, J. Kips, A. Swillens, M.L. De Buyzere, B. Suys, N. Stergiopulos, P. Segers (2010) P1.06: THE ARTERIOGRAPH: CORRELATED TO AORTIC STIFFNESS, BUT MEASURING AXILLO-BRACHIAL ARTERY STIFFNESS?, Artery Research 4:4, 154–154, DOI: <https://doi.org/10.1016/j.artres.2010.10.011>

To link to this article: <https://doi.org/10.1016/j.artres.2010.10.011>

Published online: 21 December 2019

high resolution, the near wall thickness (NW-IMT) is still challenging. New ultrasound software (CAAS-US, Pie medical Imaging) was developed for NW-FW-IMT measurement from radiofrequency signal.

Objectives: To compare CCA IMT between CAAS-US and standard Artlab, and to assess CAAS-US determination of NW-IMT and both near and far wall IMT variations during the cardiac cycle, in 38 treated hypertensive (HT) and 18 normotensive (NT) patients.

Methods: Diastolic outer diameter (Dd) and FW-IMT were determined with CAAS-US and Artlab. Bland-Altman test and Pearson's correlation coefficient (R) were used to compare methodologies. With CAAS-US, NW-IMT and FW-IMT were compared. Systolo(s)-diastolic(d) variations of IMT and outer D were calculated: $\Delta\text{IMT} ((\text{IMTs} - \text{IMTd})/\text{IMTd})$ of NW and FW were compared to $\Delta\text{D} ((\text{Ds} - \text{Dd})/\text{Dd})$.

Results: No significant differences in CCA Dd and FW-IMTd were observed between two methods ($R=0.97$, $\text{RMSE}=0.124$ mm and $R=0.91$, $\text{RMSE}=0.065$ mm, respectively $p<0.0001$). IMTd between near and far wall are significantly correlated ($R=0.73$, $\text{RMSE}=0.117$ mm, $p<0.0001$). FW ΔIMT is significantly correlated to ΔD ($R=0.57$, $\text{RMSE}=0.036$ mm, $p=0.0002$) whereas NW ΔIMT is not ($R=0.30$, $\text{RMSE}=0.073$ mm, $p=0.065$).

Conclusion: CAAS-US allows determination of both near and far wall CCA IMT. Whereas FW-IMT variation during the cardiac cycle is measurable, NW-IMT is less precise and remains a challenging. These findings added to CAAS-US plaque contours ability may be useful to estimate more completely the real state of the carotid artery atherosclerotic process.

P1.05

COMPARING AORTIC PULSE WAVE VELOCITY BY MAGNETIC RESONANCE IMAGING AND THE NEW OSCILLOMETRIC METHOD ARTERIOGRAPH

M. Rezai¹, N. Sherratt², J. K. Cruickshank¹

¹Cardiovascular Research Group, University of Manchester, Manchester, United Kingdom

²Wellcome Trust Clinical Research Facility, Manchester, United Kingdom

Aims: Comparing aortic pulse wave transit time (TT) and velocity (aPWV) between magnetic resonance imaging (MR) and the oscillometric method Arteriograph (AG) that has potential for outpatient use.

Methods: MR phase-contrast transverse slices were sequentially taken at aortic arch (pulmonary bifurcation level), and 2 cm above aortic bifurcation in 49 men (age: 53 ± 6 yr). TT was calculated using 10% of the MR flow wave amplitudes as wave-feet. The aortic root to bifurcation (aoLength) was measured in 3D volumes rendered from serial sagittal slices. Supine left-arm AG measurements were made post-MR, estimating aoLength from suprasternal-notch to symphysis-pubis surface length. Results are mean differences (95%CI).

Results: TT_{MR} and aPWV_{MR} covered a mean 85% of aoLength partly omitting the proximal ascending aorta. TT_{AG} (71 ± 10 ms) was $6.6(3.8-9.3)$ ms (10%) higher than TT_{MR} (64 ± 10 ms) and correlated ($r=0.54$, $p<0.001$). aPWV_{AG} (7.9 ± 1.4) was $1.3(0.9-1.7)$ m/s higher than aPWV_{MR} (6.6 ± 1.3) and correlated ($r=0.50$, $p<0.001$). AG's sternum-pubis length was $70(59-81)$ mm higher than MR's aoLength ($r=0.55$, $p<0.001$).

A regression model was derived from 29 cases predicting measured aoLength_{MR} using age and height. When tested in the remaining 20 cases, the predicted aoLength_{MR} was within $5.3(-22-11)$ mm of that measured by MR. Compared with original aPWV_{AG}, recalculated aPWV_{AG} using TT_{AG} and the regression-predicted aoLength was less different ($0.31(0.01-0.61)$ m/s - $p=0.02$), and more closely correlated with aPWV_{MR} ($r=0.62$).

Conclusions: TT estimations by AG and MR are close, given omitted proximal lengths by MR. More accurate length estimation can significantly improve AG's aPWV measurement using MR as a reference.

P1.06

THE ARTERIOGRAPH: CORRELATED TO AORTIC STIFFNESS, BUT MEASURING AXILLO-BRACHIAL ARTERY STIFFNESS?

B. Trachet¹, P. Reymond⁴, J. Kips¹, A. Swillens¹, M. L. De Buyzere², B. Suys³, N. Stergiopoulos⁴, P. Segers¹

¹bioMeda - IBiTech, Ghent University, Ghent, Belgium

²Department of Cardiovascular Diseases, Ghent University Hospital, Ghent, Belgium

³Department of Pediatrics, Antwerp University, Antwerp, Belgium

⁴Laboratory of Hemodynamics and Cardiovascular Technology, Swiss Federal Institute of Technology, Lausanne, Switzerland

The Arteriograph (Tensiomed) is a device that determines aortic pulse wave velocity (PWV) by recording a brachial blood pressure waveform during supra-systolic inflation of a brachial cuff. We validated the working principle of the Arteriograph in a validated 1D computer model of the arterial tree, and found an excellent correlation between Arteriograph (PWV_{ATG}) and

carotid-femoral (PWV_{car-fem}) PWV ($r = 0.98$, $p<0.01$) when homogeneously altering stiffness parameters over the complete arterial network. However, selectively altering the stiffness of the aortic or axillo-brachial pathway demonstrated that PWV_{ATG} is determined solely by axillo-brachial stiffness and not by aortic stiffness. Furthermore, wave intensity analysis shows that the secondary forward compression wave picked up by the Arteriograph and used to assess the travel time of the pressure wave over the aorta is not caused by a reflection from the lower body. Instead, this wave is the result of "trapping" of the initial forward compression wave between the occluding cuff and the axillo-aortic junction. Thus the Arteriograph measures axillo-brachial stiffness, and the good correlation between PWV_{ATG} and PWV_{car-fem} is driven by the fact that axillo-brachial and central stiffness were changed to the same extent in the model. Combining these results with earlier findings in literature of good in vivo correlations between PWV_{ATG} and PWV_{car-fem}, axillo-brachial and aortic stiffness are likely to be related. However, this does not necessarily imply that axillo-brachial and aortic segments change similarly with age (or disease). We conclude that PWV_{ATG} is, at best, an indirect and unspecific estimate of aortic stiffness.

P1.07

ULTRASOUND EVALUATION OF CAROTID STIFFNESS IN HEALTHY VOLUNTEERS DURING EXERCISE: A PILOT STUDY

E. Bianchini¹, R. M. Bruno², A. I. Corciu¹, V. Gemignani¹, F. Faita¹, M. Demi¹, L. Ghiadoni², R. Sicari¹

¹Institute of Clinical Physiology-CNR, Pisa, Italy

²Department of Internal Medicine, University of Pisa, Pisa, Italy

The study of cardiovascular parameters in exercise is intriguing, since this evaluation could provide information about dynamic conditions, mimicking patient's real life. Therefore, we evaluated carotid cross-sectional distensibility coefficient (DC), systemic vascular resistance corrected by cardiac frequency (SVRI_f), arterial elastance (Ea) and left-ventricular elastance (Elv) during graded bicycle semi-supine exercise test.

In 18 healthy subjects (9 men, 34 ± 3 years) cardiac volumes were estimated from 2D transthoracic echocardiography, right carotid diameter and distension by an automatic system (Carotid Studio, IFC-CNR) applied to ultrasound B-mode image sequences and central pressures by tonometry. All measurements were performed at 60%, 70%, 80% and 85% (peak) of the age-dependent maximal heart rate.

During exercise DC decreased (peak versus rest: -17.8% and $p<0.05$), and SVRI_f did not significantly change. Ea increased (+21.3%, $p<0.01$) and, since Elv presented a greater variation (+69.2%, $p<0.001$), arterial ventricular coupling (Ea/Elv) decreased (-22.6%, $p<0.05$). As expected, central pulse pressure was significantly increased (+81.8%, $p<0.01$).

In conclusion, carotid stiffness increased during exercise, possibly due to the recruitment of more collagen fibers and the consequent different mechanical behavior of arterial walls at higher pressures. Since SVRI_f did not change significantly, the increased arterial stiffness, observed at carotid site, might represent the main determinant of Ea variation. Finally, a decrease in arterial-ventricular coupling during exercise was confirmed.

Our results show the feasibility of a simultaneous multi-sites approach that could help in the understanding of arterial physiology and patho-physiology in stress conditions.

P1.08

THE BRACHIO-TO-RADIAL PULSE PRESSURE AMPLIFICATION AND ITS CONTRIBUTION TO CENTRAL-TO-PERIPHERAL PULSE PRESSURE AMPLIFICATION

G. Pucci^{1,2}, J. Cheriyan², L. Whittaker², S. S. Hickson², G. Schillaci¹, C. M. McEnery², I. B. Wilkinson²

¹Clinical and Experimental Medicine, University of Perugia, Perugia, Italy

²Clinical Pharmacology Unit, University of Cambridge, Cambridge, United Kingdom

Pulse pressure amplification (PPa) is the phenomenon by which systolic blood pressure (SBP) increases from aorta to periphery, whether mean (MAP) and diastolic pressure (DBP) remain constant along the arterial tree. Controversy exist in defining the extent of brachio-radial PPa and how much it contributes central-to-peripheral PPa. The SphygmoCor estimates central PP (cPP) by a transfer function applied to the radial waveform calibrated to brachial SBP/DBP, or to brachial MAP/DBP. If brachio-radial PPa was irrelevant, calibrating radial waveform to bSBP/bDBP or to bMAP/bDBP would have no measurable impact on estimated cPP.

In order to assess brachio-radial PPa and its determinants, 93 healthy subjects (42 ± 20 years, 53% men) were studied after 10 minutes of supine rest.