



## Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantispress.com/journals/artres>

---

### **3.3: ROLE OF ARTERIAL STIFFNESS AND BLOOD PRESSURE VARIABILITY IN THE DEFINITION OF SHATS (SYSTEMIC HEMODYNAMIC ATHEROTHROMBOTIC SYNDROME)**

Angelo Scuteri, Valentina Rovella, Danilo Alunni Fegatelli, Manfredi Tesauro, Marco Gabriele, Nicola Di Daniele

**To cite this article:** Angelo Scuteri, Valentina Rovella, Danilo Alunni Fegatelli, Manfredi Tesauro, Marco Gabriele, Nicola Di Daniele (2018) 3.3: ROLE OF ARTERIAL STIFFNESS AND BLOOD PRESSURE VARIABILITY IN THE DEFINITION OF SHATS (SYSTEMIC HEMODYNAMIC ATHEROTHROMBOTIC SYNDROME), Artery Research 24:C, 73–73, DOI: <https://doi.org/10.1016/j.artres.2018.10.034>

**To link to this article:** <https://doi.org/10.1016/j.artres.2018.10.034>

Published online: 7 December 2019

conventional risk factors and baseline renal function). These associations remained unchanged after further adjustments for central artery stiffness or traditional central haemodynamic parameters. No other RPA-parameters exhibited significant associations.

**Conclusions:** These findings demonstrate that baseline INTPR is independently associated with the eGFR reduction in older adults, suggesting that INTPR may play a role in the functional decline of the kidneys.

### 3.3

#### ROLE OF ARTERIAL STIFFNESS AND BLOOD PRESSURE VARIABILITY IN THE DEFINITION OF SHATS (SYSTEMIC HEMODYNAMIC ATHEROTHROMBOTIC SYNDROME)

Angelo Scuteri<sup>1</sup>, Valentina Rovella<sup>2</sup>, Danilo Alunni Fegatelli<sup>3</sup>, Manfredi Tesaro<sup>4</sup>, Marco Gabriele<sup>2</sup>, Nicola Di Daniele<sup>4</sup>

<sup>1</sup>Medicine Director School of Geriatrics University of Sassari University Hospital, Sassari, Italy

<sup>2</sup>Policlinico Tor Vergata, Rome, Italy

<sup>3</sup>University of Rome La Sapienza, Rome, Italy

<sup>4</sup>University of Rome Tor Vergata, Rome, Italy

**Background:** CV risk exponentially increases as the number of damaged organs increases. The Systemic Hemodynamic Atherosclerotic Syndrome (SHATS) represents a novel conceptualization of the CV continuum focusing on simultaneous multi-organ alteration. This is the first study operationally defining SHATS and aimed at identifying its determinants.

**Methods:** Left Ventricular Hypertrophy (echocardiography), Common Carotid Artery plaque and increased thickness (ultrasound), and Chronic Kidney Disease (estimated Glomerular Filtration Rate) indexed selective target organ damage. SHATS was operationally defined as their simultaneous presence in a patient. PWV was measured by Sphygmocor<sup>®</sup> and BP variability by 24 h ABPM.

**Results:** SHATS affected 19.9% of the 367 studied subjects. Subjects with SHATS had a similar prevalence in diabetes mellitus, but a greater prevalence of very stiff artery (84.9 vs 64.3 %,  $p < 0.01$ ) and use of antihypertensive medications. In the presence of similar office BP, SHATS was associated with higher 24 h SBP and lower 24 h DBP (a greater pulsatile pressure!), reduced nighttime SBP fall, and a twofold greater prevalence of reverse dipper status (48.2 vs 20.2 %,  $p < 0.001$ ). BMI (positive correlation) and DBP (negative correlation) were the only traditional CV risk factors significantly associated with the odds of having SHATS. Very stiff artery and BP variability were significant independent determinants of SHATS, with highly predictive accuracy.

**Conclusion:** SHATS, the simultaneous damage of multiple target organs, may easily operationally be defined. Very stiff artery and BP variability represent key factors for SHATS. The present results support the hypothesis of SHATS as a systemic condition, needing further characterization.

### 3.4

#### A CLINICAL SCORE TO PREDICT ELEVATED ARTERIAL STIFFNESS: DERIVATION AND VALIDATION IN 3,943 HYPERTENSIVE PATIENTS

Panagiotis Xaplanteris<sup>1</sup>, Charalambos Vlachopoulos<sup>2</sup>, Athanasios Protogerou<sup>3</sup>, Konstantinos Aznaouridis<sup>2</sup>, Dimitris Terentes-Printzios<sup>2</sup>, Antonis Argyris<sup>3</sup>, Nikolaos Tentolouris<sup>3</sup>, Petros Sfikakis<sup>3</sup>, Dimitris Tousoulis<sup>2</sup>

<sup>1</sup>CHU Saint Pierre, Brussels, Belgium

<sup>2</sup>Hippokraton Hospital, University of Athens, 1st Department of Cardiology, Athens, Greece

<sup>3</sup>Laiko University General Hospital, Cardiovascular Prevention and Research Unit, Department of Pathophysiology, Athens, Greece


**Purpose/Background/Objectives:** Aortic stiffness assessed by carotid-femoral pulse wave velocity (PWV) is an important predictor to gauge the overall risk of hypertensive patients; nonetheless, it is underutilized in everyday practice. We derived a simple scoring system based on clinical variables that can identify patients with a priority for measurement of PWV, i.e. those with elevated PWV ( $\geq 10$  m/sec) and at higher risk for events.

**Methods:** Patient data from three outpatient clinics ( $n = 3,943$ ) were used to form a derivation, internal and external validation cohort. For derivation, independent predictors of high PWV from a binary logistic regression model were dichotomized and implemented in a clinical prediction scoring system with the acronym SAGE (office systolic blood pressure  $\geq 160$  mmHg: 4 points,

age  $\geq 60$  years: 4 points, glycemia [blood glucose  $\geq 126$  mg/dl]: 1 point, eGFR  $\leq 60$ : 2 points).

**Results:** Its performance was validated at the internal and external validation cohorts with c-statistics being 0.781 (95% CI: 0.753–0.808) and 0.718 (95% CI: 0.682–0.755) respectively (Figure 1). A cut-off of 5 points to identify patients with high PWV in the external validation cohort yielded a positive predictive value, negative predictive value, sensitivity and specificity of 60.7%, 84.8%, 51.9% and 78.3% respectively.

**Conclusions:** The SAGE score that takes into account easily measured clinical variables (systolic blood pressure, age, glucose and eGFR) can be used to predict elevated levels of PWV and prioritize its measurement in specific patients. Its use will result in greater acknowledgement of the role of aortic stiffness and aid physicians in implementing it in clinical practice.



### References

- Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol* 2010; 55 (13):1318-1327.
- Ben-Shlomo Y, Spears M, Boustred C, May M, Anderson SG, Benjamin EJ, et al. Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects. *J Am Coll Cardiol* 2014; 63 (7):636-646.
- Vlachopoulos C, Xaplanteris P, Aboyans V, Brodmann M, Cifkova R, Cosentino F, et al. The role of vascular biomarkers for primary and secondary prevention. A position paper from the European Society of Cardiology Working Group on peripheral circulation: Endorsed by the Association for Research into Arterial Structure and Physiology (ARTERY) Society. *Atherosclerosis* 2015; 241 (2):507-532.
- Reference Values for Arterial Stiffness C. Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: 'establishing normal and reference values'. *Eur Heart J* 2010; 31 (19):2338-2350.
- McEnery CM, Spratt M, Munnery M, Yarnell J, Lowe GD, Rumley A, et al. An analysis of prospective risk factors for aortic stiffness in men: 20-year follow-up from the Caerphilly prospective study. *Hypertension* 2010; 56 (1):36-43.

### 3.5

#### CUFF BLOOD PRESSURE IS PROGRESSIVELY MORE BIASED WITH INCREASING AGE: INDIVIDUAL PARTICIPANT LEVEL ANALYSIS FROM THE INSPECT CONSORTIUM

Dean Picone<sup>1</sup>, Martin Schultz<sup>2</sup>, Petr Otahal<sup>3</sup>, Ahmed Al-Jumaily<sup>4</sup>, J. Andrew Black<sup>2</sup>, Willem Bos<sup>5</sup>, Chen-Huan Chen<sup>6</sup>, Hao-Min Chen<sup>7</sup>, Antoine Cremer<sup>8</sup>, Nathan Dwyer<sup>2</sup>, Ricardo Fonseca Diaz<sup>2</sup>, Brian Gould<sup>9</sup>, Alun Hughes<sup>10</sup>, Hack-Lyoung Kim<sup>11</sup>, Peter Lacy<sup>12</sup>, Esben Laugesen<sup>13</sup>, Sandy Muecke<sup>14</sup>, Nobuyuki Ohte<sup>15</sup>, Stefano Omboni<sup>16</sup>, Christian Ott<sup>17</sup>, Xiaoqiong Peng<sup>2</sup>, Telmo Pereira<sup>18</sup>, Giacomo Pucci<sup>19</sup>, Philip Roberts-Thomson<sup>2</sup>, Niklas Rossen<sup>13</sup>, Roland Schmieder<sup>17</sup>, Velandai Srikanth<sup>2</sup>,