



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P122: CALCULATING RESERVOIR PRESSURE WITH OR WITHOUT FLOW INFORMATION: SIMILARITY AND ALGORITHMIC SENSITIVITY AT RADIAL ARTERY

Michael Ebner, Kim Parker, Tom Vercauteren, Sébastien Ourselin, Siegfried Wassertheurer, Alun Hughes, Bernhard Hametner

To cite this article: Michael Ebner, Kim Parker, Tom Vercauteren, Sébastien Ourselin, Siegfried Wassertheurer, Alun Hughes, Bernhard Hametner (2017) P122: CALCULATING RESERVOIR PRESSURE WITH OR WITHOUT FLOW INFORMATION: SIMILARITY AND ALGORITHMIC SENSITIVITY AT RADIAL ARTERY, Artery Research 20:C, 78–79, DOI: <https://doi.org/10.1016/j.artres.2017.10.104>

To link to this article: <https://doi.org/10.1016/j.artres.2017.10.104>

Published online: 7 December 2019

hypertension, with investigators, however, not reaching a consensus on the relative importance of each wave component (1,2).

Objective: The aim of the current investigation was to examine the wave profile over time after developing an age-adapted, mathematical, one-dimensional model of the cardiovascular system.

Methods: Our state-of-the-art 1-D model (3,4) was extended to include turbulence and inertial effects of the flow exiting the left ventricle. Literature data on the age-associated changes in arterial stiffness, peripheral resistance and cardiac contractility were gathered and used as an input for the simulation.

Results: The predicted evolution of pressure and augmentation index with age followed accurately the curves obtained in a number of large-scale clinical studies. Analysis of the relative contribution of the forward and backward wave components showed that the forward wave becomes the major determinant of the increase in central and peripheral SBP and PP with advancing age.

Conclusions: The 1-D model of the ageing tree and heart captures faithfully and with great accuracy the central pressure evolution with ageing. The stiffening of the proximal aorta and the resulting augmentation of the forward pressure wave is the major contributor of the systolic pressure augmentation with age.

References

1. O'Rourke MF, Nichols WW. Changes in Wave Reflection With Advancing Age in Normal Subjects. *Hypertension*. 2004 Dec 1;44(6):E10–1.
2. Mitchell GF, Conlin PR, Dunlap ME, Lacourcière Y, Arnold JMO, Ogilvie RI, et al. Aortic diameter, wall stiffness, and wave reflection in systolic hypertension. *Hypertens Dallas Tex* 1979. 2008 Jan;51(1):105–11.
3. Reymond P, Bohraus Y, Perren F, Lazeyras F, Stergiopoulos N. Validation of a patient-specific one-dimensional model of the systemic arterial tree. *Am J Physiol Heart Circ Physiol*. 2011 Sep; 301(3):H1173–1182.
4. Reymond P, Merenda F, Perren F, Rüfenacht D, Stergiopoulos N. Validation of a one-dimensional model of the systemic arterial tree. *Am J Physiol – Heart Circ Physiol*. 2009 Jul 1;297(1):H208–22.

P121

IDENTIFYING HAEMODYNAMIC DETERMINANTS OF PULSE PRESSURE: AN INTEGRATED NUMERICAL AND PHYSIOLOGICAL APPROACH

Samuel Vennin, Ye Li, Marie Willemet, Henry Fok, Haotian Gu, Peter Charlton, Jordi Alastruey, Phil Chowiecnyk
King's College London, UK

Purpose: Hypertension, the single biggest killer worldwide¹, arises mainly as a result of an increase in central pulse pressure (PP)², yet haemodynamic basis of that increase is still disputed. We examined the ability of a simple "reduced" model comprising a proximal characteristic impedance linked to a Windkessel element to accurately predict PP from aortic blood flow and applied the model to examine PP dependence on cardiac and vascular properties.

Method: PP obtained from the model was compared with theoretical values obtained in silico and in vivo. Theoretical values were obtained using a distributed multi-segment model in a population of "virtual" subjects ($n = 3,095$) in which cardiovascular properties were varied over the pathophysiological range. In vivo measurements were in normotensive subject ($n = 13$) during modulation of physiology with vasoactive drugs with divergent actions on cardiac and vascular properties and in hypertensive subjects ($n = 156$).

Results: PP derived from the model agreed with theoretical values (mean difference SD, -0.09 ± 1.96 mmHg) and with measured values (-1.95 ± 3.74 and -1.18 ± 3.67 mmHg for normotensive and hypertensive subjects respectively). Parameters extracted from the model agreed closely with theoretical and measured physical properties. PP was seen to be determined mainly by total arterial compliance (inversely associated with arterial stiffness) and ventricular dynamics: the volume of blood ejected up to time of pulse pressure and the rate of ventricular ejection up to this point.

Conclusion: Increased flow and/or volume accounted for 20.1 mmHg (52%) of the 39.0 mmHg difference in pulse pressure between the upper and lower tertiles of the hypertensive subjects.

References

1. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the global burden of disease study 2010. *Lancet*. 2012;380:2224–2260
2. Franklin SS, Gustin W, Wong ND, Larson MG, Weber MA, Kannel WB, Levy D. Hemodynamic patterns of age-related changes in blood pressure. The framingham heart study. *Circulation*. 1997;96:308–315

P122

CALCULATING RESERVOIR PRESSURE WITH OR WITHOUT FLOW INFORMATION: SIMILARITY AND ALGORITHMIC SENSITIVITY AT RADIAL ARTERY

Michael Ebner¹, Kim Parker², Tom Vercauteren^{3,1}, Sébastien Ourselin^{3,1}, Siegfried Wassertheurer⁴, Alun Hughes⁵, Bernhard Hametner⁴
¹Translational Imaging Group, Centre for Medical Image Computing, Department of Medical Physics and Biomedical Engineering, University College London, London, UK

²Department of Bioengineering, Imperial College London, London, UK

³Wellcome / EPSRC Centre for Interventional and Surgical Sciences, University College London, London, UK

⁴Center for Health & Bioresources, AIT Austrian Institute of Technology, Vienna, Austria

⁵Institute of Cardiovascular Science, University College London, London, UK

Background: Reservoir pressure is typically estimated from the pressure waveform information only. Comparability with estimates made using pressure and flow depend on assumptions, e.g. a proportional relationship between excess pressure and flow [1]. In this study, we compared (i) results using flow and pressure versus pressure-only at the radial artery, and (ii) two different algorithms used in the literature for pressure-only analysis.

Methods: Reservoir pressure separations were performed on 95 hypertensive individuals where radial pressure and flow velocity waveform measurements were available [2]. Algorithm (F) used flow and pressure information [3]. Algorithms (P1) and (P2) refer to the two different pressure-only implementations as used in [4, 5], and [1, 6], respectively. Reservoir curves characterized by physiologically implausible parameters, i.e. a rate constant $b < 0$ or an asymptotic pressure $P_{\infty} < 0$, were discarded, leaving 63 subjects with valid reservoir pressure data.

Results: Estimated reservoir parameters are shown in Table 1. Algorithm (F) showed statistically significant differences in most of the parameters compared to (P1) and (P2), although, except time constant τ and asymptotic pressure P_{∞} , there was a strong correlation between methods. Significant differences were observed in reservoir pulse pressure and area estimates between (P1) and (P2) despite their, in general, high correlation.

Table 1. Quantification of reservoir pressures p_{res} obtained by methods (F), (P1) and (P2) at radial artery in the format of mean $\pm$ standard deviation based on 63 subjects whereby PP denotes the reservoir pulse pressure, A_p the area of reservoir pressure above diastolic blood pressure, τ the time constant describing the diastolic pressure decay, P_{∞} the asymptotic blood pressure and $a, b = 1/\tau$ the rate constants. Peripheral (area) resistance and compliance, i.e. R and C, were estimated from the rate constants a and b for (P1) and (P2) using flow information. The correlation coefficient r was computed between relevant methods. The statistical significance of the differences between methods was based on a paired t-test with * indicating $p < 0.05$.

Radial artery	p_{res} (F)	p_{res} (P1)	p_{res} (P2)	$r(F,P1)$	$r(F,P2)$	$r(P1,P2)$
PP [mmHg]	41.5 $\pm$ 10.0	36.3 $\pm$ 7.2	35.7 $\pm$ 7.0	0.82*	0.82*	0.96*
A_p [mmHg s]	17.5 $\pm$ 4.3	15.6 $\pm$ 3.7	15.5 $\pm$ 3.7	0.94*	0.94*	1.00*
τ [s]	0.3 $\pm$ 0.1	0.6 $\pm$ 0.4	0.6 $\pm$ 0.3	0.36*	0.42*	0.88
P_{∞} [mmHg]	65.7 $\pm$ 10.3	63.9 $\pm$ 15.2	64.8 $\pm$ 12.6	0.45	0.53	0.79
a [1/s]	—	8.1 $\pm$ 5.2	7.4 $\pm$ 2.7	—	—	0.93
b [1/s]	—	2.2 $\pm$ 1.1	2.1 $\pm$ 0.8	—	—	0.84
R [mmHg s/m]	419.0 $\pm$ 188.8	453.7 $\pm$ 348.2	436.7 $\pm$ 302.6	0.68	0.75	0.92
C [mm/mmHg]	0.8 $\pm$ 0.3	1.7 $\pm$ 1.0	1.7 $\pm$ 1.0	0.70*	0.70*	1.00

Conclusions: The discrepancies between (F) and (P1), (P2) raise concerns about the validity of the implicit assumptions in pressure-only reservoir pressure separation at the radial artery. Differences in (P1) and (P2) indicate some sensitivity of derived parameters to the algorithm employed.

References

- [1] J. Aguado-Sierra, J. Alastruey, J.-J. Wang, N. Hadjiloizou, J. Davies, and K. H. Parker, "Separation of the reservoir and wave pressure and velocity from measurements at an arbitrary location in arteries," *Proc. Inst. Mech. Eng. Part H J. Eng. Med.*, vol. 222, no. 4, pp. 403–416, 2008.
- [2] C. H. Manisty, A. Zambanini, K. H. Parker, J. E. Davies, D. P. Francis, J. Mayet, S. A. McG Thom, and A. D. Hughes, "Differences in the magnitude of wave reflection account for differential effects of amlodipine- versus atenolol-based regimens on central blood pressure: an Anglo-Scandinavian Cardiac Outcome Trial substudy," *Hypertension*, vol. 54, no. 4, pp. 724–30, Oct. 2009.
- [3] J.-J. Wang, A. B. O'Brien, N. G. Shrive, K. H. Parker, and J. V. Tyberg, "Time-domain representation of ventricular-arterial coupling as a wind-kessel and wave system," *Am. J. Physiol. Heart Circ. Physiol.*, vol. 284, no. 4, pp. H1358–68, Apr. 2003.
- [4] M. Ebner, B. Hametner, S. Parragh, and S. Wassertheurer, "Reservoir Wave Paradigm: An Implementation and Sensitivity Analysis," *SNE Simul. Notes Eur.*, vol. 25, no. 3–4, pp. 151–156, 2015.
- [5] B. Hametner, S. Wassertheurer, A. D. Hughes, K. H. Parker, T. Weber, and B. Eber, "Reservoir and excess pressures predict cardiovascular events in high-risk patients," *Int. J. Cardiol.*, vol. 171, no. 1, pp. 31–36, 2014.
- [6] J. E. Davies, P. Lacy, T. Tillin, D. Collier, J. K. Cruickshank, D. P. Francis, A. Malaweera, J. Mayet, A. Stanton, B. Williams, K. H. Parker, S. A. McG Thom, and A. D. Hughes, "Excess pressure integral predicts cardiovascular events independent of other risk factors in the conduit artery functional evaluation substudy of Anglo-Scandinavian Cardiac Outcomes Trial," *Hypertension*, vol. 64, no. 1, pp. 60–8, Jul. 2014.

P123

RESERVOIR PRESSURE IS INDEPENDENTLY ASSOCIATED WITH 11–12 YEAR OLD'S KIDNEY FUNCTION: POPULATION-DERIVED STUDY

Xiaoqing Peng¹, Martin Schultz¹, Michael Cheung^{2,3,4}, Melissa Wake^{2,3,5}, Jonathan Mynard^{2,3,4}, David Burgner^{2,3,6}, Richard Liu³, James Sharman¹

¹Menzies Institute for Medical Research, University of Tasmania, Hobart, Tas, Australia

²The University of Melbourne, Parkville, Victoria, Australia

³Murdoch Children's Research Institute, Parkville, Victoria, Australia

⁴The Royal Children's Hospital, Melbourne, Victoria, Australia

⁵The University of Auckland, Grafton, Auckland, New Zealand

⁶Monash University, Clayton, Victoria, Australia

Introduction: Reservoir pressure (RP) and excess pressure (XSP) independently predict cardiovascular events in adults, but have never been investigated as markers of cardiovascular risk among children. This study aimed to determine the association of RP and XSP with end-organ markers of cardiovascular risk related to kidney function and large artery pre-atherosclerosis.

Methods: Participants were 1874 11–12 year-old children (50% male) from the Longitudinal Study of Australian Children's Child Health CheckPoint study. Brachial blood pressure was measured by cuff oscillometric device (SphygmoCor XCEL, AtCor, Sydney). The same device was used to derive reservoir pressure (RP) and excess pressure (XSP) from the brachial pressure waveform. Kidney function was determined from estimated glomerular filtration rate (eGFR, $n = 926$) and large artery pre-atherosclerosis was determined from carotid intima media thickness (cIMT, $n = 1131$) using ultrasound.

Results: The eGFR was significantly correlated with RP peak ($r = -0.109$, $p = 0.001$), RP integral ($r = -0.136$, $p < 0.001$), XSP peak ($r = 0.096$, $p = 0.004$) and XSP integral ($r = 0.102$, $p = 0.002$). The RP (whether expressed as peak or integral) was significantly associated with eGFR after adjusting for sex, waist-to-hip ratio, heart rate and brachial BP indices (RP peak $\beta = -0.079$, $p = 0.02$, partial $R^2 = 0.006$ and RP integral $\beta = -0.079$, $p = 0.02$, partial $R^2 = 0.007$). XSP was not independently associated with eGFR after adjusting for the above variables. Neither RP nor XSP were significantly associated with cIMT.

Conclusion: Independent of conventional risk factors, RP was significantly associated with kidney function among a large population of Australian

children. The non-invasive method to derive RP using an oscillometric cuff device could provide useful clinical information in children.

P124

VALIDITY AND RELIABILITY OF PULSE WAVE ANALYSIS ESTIMATED BY A NOVEL WRIST-WORN TONOMETER

Luis Garcia-Ortiz^{1,2}, Jose Recio-Rodriguez^{1,3}, Cristina Agudo-Conde¹, Rita Salvado¹, Jose Maderuelo-Fernandez¹, Jesus Gonzalez-Sanchez^{4,5}, Emiliano Rodriguez-Sanches^{1,6}, Manuel Gómez-Marcos^{1,6} on behalf EVA_Investigators⁶

¹Primary Health Care Research Unit, La Alamedilla Health Center, Health Service of Castilla y León (SACYL), Biomedical Research Institute of Salamanca (IBSAL), Salamanca, Spain

²Department of Biomedical and Diagnostic Sciences, University of Salamanca, Salamanca, Spain

³Department of Nurse and Physiotherapy, University of Salamanca, Salamanca, Spain

⁴Primary Care Research Unit, The Alamedilla Health Center, Castilla and León Health Service (SACYL), Biomedical Research Institute of Salamanca (IBSAL)

⁵Department of Nursing, University of Extremadura, Plasencia, Cáceres, Spain

⁶Department of Medicine, University of Salamanca, Salamanca, Spain

Objective: To analyze the reliability and validity of Pulse Wave Analysis determined with the new wrist-worn tonometry.

Methods: Cross sectional study including 254 subjects. Aged 51.9 ± 13.4 , being women 53%. Main measurements: Peripheral Alx (PALx) and Central Alx (CALx) by wrist-worn tonometry and Sphygmocor. Carotid femoral (cf) pulse wave velocity (PWV) by Sphygmocor, Cardio Ankle Vascular index (CAVI), ankle brachial index (ABI) and brachial ankle (ba) PWV by Vasera device. Carotid intima media thickness (IMT) by ultrasonography.

Results: Intra-class correlation coefficient (ICC) intraobserver for the PALx was 0.886 (95% CI 0.803 to 0.934) and for the CALx 0.943 (0.901 to 0.968) with a Bland Altman agreement limit of -0.75 (-23.8 to 21.8) and 0.08 (-15.7 to 15.9) respectively. ICC interobserver for PALx was 0.952 (95% CI 0.915 to 0.972) and CALx 0.893 (0.811 to 0.939) with an agreement limit of -0.45 (-13.7 to 12.8) and 0.43 (-17.7 to 1835) respectively. We found, compared with Sphygmocor, an ICC of 0.849 (0.798 to 0.887) for PALx, and 0.783 (0.711 to 0.838) for CALx. The agreement limit for PALx was -1.03 (-22.73 to 20.67) and CALx 2.14 (-20.50 to 24.79). We found positive correlation with PALx, CALx and CALx HR75 by Aurora with age, CAVI, ABI, baPWV, cfPWV, IMT and cardiovascular risk and negative with glomerular filtration rate.

Conclusions: The wrist-worn tonometry shows an adequate reliability intra and interobserver, and interdevice when compared to Sphygmocor, and an adequate validity when compared with other measures that evaluate arterial stiffness, target organ damage and cardiovascular risk.

P125

USE OF VASCULAR ADAPTATION IN RESPONSE TO MECHANICAL LOADING FACILITATES PERSONALISATION OF A ONE-DIMENSIONAL PULSE WAVE PROPAGATION MODEL

Maarten Heusinkveld, Koen Reesink, Theo Arts, Wouter Huberts, Tammo Delhaas

Department of Biomedical Engineering, CARIM School for Cardiovascular Diseases, Maastricht University, The Netherlands

Background: Mathematical modelling of pressure and flow waveforms in blood vessels using pulse wave propagation (PWP) models could support clinical decision-making. For a personalised model outcome, measurements of all modelled vessel radii and wall thicknesses are required. In clinical practice, however, datasets are often incomplete. To overcome this problem, we hypothesised that the adaptive capacity of blood vessels in response to mechanical load can be utilised to fill in the gaps of incomplete patient-specific datasets.

Methods: We implemented homeostatic feedback loops in a validated PWP model [1] to allow adaptation of vessel geometry to maintain wall stress and wall shear stress. To evaluate our approach, we utilised complete