



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantispress.com/journals/artres>

P124: VALIDITY AND RELIABILITY OF PULSE WAVE ANALYSIS ESTIMATED BY A NOVEL WRIST-WORN TONOMETER

Luis Garcia-Ortiz, Jose Recio-Rodriguez, Cristina Agudo-Conde, Rita Salvado, Jose Maderuelo-Fernandez, Jesus Gonzalez-Sanchez, Emiliano Rodriguez- Sanches, Manuel Gómez-Marcos

To cite this article: Luis Garcia-Ortiz, Jose Recio-Rodriguez, Cristina Agudo-Conde, Rita Salvado, Jose Maderuelo-Fernandez, Jesus Gonzalez-Sanchez, Emiliano Rodriguez-Sanches, Manuel Gómez-Marcos (2017) P124: VALIDITY AND RELIABILITY OF PULSE WAVE ANALYSIS ESTIMATED BY A NOVEL WRIST-WORN TONOMETER, Artery Research 20:C, 79–79, DOI: <https://doi.org/10.1016/j.artres.2017.10.106>

To link to this article: <https://doi.org/10.1016/j.artres.2017.10.106>

Published online: 7 December 2019

Conclusions: The discrepancies between (F) and (P1), (P2) raise concerns about the validity of the implicit assumptions in pressure-only reservoir pressure separation at the radial artery. Differences in (P1) and (P2) indicate some sensitivity of derived parameters to the algorithm employed.

References

- [1] J. Aguado-Sierra, J. Alastruey, J.-J. Wang, N. Hadjiloizou, J. Davies, and K. H. Parker, "Separation of the reservoir and wave pressure and velocity from measurements at an arbitrary location in arteries," *Proc. Inst. Mech. Eng. Part H J. Eng. Med.*, vol. 222, no. 4, pp. 403–416, 2008.
- [2] C. H. Manisty, A. Zambanini, K. H. Parker, J. E. Davies, D. P. Francis, J. Mayet, S. A. McG Thom, and A. D. Hughes, "Differences in the magnitude of wave reflection account for differential effects of amlodipine- versus atenolol-based regimens on central blood pressure: an Anglo-Scandinavian Cardiac Outcome Trial substudy," *Hypertension*, vol. 54, no. 4, pp. 724–30, Oct. 2009.
- [3] J.-J. Wang, A. B. O'Brien, N. G. Shrive, K. H. Parker, and J. V. Tyberg, "Time-domain representation of ventricular-arterial coupling as a wind-kessel and wave system," *Am. J. Physiol. Heart Circ. Physiol.*, vol. 284, no. 4, pp. H1358–68, Apr. 2003.
- [4] M. Ebner, B. Hametner, S. Parragh, and S. Wassertheurer, "Reservoir Wave Paradigm: An Implementation and Sensitivity Analysis," *SNE Simul. Notes Eur.*, vol. 25, no. 3–4, pp. 151–156, 2015.
- [5] B. Hametner, S. Wassertheurer, A. D. Hughes, K. H. Parker, T. Weber, and B. Eber, "Reservoir and excess pressures predict cardiovascular events in high-risk patients," *Int. J. Cardiol.*, vol. 171, no. 1, pp. 31–36, 2014.
- [6] J. E. Davies, P. Lacy, T. Tillin, D. Collier, J. K. Cruickshank, D. P. Francis, A. Malaweera, J. Mayet, A. Stanton, B. Williams, K. H. Parker, S. A. McG Thom, and A. D. Hughes, "Excess pressure integral predicts cardiovascular events independent of other risk factors in the conduit artery functional evaluation substudy of Anglo-Scandinavian Cardiac Outcomes Trial," *Hypertension*, vol. 64, no. 1, pp. 60–8, Jul. 2014.

P123

RESERVOIR PRESSURE IS INDEPENDENTLY ASSOCIATED WITH 11–12 YEAR OLD'S KIDNEY FUNCTION: POPULATION-DERIVED STUDY

Xiaoqing Peng¹, Martin Schultz¹, Michael Cheung^{2,3,4}, Melissa Wake^{2,3,5}, Jonathan Mynard^{2,3,4}, David Burgner^{2,3,6}, Richard Liu³, James Sharman¹

¹Menzies Institute for Medical Research, University of Tasmania, Hobart, Tas, Australia

²The University of Melbourne, Parkville, Victoria, Australia

³Murdoch Children's Research Institute, Parkville, Victoria, Australia

⁴The Royal Children's Hospital, Melbourne, Victoria, Australia

⁵The University of Auckland, Grafton, Auckland, New Zealand

⁶Monash University, Clayton, Victoria, Australia

Introduction: Reservoir pressure (RP) and excess pressure (XSP) independently predict cardiovascular events in adults, but have never been investigated as markers of cardiovascular risk among children. This study aimed to determine the association of RP and XSP with end-organ markers of cardiovascular risk related to kidney function and large artery pre-atherosclerosis. **Methods:** Participants were 1874 11–12 year-old children (50% male) from the Longitudinal Study of Australian Children's Child Health CheckPoint study. Brachial blood pressure was measured by cuff oscillometric device (SphygmoCor XCEL, AtCor, Sydney). The same device was used to derive reservoir pressure (RP) and excess pressure (XSP) from the brachial pressure waveform. Kidney function was determined from estimated glomerular filtration rate (eGFR, $n = 926$) and large artery pre-atherosclerosis was determined from carotid intima media thickness (cIMT, $n = 1131$) using ultrasound.

Results: The eGFR was significantly correlated with RP peak ($r = -0.109$, $p = 0.001$), RP integral ($r = -0.136$, $p < 0.001$), XSP peak ($r = 0.096$, $p = 0.004$) and XSP integral ($r = 0.102$, $p = 0.002$). The RP (whether expressed as peak or integral) was significantly associated with eGFR after adjusting for sex, waist-to-hip ratio, heart rate and brachial BP indices (RP peak $\beta = -0.079$, $p = 0.02$, partial $R^2 = 0.006$ and RP integral $\beta = -0.079$, $p = 0.02$, partial $R^2 = 0.007$). XSP was not independently associated with eGFR after adjusting for the above variables. Neither RP nor XSP were significantly associated with cIMT.

Conclusion: Independent of conventional risk factors, RP was significantly associated with kidney function among a large population of Australian

children. The non-invasive method to derive RP using an oscillometric cuff device could provide useful clinical information in children.

P124

VALIDITY AND RELIABILITY OF PULSE WAVE ANALYSIS ESTIMATED BY A NOVEL WRIST-WORN TONOMETER

Luis Garcia-Ortiz^{1,2}, Jose Recio-Rodriguez^{1,3}, Cristina Agudo-Conde¹, Rita Salvado¹, Jose Maderuelo-Fernandez¹, Jesus Gonzalez-Sanchez^{4,5}, Emiliano Rodriguez-Sanches^{1,6}, Manuel Gómez-Marcos^{1,6} on behalf EVA_Investigators⁶

¹Primary Health Care Research Unit, La Alamedilla Health Center, Health Service of Castilla y León (SACYL), Biomedical Research Institute of Salamanca (IBSAL), Salamanca, Spain

²Department of Biomedical and Diagnostic Sciences, University of Salamanca, Salamanca, Spain

³Department of Nurse and Physiotherapy, University of Salamanca, Salamanca, Spain

⁴Primary Care Research Unit, The Alamedilla Health Center, Castilla and León Health Service (SACYL), Biomedical Research Institute of Salamanca (IBSAL)

⁵Department of Nursing, University of Extremadura, Plasencia, Cáceres, Spain

⁶Department of Medicine, University of Salamanca, Salamanca, Spain

Objective: To analyze the reliability and validity of Pulse Wave Analysis determined with the new wrist-worn tonometry.

Methods: Cross sectional study including 254 subjects. Aged 51.9 ± 13.4 , being women 53%. Main measurements: Peripheral Alx (PALx) and Central Alx (CALx) by wrist-worn tonometry and Sphygmocor. Carotid femoral (cf) pulse wave velocity (PWV) by Sphygmocor, Cardio Ankle Vascular index (CAVI), ankle brachial index (ABI) and brachial ankle (ba) PWV by Vasera device. Carotid intima media thickness (IMT) by ultrasonography.

Results: Intra-class correlation coefficient (ICC) intraobserver for the PALx was 0.886 (95% CI 0.803 to 0.934) and for the CALx 0.943 (0.901 to 0.968) with a Bland Altman agreement limit of -0.75 (-23.8 to 21.8) and 0.08 (-15.7 to 15.9) respectively. ICC interobserver for PALx was 0.952 (95% CI 0.915 to 0.972) and CALx 0.893 (0.811 to 0.939) with an agreement limit of -0.45 (-13.7 to 12.8) and 0.43 (-17.7 to 1835) respectively. We found, compared with Sphygmocor, an ICC of 0.849 (0.798 to 0.887) for PALx, and 0.783 (0.711 to 0.838) for CALx. The agreement limit for PALx was -1.03 (-22.73 to 20.67) and CALx 2.14 (-20.50 to 24.79). We found positive correlation with PALx, CALx and CALx HR75 by Aurora with age, CAVI, ABI, baPWV, cfPWV, IMT and cardiovascular risk and negative with glomerular filtration rate.

Conclusions: The wrist-worn tonometry shows an adequate reliability intra and interobserver, and interdevice when compared to Sphygmocor, and an adequate validity when compared with other measures that evaluate arterial stiffness, target organ damage and cardiovascular risk.

P125

USE OF VASCULAR ADAPTATION IN RESPONSE TO MECHANICAL LOADING FACILITATES PERSONALISATION OF A ONE-DIMENSIONAL PULSE WAVE PROPAGATION MODEL

Maarten Heusinkveld, Koen Reesink, Theo Arts, Wouter Huberts, Tammo Delhaas

Department of Biomedical Engineering, CARIM School for Cardiovascular Diseases, Maastricht University, The Netherlands

Background: Mathematical modelling of pressure and flow waveforms in blood vessels using pulse wave propagation (PWP) models could support clinical decision-making. For a personalised model outcome, measurements of all modelled vessel radii and wall thicknesses are required. In clinical practice, however, datasets are often incomplete. To overcome this problem, we hypothesised that the adaptive capacity of blood vessels in response to mechanical load can be utilised to fill in the gaps of incomplete patient-specific datasets.

Methods: We implemented homeostatic feedback loops in a validated PWP model [1] to allow adaptation of vessel geometry to maintain wall stress and wall shear stress. To evaluate our approach, we utilised complete