



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

5.4: HIGH PWV IS ASSOCIATED WITH NANO-SCALE CHANGES IN THE MEDIAL LAYER OF THE INTERNAL MAMMARY ARTERY

Riaz Akhtar, Zhuo Chang, Maria Lyck Hansen, Lars Melholt Rasmussen

To cite this article: Riaz Akhtar, Zhuo Chang, Maria Lyck Hansen, Lars Melholt Rasmussen (2016) 5.4: HIGH PWV IS ASSOCIATED WITH NANO-SCALE CHANGES IN THE MEDIAL LAYER OF THE INTERNAL MAMMARY ARTERY, Artery Research 16:C, 57–57, DOI: <https://doi.org/10.1016/j.artres.2016.10.032>

To link to this article: <https://doi.org/10.1016/j.artres.2016.10.032>

Published online: 7 December 2019

Methods: We studied 561 (50% male) normotensive and hypertensive subjects without kidney or other cardiovascular diseases or antihypertensive treatment. Supine and upright hemodynamics were recorded using continuous pulse wave analysis, whole body impedance cardiography and heart rate variability analysis. Estimated glomerular filtration rate (eGFR) was calculated using the CKD-EPI cystatin C equation.

Results: Mean eGFR was 99 (range 53-152) ml/min/1.73 m² and one third of the patients had values below 90. After adjustments for age, sex, body mass index and low density lipoprotein cholesterol level, regression analysis indicated significant associations between lower eGFR and higher systolic ($p < 0.001$) and diastolic blood pressure ($p < 0.001$) and systemic vascular resistance ($p = 0.001$) regardless of body position. Lower eGFR was associated with higher low frequency to high frequency ratio of heart rate variability in supine but not in upright position. The level of eGFR was not associated with the level of cardiac output.

Conclusions: Even mild kidney impairment is associated with higher systemic vascular resistance and increased supine sympathovagal balance. However, changes in autonomic tone, as based on analysis of heart rate variability, do not seem to explain the relation between lower eGFR and higher systemic vascular resistance in the upright position. The close relationship between the regulation of GFR and systemic vascular resistance may play a role in the pathogenesis of primary hypertension.

5.2

AN ASSOCIATED WITH FAMILIAL HEMIPLEGIC MIGRAINE TYPE 2 MUTATION IN THE ALPHA-2 ISOFORM NA,K-ATPASE DISTURBS VASCULAR RESPONSES IN MOUSE BRAIN

Mr Christian Staehr, Mrs Lise Hangaard, Dr Karin Lykke-Hartmann, Dr Elena Bouzinova, Professor Christian Aalkjaer, [Dr Vladimir Matchkov](#)
Aarhus University, Aarhus, Denmark

Objectives: Migraine attack is associated with severe changes in brain perfusion vasoconstriction-induced hypoxemia during aura and rebound vasodilation in subsequent headache. Familial Hemiplegic Migraine Type 2 is associated with point mutations (including G301R) in the $\alpha 2$ isoform Na,K-ATPase. Heterozygote mice bearing G301R mutation (FHM2) were recently characterized for several behavioral and neuronal abnormalities.

Methods: Vascular function of wild type (WT) and FHM2 mice was compared in vivo (telemetry and Laser Speckle measurements of brain perfusion), in vitro (myography) and in situ (changes in astrocytic $[Ca^{2+}]_i$ and parenchymal arteriole diameter in brain slices to electric field stimulation (EFS)).

Results: Vascular abnormalities were shown for cerebral circulation while only minor or no significant changes were found in peripheral arteries. Accordingly, no difference in blood pressure was seen under resting conditions. Middle cerebral artery from FHM2 mice had large inner diameter and constricted stronger to U46619, endothelin and K⁺-depolarization. This was associated with increased depolarization and Src-kinase-dependent sensitization to $[Ca^{2+}]_i$.

Isolated cerebral arteries from FHM2 mice have exaggerated relaxation to elevated $[K^+]_{out}$ (4-12mM) due to increased role of the inward-rectifying K⁺ channels. Repeated EFS (>3 times) reduced the $[Ca^{2+}]_i$ responses in astrocytic endfeet and increased relaxation of parenchymal arterioles in the FHM2 in comparison with WT. Flow responses to whiskers stimulation were also potentiated in FHM2 mice.

Conclusions: A knock-out mutation of the $\alpha 2$ Na,K-ATPase leads to both elevated contractility and increased relaxation of cerebral arteries. These dysfunctions could affect the blood supply to active neurons and thus disturb neurovascular coupling.

5.3

REVERSIBILITY OF ARTERIAL STIFFNESS AFTER KIDNEY TRANSPLANTATION: SYSTEMATIC REVIEW AND META-ANALYSIS

Ms Aboubacar Sidibe, Ms Catherine Fortier, Miss Marie-Pier Desjardins, Dr Fabrice Mac-Way, Dr Sacha De Serres, [Dr Mohsen Agharazii](#)
Centre de Recherche du CHU de Quebec, Division of Nephrology, Faculty of Medicine, Department of Medicine and Social and Preventive Medicine, Université Laval, Quebec City, Canada

Background: Chronic kidney disease is associated with increased arterial stiffness. Correction of the uremic milieu by kidney transplantation (KTx) may be improve arterial stiffness. However, results from clinical studies are not uniformly convincing. This could be related to small sample size of studies,

heterogeneity in methods and timing of assessment of arterial stiffness after KTx. We aim to measure the reversibility of arterial stiffness after KTx.

Design and Method: Observational studies and randomized controlled trials with measurements of pulse wave velocity (PWV), pulse pressure (PP) and/or augmentation index (Alx) were extracted from MEDLINE, EMBASE, COCHRANE LIBRARY, and Web of Science from their inception to January 2016. Two reviewers independently identified eligible studies comparing PW, PP and/or Alx pre to post KTx and extracted data including population characteristics, interventions and outcomes.

Results: 13 studies of 981 met our inclusion criteria. 11 Studies (408 renal transplant) have been included in meta-analysis. There was a standard mean change of PWV by -0.45 (95% CI: - 0.68 -0.20, I²=58%) post-KTx. Both studies using aortic PWV (5 studies, 160 patients) and those using brachial-ankle PWV, showed a significant decrease of PWV by -1.58 m/s (95% CI: -2.97 - 0.19, I²= 87%) and by -1.21 m/s (95% CI: - 1.89 - 0.54, I²=0 %) post-KTx, respectively. Analysis of central PP and Alx showed significant reduction post-KTx by -4.77 (95% CI: -9.19 -0.35, I²=55%) and by -11.59 (95% CI: -15.64 -7.53, I²=43%), respectively. Only two studies have reported adjusted parameters for mean arterial pressure.

Conclusions: There is a significant reduction in PWV, central PP and Alx after KTx. Heterogeneity among studies are globally moderate. Further analysis is required to examine the importance of changes in different vascular beds taking into account changes in blood pressure.

5.4

HIGH PWV IS ASSOCIATED WITH NANO-SCALE CHANGES IN THE MEDIAL LAYER OF THE INTERNAL MAMMARY ARTERY

[Dr Riaz Akhtar](#)¹, [Mr Zhuo Chang](#)¹, [Dr Maria Lyck Hansen](#)², Professor Lars Melholt Rasmussen²

¹University of Liverpool, UK

²Odense University Hospital, Odense, Denmark

Background: Arterial stiffening occurs as part of the natural ageing process. Degradation of the extracellular matrix (ECM) in the medial layer is typically implicated in arterial stiffening. However, little is known about how localised changes in arteries in terms of both structure and mechanical properties contribute to the overall stiffening of arteries.

Aim: To determine localised differences in the nano-structure and mechanical properties in the medial layer of internal mammary arteries (IMA) in patients with high and low pulse wave velocity (PWV).

Methods: IMAs were collected from coronary bypass operations from 7 patients with high (13.8 3.3 m/s) and 7 patients with low (8.6 0.7 m/s) PWV. The samples were cryo-sectioned to a nominal thickness of 5 μ m for atomic force microscopy (AFM) measurement. All the samples were tested hydrated. Histological analysis was used to determine collagen and elastin content. Data are presented as means SEMs.

Results: The medial layers of IMAs in the high PWV group were significant stiffer than in the low PWV group (Low 228.4 15.6 kPa, High 735.8 108.8 kPa,) ($p < 0.0001$). Topographical features as visualised with AFM were similar in both groups but the higher nanomechanical stiffness was found to correlate with histological data.

Conclusions: Nanomechanical properties of the medial layer in the IMA associate with PWV data. Changes in composition in the ECM drive the profound localized changes in tissue stiffness.

5.5

AGE-DEPENDENT TELOMERE ATTRITION, SHORT TELOMERES AND ATHEROSCLEROSIS

[Dr Simon Toupance](#)¹, [Dr Anna Kearney-Schwartz](#)², [Dr Mohamed Temmar](#)³, [Mrs Cécile Lakomy](#)¹, [Mr Carlos Labat](#)¹, [Professor Patrick Rossignol](#)⁴, [Professor Faiez Zannad](#)⁴, [Dr Patrick Lacolley](#)¹, [Professor Abraham Aviv](#)⁵, [Professor Athanase Benetos](#)²

¹INSERM UMRS 1116, Université de Lorraine, Nancy, France

²Département de Médecine Gériatrique, CHU de Nancy, France

³Cardiology Center, Ghardaia, Algeria

⁴CIC-P Pierre Drouin, CHU de Nancy, France

⁵Center of Human Development and Aging, Rutgers University, NJ, USA

Background: Short leukocyte telomere length (LTL) is associated with atherosclerosis. The prevailing view is this association exists since LTL is a biomarker of cumulative inflammation and oxidative stress during adult life. However recent studies show that LTL in adults is defined