



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

8.5: HEMODYNAMICS DURING INTRA- AND INTERDIALYTIC PERIODS DEPEND ON ULTRAFILTRATION VOLUME

Christopher Mayer, Stephan Geilert, Julia Matschkal, Uwe Heemann, Marcus Baumann, Christoph Schmaderer

To cite this article: Christopher Mayer, Stephan Geilert, Julia Matschkal, Uwe Heemann, Marcus Baumann, Christoph Schmaderer (2016) 8.5: HEMODYNAMICS DURING INTRA- AND INTERDIALYTIC PERIODS DEPEND ON ULTRAFILTRATION VOLUME, Artery Research 16:C, 66–66, DOI: <https://doi.org/10.1016/j.artres.2016.10.061>

To link to this article: <https://doi.org/10.1016/j.artres.2016.10.061>

Published online: 7 December 2019

8.5 HEMODYNAMICS DURING INTRA- AND INTERDIALYTIC PERIODS DEPEND ON ULTRAFILTRATION VOLUME

Christopher Mayer¹, Stephan Geilert², Julia Matschkal², Uwe Heemann², Marcus Baumann², Christoph Schmaderer²
¹AIT Austrian Institute of Technology GmbH, Vienna, Austria
²Technical University of Munich, Germany

Introduction: Parameters of arterial stiffness are independent cardiovascular risk factors for end-stage renal disease patients. Significant changes of these parameters between intra- and interdialytic periods have been reported previously [1]. The aim of this cross-sectional study is to describe the influence of the ultrafiltration volume on hemodynamic parameters.

Methods: All measurements were obtained with the Mobil-O-Graph 24h PWA (I.E.M. GmbH, Germany) within the ISAR hemodialysis study. Measurement started before the midweek dialysis session and lasted for 24-hours. 348 patients (238 male / 110 female 65 ± 18 years) were included. Intra- and interdialytic parameters were averaged and compared for three subgroups (ultrafiltration volume (UFV) ≤ 500 ml (N = 50) 500 < UFV < 2000 ml (N = 159) UFV > 2000 ml (N = 139)) and all subjects.

Results: The results for all patients support the findings of Karpetas et al. [1] (see Table). Beyond [1], the results underpin the differences between subgroups for intra- and interdialytic periods. Furthermore, there are significant differences between intra- and interdialytic periods depending on the ultrafiltration volume (see Table). Exemplarily, there is a significant rise in the augmentation index (26.0 vs. 28.5%, $p < 0.05$) for UFV > 2000 ml and for central pulse pressure (39.6 vs. 43.4 mmHg and 36.0 vs. 38.3, $p < 0.05$) for UFV ≤ 2000 ml opposed to non-significance for the other subgroups.

Conclusions: Our findings support the hypothesis that hemodynamic parameters depend on ultrafiltration volume. Further studies should investigate their prognostic value considering the ultrafiltration volume.

compliance, but its effect on peripheral arterial compliance (pC) is controversial. We aimed to test the hypotheses that aerobic training augments pC at rest and during different autonomic nervous system provocations (ANSP) in young healthy men.

We enrolled 44 males, 19-24 years old (22 trained, VO₂max=485ml/kg/min – group A, 22 sedentary controls, VO₂max=303ml/kg/min – group B). VO₂max was determined using cycloergometry (QuarkCPET, Cosmed). On the testing day, ECG, arterial blood pressure (Finapres, Ohmeda) and finger artery compliance at rest, 3 minutes during 0.1Hz breathing and 3 min during mental stress were measured. A noninvasive method was used to determine compliance index (CI), calculated as an average of the pressure dependant compliance curve in the range of arterial pressures from 97 to 105mmHg.

Our results revealed elevated CI in group A compared to group B (4.18±0.38 and 1.28±0.25, $p=0.004$) at rest and no significant differences in CI between groups during ANSP (1.34±0.20 at 0.1Hz breathing and 0.82±0.18 during mental stress in group A compared to 1.09±0.21, $p=0.06$, and 0.60±0.12, $p=0.08$ in group B). A statistically significant positive linear correlation existed between CI and VO₂max ($P < 0.001$) at rest in group A, however, no correlation was found at both ANSP.

Regular aerobic training increases pC in healthy young subjects at rest, but not during 0.1Hz breathing or mental stress. Our findings indicate that peripheral and not central autonomic mechanisms govern pC in young healthy males.

References

Terenzi TJ. An alteration in arterial compliance associated with elevated aerobic fitness. *J Manipulative Physiol Ther.* 2000 Jan23(1):27-31.
 Ashor AW, Lara J, Siervo M, Celis-Morales C, Mathers JC. Effects of exercise modalities on arterial stiffness and wave reflection: a systematic review and meta-analysis of randomized controlled trials. *PLoS One.* 2014 Oct 15(10).
 Boutouyrie P, Lacolley P, Girerd X, Beck L, Safar M, Laurent S. Sympathetic activation decreases medium-sized arterial compliance in humans. *Am J Physiol.* 1994 Oct267(4 Pt 2):H1368-76.

	UFV ≤ 500 ml		500 < UFV ≤ 2000 ml		UFV > 2000 ml		All	
	In	Out	In	Out	In	Out	In	Out
pSBP (mmHg)	128.9	131.0	123.7	122.3	124.5	123.0	124.8	123.8
pDBP (mmHg)	75.9	74.5	74.9	72.3	75.6	73.1	75.3	* 72.9
pPP (mmHg)	53.0	56.5	48.8	50.0	478.8	49.9	49.4	50.9
HR (bpm)	69.4	70.2	68.1	70.8	71.8	73.7	69.8	* 71.9
cSBP (mmHg)	116.9	119.3	112.4	111.9	113.4	112.6	113.4	113.2
cDBP (mmHg)	77.3	75.9	76.3	73.6	77.2	74.6	76.8	* 74.3
cPP (mmHg)	39.6	*	43.4	38.3	36.2	38.0	36.6	** 38.9
Alx (%)	29.7	31.8	29.5	30.2	26.0	*	28.5	* 29.8
Alx75 (%)	26.4	28.8	25.5	27.7	24.0	**	27.6	** 27.8
PWV (m/s)	10.03	10.14	9.88	9.85	9.08	9.07	9.58	9.58

Table: Averaged hemodynamic parameters for intra- and interdialytic periods (In vs. Out) for different subgroups based on ultrafiltration volume (UF) and all subjects. Abbreviations: peripheral diastolic blood pressure (pDBP), peripheral systolic blood pressure (pSBP), peripheral pulse pressure (pPP), heart rate (HR), central diastolic blood pressure (cDBP), central systolic blood pressure (cSBP), peripheral pulse pressure (pPP), augmentation index (Alx, Alx75) and pulse wave velocity (PWV) */** marks a significant difference between intra- and interdialytic periods ($p < 0.05$ and $p < 0.01$, respectively).

References

1. Karpetas A, Sarafidis PA, Georgianos PI, Protogerou A, Vakianis P, Koutroumpas G, Raptis V, Stamatiadis DN, Syrganis C, Liakopoulos V, Efstratiadis G, Lasaridis AN. Ambulatory Recording of Wave Reflections and Arterial Stiffness during Intra- and Interdialytic Periods in Patients Treated with Dialysis. *Clin J Am Soc Nephrol.* 2015 10(4):630-638.

8.6 AEROBIC FITNESS LEVEL AND PERIPHERAL ARTERIAL COMPLIANCE – THE ROLE OF AUTONOMIC NERVOUS SYSTEM TONE

Nejka Potocnik, Ziva Melik, Ksenija Cankar, Martin Strucl
 Medical Faculty, Institute of Physiology, University of Ljubljana, Slovenia

Physical activity has beneficial effects on prevention of cardiovascular disease. Aerobic fitness is associated with higher central arterial

8.7 CHANGES IN CARDIAC FUNCTION BUT NOT STRUCTURE IN HEALTHY SUBJECTS WITH PREMATURE VASCULAR AGEING

Oscar Mac Ananey¹, Vincent Maher²

¹School of Biological Science, Dublin Institute of Technology, Dublin, Ireland

²Department of Cardiology, Tallaght Hospital, Dublin, Ireland

Purpose: Changes in myocardial and arterial wall properties/function are consistently reported in patients with established cardiovascular disease¹. However, few studies have reported these changes in early subclinical disease. The aim of the present study was to examine cardiac and vascular changes in early subclinical disease and to determine whether these changes occur in parallel.

Methods: For this study, 98 healthy lifelong never smokers were recruited. Subjects were categorised as having normal (Norm, n=71) or abnormal (High, n=27) arterial stiffness (carotid-femoral pulse wave velocity, PWV Vicorder, Skidmore, UK) for their age and blood pressure. M-mode Doppler echocardiography (Vivid 7 Dimension, GE, USA) was used to assess heart structure (interventricular septal thickness, IVSd left ventricular internal diameter, LVIDd left ventricular posterior wall thickness, LVPW left ventricular mass, LV Mass) and function (left ventricular isovolumetric relaxation time, LV IVRT mitral valve early/late filling velocity, MV E/A).