

## Artery Research Vol. **25(S1)**; 2019, *p.* S61 DOI: https://doi.org/10.2991/artres.k.191224.052; ISSN 1872-9312; eISSN 1876-4401 https://www.atlantis-press.com/journals/artres



## P18 Vascular Aging Index of the Finger Photoplethysmogram: A Validation Study with Vascular Stiffness, Mental Stress, and Day-to-Day Variability

Takashi Tarumi<sup>1,\*</sup>, Takayuki Yamabe<sup>2</sup>, Jun Sugawara<sup>1</sup>

<sup>1</sup>National Institute of Advanced Industrial Science and Technology, Tsukuba, Japan <sup>2</sup>University of Tsukuba, Tsukuba, Japan

## **ABSTRACT**

**Background:** Daily assessment of vascular health may predict cardiovascular incidence. Vascular aging index (VAI) calculated from second derivative of photoplethysmogram (SDPTG) is a simple, non-invasive measure possibly reflecting vascular stiffness. However, the effects of daily life events such as mental stress and day-to-day variability as well as its relation to other indices of vascular stiffness remain unclear.

**Purpose:** To determine whether VAI measured by finger SDPTG is 1) correlated with peripheral augmentation index (pAI), 2) altered by acute mental stress, and 3) affected by day-to-day variability.

**Methods:** Simultaneous measurements of finger photoplethysmogram and radial artery tonometry were performed in 68 healthy subjects (age = 22–64 years) of whom 31 subjects were further tested during a 30-second mental arithmetic and 10 subjects underwent day-to-day variability assessment for 5 consecutive days. VAI was calculated from a 20-second segment of photoplethysmogram data [1].

**Results:** At rest, VAI was positively correlated with pAI (r = 0.62, p < 0.001). During mental arithmetic, VAI increased when compared with the control condition (p = 0.032) whereas pAI did not show significant change. Five day measurements of VAI demonstrated the overall coefficient of variation of 21.1  $\pm$  13.7% across all subjects.

**Conclusion:** VAI calculated from the finger SDPTG is related to a measure of vascular stiffness and sensitive to mental stress with fair day-to-day variability. These findings suggest that VAI assessment needs to be performed at the quiet resting condition.

## REFERENCE

- [1] Takazawa K, Tanaka N, Fujita M, Matsuoka O, Saiki T, Aikawa M, et al. Assessment of vasoactive agents and vascular aging by the second derivative of photoplethysmogram waveform. Hypertension 1998;32:365–70.
  - © 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (http://creativecommons.org/licenses/by-nc/4.0/).