

Artery Research Vol. **25(S1)**; 2019, *p.* S124 DOI: https://doi.org/10.2991/artres.k.191224.111; ISSN 1872-9312; eISSN 1876-4401 https://www.atlantis-press.com/journals/artres



P81 Antibody to Cardiotonic Steroid Reduces Blood Pressure and Vascular Fibrosis in Preeclampsia

NI Agalakova³, VA Reznik¹, OV Nadei³, IA Ershov¹, OS Rassokha¹, ML Vasyutina², MM Galagudza², AY Bagrov^{1,3,*}

- ¹Department of Obstetrics and Gynecology, School of Pediatric Medicine
- ²Institute of Experimental Medicine, Almazov National Medical Research Centre
- ³Sechenov Institute of Evolutionary Physiology and Biochemistry, St. Petersburg, Russia

ABSTRACT

Background: Previous studies implicated cardiotonic steroids, including Na/K-ATPase inhibitor marinobufagenin (MBG), in the pathogenesis of preeclampsia (PE). We demonstrated that MBG induces fibrosis via mechanism involving inhibition of Fli1, a nuclear transcription factor and a negative regulator of collagen-1 synthesis.

Objectives and Methods: We hypothesized that PE blockade of increased MBG with antibody would lessen the fibrosis of umbilical arteries and lower the blood pressure in rats with PE. We tested 36 pregnant Sprague-Dawley rats in which 12 were made hypertensive by 1.8% Na supplementation (days 6–19 of gestation), 12 pregnant rats served controls. At day 19, PE rats received one intraperitoneal injection of polyclonal anti-MBG-4 antibody (0.5 ug/mL) for 4 hours.

Results: PE was associated with higher blood pressure (117 \pm 2 vs 107 \pm 2 mmHg; p < 0.01), plasma MBG levels (1.54 \pm 0.34 vs 0.49 \pm 0.11 nmol/L; p < 0.01), protein excretion (26 vs 12 mg/24 hours), sFlt-1 (4-fold), decrease in Fli1 (7-fold) and increase in collagen-1 in aorta (4-fold) vs. control rats (all p < 0.01). In 12 rats was treated with polyclonal anti-MBG-4 antibody blood pressure dropped (93 \pm 3 mmHg) and Fli1 was decreased much less (2-fold; p < 0.01 vs nontreated rats).

Conclusion: These results demonstrate that in experimental PE elevated MBG level is implicated in umbilical fibrosis via suppression of Fli1. Supported by Russian Scientific Foundation grant № 18-15-00222.

© 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (http://creativecommons.org/licenses/by-nc/4.0/).