



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P5.32: WAON-THERAPY (WARMING THERAPY) IMPROVED CARDIAC FUNCTION ASSOCIATING WITH DECREASING CARDIO-ANKLE VASCULAR INDEX (CAVI) IN HEART FAILURE PATIENTS

Kohji Shirai, Nagakawa Osamu, Hirano Keiichi, Noike Hirohumi, Takahashi Mao, Shimizu Kazuhiro, Yoshida Toyohiko

To cite this article: Kohji Shirai, Nagakawa Osamu, Hirano Keiichi, Noike Hirohumi, Takahashi Mao, Shimizu Kazuhiro, Yoshida Toyohiko (2013) P5.32: WAON-THERAPY (WARMING THERAPY) IMPROVED CARDIAC FUNCTION ASSOCIATING WITH DECREASING CARDIO-ANKLE VASCULAR INDEX (CAVI) IN HEART FAILURE PATIENTS, Artery Research 7:3_4, 152–152, DOI: <https://doi.org/10.1016/j.artres.2013.10.180>

To link to this article: <https://doi.org/10.1016/j.artres.2013.10.180>

Published online: 14 December 2019

Methods: Seventy-nine (51 male & 28 female) healthy subjects were recruited. All were lifelong non-smokers, normotensive, normolipidaemic and normoglycaemic. Augmentation index (Alx) and pulse wave velocity (PWV) were measured using applanation tonometry (Sphygmocor & Vicorder). Daily physical activity patterns were measured using triaxial accelerometry and expressed as time spent being sedentary, lightly active and vigorously active. Body composition was estimated using bioimpedance. The adipose associated markers, leptin & adiponectin, and inflammatory markers, IL-6, TNF α & MCP-1 were also measured.

Results: Time spent being sedentary was significantly associated with Alx ($r=0.38$, $P<0.001$), PWV ($r=0.33$, $P<0.01$), body fat composition ($r=0.40$, $P<0.001$) and age ($r=0.30$, $P<0.01$). Vigorous activity was inversely correlated with Alx ($r=-0.28$, $P<0.05$) body fat composition ($r=-0.30$, $P<0.01$), post prandial insulin ($r=-0.35$, $P<0.01$) and leptin/adiponectin ratio ($r=-0.28$, $P<0.05$). The inflammatory cytokines, IL-6, TNF α and MCP-1, were not significantly correlated with any activity categories.

Conclusion: The main findings of the study are that people, who spend more time being sedentary, despite being clinically healthy, have stiffer arteries and more body fat compared to those who spend less time being sedentary. Conversely, people that spend more time being active have less arterial stiffness, lower body fat, post prandial insulin and leptin/adiponectin ratio. These results suggest that activity related changes in arterial stiffness in healthy people may be mediated by the hormonal/metabolic consequences of inactivity rather than pro-inflammatory mechanisms.

P5.31

CENTRAL PRESSURES ARE HIGHER IN PATIENTS WITH TRANSIENT ISCHAEMIC ATTACK (TIA) THAN HEALTHY ADULTS

S. R. U. Patel¹, E. Bunting², E. Bodenham², S. A. Hopkins², E. Ormerod², J. Timeyin², K. Ali¹, C. Rajkumar¹

¹Brighton and Sussex Medical School, Brighton, United Kingdom

²Brighton and Sussex University Hospitals Trust, Brighton, United Kingdom

Introduction: The increased risk of stroke following a Transient Ischaemic Attack (TIA) is well established. Central aortic pressure, a marker of arterial stiffness, is emerging as a predictor of cardiovascular events including stroke. Additionally brachial Blood Pressure (BP) measurements may not accurately reflect the haemodynamic status of an individual, as systolic BP is amplified peripherally. The association and interplay of central aortic pressure and brachial BP parameters in TIA patients have not previously been explored.

Methods: Twenty-six participants with a mean age of 70 years ($SD\pm 10.1$) and a confirmed diagnosis of TIA were recruited from Brighton and Sussex University Hospitals Trust. Central aortic BP and brachial BP were measured using Arteriograph[®]. Carotid-femoral pulse wave velocity (C-F PWV) and carotid-radial pulse wave velocity (C-R PWV) were measured using Complior[®].

Results: Twenty participants had complete BP recordings. Central aortic BP was higher than peripheral BP (mean difference = 2 mmHg). Healthy patients are expected to have a central BP 10-15mmHg lower than peripheral BP.^[1,2] C-F PWV was found to be higher in TIA participants compared to published normal values (C-F 10.9 m/s).^[3]

Measurement	Mean \pm SD	Range	n
Peripheral BP (mmHg)	128 \pm 14.7	102-163	20
Central BP (mmHg)	130 \pm 19.3	97-171	20
C-F PWV (m/s)	11.5 \pm 3.18	7.2-20.6	22
C-R PWV (m/s)	10.8 \pm 1.26	8.3-13.7	22

Conclusions: In this TIA cohort, there was a tendency towards an increase in central BP and aortic stiffness in comparison to published normal values.

References

- McEnery CM, et al. Hypertension.2008 Jun;51(6):1476-82
- Williams B, et al. Circulation.2009 Jan 6;119(1):53-61
- Mattace-Raso F, et al. Eur Heart J. 2010 Oct;31(19):2338-50

P5.32

WAON-THERAPY (WARMING THERAPY) IMPROVED CARDIAC FUNCTION ASSOCIATING WITH DECREASING CARDIO-ANKLE VASCULAR INDEX (CAVI) IN HEART FAILURE PATIENTS

Kohji Shirai¹, Nagakawa Osamu¹, Hirano Keiichi², Noike Hirohumi², Takahashi Mao², Shimizu Kazuhiro², Yoshida Toyohiko¹

¹Mihama Hospital, Chiba, Japan

²Cardiovascular center, Sakura Hospital, Toho Univ., Chiba, Japan

Objects: Waon-Therapy was invented for the therapy of heart failure by Dr. Tei. The purpose of this study was to confirm the efficacy of Waon therapy and to clarify the role of Waon-therapy in the arterial stiffness monitoring with cardio-ankle vascular index (CAVI).

Subjects and methods: Study 1 Subjects were 7 patients with chronic heart failure. They were divided randomly into Waon-therapy group and conservative therapy group. Waon therapy was performed as follows; warming up at 60 for 15 min in a chamber, and then, laying down in supine position with covering warm blanket for 30 min. They took this therapy once a day for 14 days. Study 2. A person, 64 years old man with diabetes mellitus, who is taking hemodialysis therapy. He also was suffering from systemic arteriosclerosis and heart failure. He took Waon therapy for 3 months. CAVI was measured using Vasela1500 (Fukuda Denshi. Co.LTD).

Results: Study 1; Among 7 heart failure patients, Waon therapy group 4 patients showed improved BNP from 1220- 780 pg/ml, whereas conservative group 720- 920 pg/ml. CAVI improved in Waon group from 10.5 to 9.3, but in conservative group, CAVI did not change Study 2: 65 year old man taking maintenance hemodialysis were suffering from necrosis of penis because of arteriosclerosis. He took Waon therapy for 3 months. At every week, BNP decreased from 4200, 3400, 2600 to 1560 pg/ml after 3 months. CAVI also decreased 13.4, 13.2 12.3, 11.2, 10.8, 11.2. Penis necrosis was improved completely.

Conclusion: Waon therapy improved heart failure, associating with improvement of CAVI. Improved CAVI which means decreased arterial stiffness, and also decreased after-load, might be involved in improving left ventricular dysfunction.

P5.33

REVERSE DIPPING STATUS IS INVERSELY ASSOCIATED WITH HEALTH-RELATED QUALITY OF LIFE

D. A. Vrachatis^{1,2,3}, A. Argyris¹, A. Papadopoulos², T. G. Papaioannou³, E. Aissopou¹, G. Konstantonis¹, D. Niakas², P. Sfikakis¹, A. Protogerou¹

¹Cardiovascular Research Laboratory, First Dpt. of Propedeutic Medicine, National & Kapodistrian University of Athens, Athens, Greece

²Faculty of Social Sciences, Hellenic Open University, Patras, Greece

³Fist Dpt. of Cardiology, National & Kapodistrian University of Athens, Athens, Greece

Objectives: We sought to investigate whether reverse dipping status is associated with self-reported health-related quality of life (HRQL).

Methods: BP was evaluated non-invasively under 24-hour ABPM with a brachial cuff-based automatic oscillometric device (Mobil-O-Graph, IEM); all subjects maintained a time-log indicating awake/asleep periods. Subjects were classified as Reverse Dippers (Group-A) if SBP-asleep > SBP-awake; the rest were included in Group-B. For HRQL assessment the EQ5D instrument was utilized prior to BP evaluation. EQ5D estimates the following dimensions: mobility (MO), self-care (SC), usual activities (UA), pain/discomfort (PD), anxiety/depression (AD); each is subdivided in three severity levels (1=no, 2=some/moderate and 3=extreme problems). Moreover, an index (MVH-York-A1-tariff) is calculated by the combination of these dimensions. Additionally, EQ5D involves a visual analogue scale (VAS) in which respondents self-rate their health state (0=worst, 100=best imaginable).

Results: Group-A consisted of 10 subjects and Group-B of 139. The two groups were comparable regarding demographic and clinical characteristics (males 50% vs. 53%, 63 vs. 55 years old, BMI 26.5 vs. 27.2, hypertension 60% vs. 48.2%, diabetes mellitus 10% vs. 10%, respectively; $p=ns$ for all). MO, SC, UA, AD and MVH-York-A1-tariff (A: 0.74 ± 0.23 vs. B: 0.86 ± 0.74) did not differ between the two groups. Yet, a higher percentage of Group-A stated some/moderate or severe problems in PD dimension compared to Group-B (Figures A, B; $p<0.001$). Moreover, lower levels of VAS were observed in Group-A compared to Group-B (72.5 ± 10.9 vs. 78.7 ± 13.7 , respectively; $p=0.037$).

