

A New Stochastic Process with Long-Range Dependence

Sung Ik Kim^{1,*}, Young Shin Kim²

¹College of Business, Louisiana State University Shreveport, 1 University Place, Shreveport, LA 71115, USA

²College of Business, Stony Brook University, 100 Nicolls Road., Stony Brook, NY 11794, USA

ARTICLE INFO

Article History

Received 10 Jun 2020

Accepted 15 Sep 2020

Keywords

Generalized hyperbolic process
 Lévy process
 Time-changed Brownian motion
 Long-range dependence
 Fractional Brownian motion

ABSTRACT

In this paper, we introduce a fractional Generalized Hyperbolic process, a new stochastic process with long-range dependence obtained by subordinating fractional Brownian motion to a fractional Generalized Inverse Gaussian process. The basic properties and covariance structure between the elements of the processes are discussed, and we present numerical methods to generate the sample paths for the processes.

© 2020 The Authors. Published by Atlantis Press B.V.

This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).

1. INTRODUCTION

The fractional Brownian motion $\{B_H(t)\}_{t \geq 0}$ with Hurst parameter $H \in (0, 1)$ is a continuous zero mean Gaussian process with stationary increments and covariance function

$$\text{Cov}(B_H(t), B_H(s)) = \frac{1}{2} (|t|^{2H} + |s|^{2H} - |t - s|^{2H}), \quad \text{for } t, s \in \mathbb{R}.$$

For $H = 1/2$, the fractional Brownian motion is the same as ordinary Brownian motion which has independent increments. The fractional Brownian motion is first introduced by Mandelbrot and Van Ness [1] and has been widely used in many areas, such as theoretical physics, probability, hydrology, biology, finance, and many others, due to growing interest in the simulation of long-range dependence processes. In finance, especially, as a subclass of the fractional Stable process (See Samorodnitsky and Taqqu [2]), it has been applied to financial time series models having long-range dependence (See Willinger *et al.* [3], Lo [4], Cutland *et al.* [5]). Indeed, Kim [6] introduces the fractional multivariate Normal Tempered Stable process by using the time-changed fractional Brownian motion with the fractional Tempered Stable subordinator. Kim [7] redefines a fractional multivariate Normal Tempered Stable process and constructs new market model by applying the process to innovations on the multivariate ARMA-GARCH model.¹ Furthermore, Kim *et al.* [10] use a fractional Tempered Stable process in option pricing and compare its performance with that of the models using other types of stochastic processes. Other than Tempered Stable process, Meerschaert *et al.* [11] obtains fractional Laplace motion by subordinating fractional Brownian motion to a Gamma process to model hydraulic conductivity fields in geophysics, and Kozubowski *et al.* [12] applies it to modeling financial time series. Fractional Normal Inverse Gaussian process is also proposed as a simple alternative to the Normal Inverse Gaussian process with long-range dependence (See Kumar *et al.* [13], Kumar and Vellaisamy [14]).

Financial data have typically exhibited distinct nonperiodic cyclical patterns which are indicative of the presence of long-range dependence. In this paper, we introduce a fractional Generalized Hyperbolic process, a new stochastic process with the long-range dependence. The process is defined by taking the fractional Brownian motion that replaces the time variable to a fractional Generalized Inverse Gaussian process. It is noted that using the time-changed fractional Brownian motion with another long-range dependent stochastic process makes it possible to capture endogenous as well as exogenous long-range dependence (See Kim [6]). We discuss the basic properties of this process and obtain covariance structure between two elements of the processes from the covariance matrix of the fractional multivariate Brownian motion.

*Corresponding author. Email: sung.kim@lsus.edu

¹Please refer to Rachev and Mittnik [8], Sun *et al.* [9] for other applications of Stable process to the fractional models.

Our paper is organized as follows: In Section 2, we review Generalized Inverse Gaussian distribution. In Section 3, the fractional Generalized Inverse Gaussian process is defined, and its basic properties are discussed. In Section 4, we study the corresponding fractional univariate Generalized Hyperbolic process and discuss long-range dependence. Section 5 is devoted to the presentation of the fractional multivariate Generalized Hyperbolic process. In Section 6, we simulate the fractional Generalized Hyperbolic processes and illustrate sample paths for representative values of parameters. The principal findings are summarized in Section 7.

2. GENERALIZED INVERSE GAUSSIAN DISTRIBUTION

The class of Generalized Inverse Gaussian distribution which has been extensively studied by Jørgensen [15] is described by three parameters $(\lambda, \delta, \gamma)$. Its density function has support on the positive axis and is given by

$$f_{GIG}(x) = \frac{(\gamma/\delta)^\lambda}{2N_\lambda(\delta\gamma)} x^{\lambda-1} \exp\left(-\frac{1}{2}(\delta^2 x^{-1} + \gamma^2 x)\right), \quad x > 0, \quad (1)$$

where N_λ is the modified Bessel function of the second kind with index λ given by $N_\lambda(x) = \int_0^\infty u^{\lambda-1} e^{-\frac{1}{2}x(u^{-1}+u)} du$ for $x > 0$. The parameter domain of the Generalized Inverse Gaussian distribution is

$$\begin{aligned} \delta > 0, \quad \gamma \geq 0, & \quad \text{if } \lambda < 0, \\ \delta > 0, \quad \gamma > 0, & \quad \text{if } \lambda = 0, \\ \delta \geq 0, \quad \gamma > 0, & \quad \text{if } \lambda > 0. \end{aligned}$$

If $\lambda = -\frac{1}{2}$, the density function in Equation (1) reduces to that of the Inverse Gaussian distribution. The Gamma distribution is a limiting case of the Generalized Inverse Gaussian distribution for $\lambda > 0$ and $\gamma > 0$ and $\delta \rightarrow 0$. The mean and the variance of a Generalized Inverse Gaussian random variable G can easily be obtained from the Laplace transform. They are given, respectively, by $E[G] = \frac{\delta}{\gamma} \frac{N_{\lambda+1}(\delta\gamma)}{N_\lambda(\delta\gamma)}$ and $\text{Var}(G) = \frac{\delta^2}{\gamma^2} \left[\frac{N_{\lambda+2}(\delta\gamma)}{N_\lambda(\delta\gamma)} - \frac{N_{\lambda+1}^2(\delta\gamma)}{N_\lambda^2(\delta\gamma)} \right]$. Proposition 2.1 defines characteristic function of the Generalized Inverse Gaussian G .

Proposition 2.1. *The characteristic function of a Generalized Inverse Gaussian random variable G is given by*

$$\phi_G(u) = \left(\frac{\gamma}{\sqrt{\gamma^2 - 2iu}} \right)^\lambda \frac{N_\lambda(\delta\sqrt{\gamma^2 - 2iu})}{N_\lambda(\delta\gamma)}, \quad \delta, \gamma > 0.$$

Proof. Let $q(\lambda, \delta, \gamma) = \left(\frac{\gamma}{\delta}\right)^\lambda \frac{1}{2N_\lambda(\delta\gamma)}$ denote the norming constant of the Generalized Inverse Gaussian density, then the characteristic function of G is

$$\begin{aligned} \phi_G(u) &= E[e^{iuG}] = \int_0^\infty q(\lambda, \delta, \gamma) x^{\lambda-1} \exp\left(-\frac{1}{2}(\delta^2 x^{-1} + (\gamma^2 - 2iu)x)\right) dx \\ &= \frac{q(\lambda, \delta, \gamma)}{q(\lambda, \delta, \sqrt{\gamma^2 - 2iu})} = \left(\frac{\gamma}{\sqrt{\gamma^2 - 2iu}} \right)^\lambda \frac{N_\lambda(\delta\sqrt{\gamma^2 - 2iu})}{N_\lambda(\delta\gamma)}. \end{aligned}$$

□

Since Generalized Inverse Gaussian distribution is infinitely divisible, we can define one Lévy process $\{G(t)\}_{t \geq 0}$ such that the characteristic function of $G(t)$ is given by $\phi_{G(t)}(u) = E[\exp(iuG(t))] = \exp(t \log(\phi_G(u)))$, where $\phi_G(u)$ is given by Proposition 2.1. In this case, $\{G(t)\}_{t \geq 0}$ is referred to as *Generalized Inverse Gaussian process* with parameters $(\lambda, \delta, \gamma)$.

3. FRACTIONAL GENERALIZED INVERSE GAUSSIAN PROCESS

To define a fractional Generalized Inverse Gaussian process, we use the Volterra kernel $K_H : [0, \infty] \times [0, \infty] \rightarrow [0, \infty]$, given by

$$K_H(t, s) = c_H \left(\left(\frac{t}{s}\right)^{H-\frac{1}{2}} (t-s)^{H-\frac{1}{2}} - \left(H - \frac{1}{2}\right) s^{\frac{1}{2}-H} \int_s^t u^{H-\frac{3}{2}} (u-s)^{H-\frac{1}{2}} du \right) \mathbb{1}_{[0,t]}(s) \quad (2)$$

with

$$c_H = \left(\frac{H(1-2H)\Gamma(\frac{1}{2}-H)}{\Gamma(2-2H)\Gamma(H+\frac{1}{2})} \right)^{\frac{1}{2}} \quad \text{and} \quad H \in (0, 1).$$

According to Houdre and Kawai [16] and Nualart [17], we have the following facts:

1. For $t, s > 0$, $\int_0^{t \wedge s} K_H(t, u)K_H(s, u)du = \frac{1}{2} (t^{2H} + s^{2H} - |t - s|^{2H})$ and $\int_0^t K_H(t, s)^2 ds = t^{2H}$.
2. If $H \in (\frac{1}{2}, 1)$, then $K_H(t, s) = c_H (H - \frac{1}{2}) s^{\frac{1}{2}-H} \int_s^t (u - s)^{H-\frac{3}{2}} u^{H-\frac{1}{2}} du \mathbb{1}_{[0,t]}(s)$.
3. Let $t > 0$ and $p \geq 2$. $K_H(t, \cdot) \in L^p([0, t])$ if and only if $H \in (\frac{1}{2} - \frac{1}{p}, \frac{1}{2} + \frac{1}{p})$. When $K_H(t, \cdot) \in L^p([0, t])$, $\int_0^t K_H(t, s)^p ds = C_{H,p} t^{p(H-\frac{1}{2})+1}$, where $C_{H,p} = c_H^p \int_0^1 v^{p(\frac{1}{2}-H)} \left[(1-v)^{H-\frac{1}{2}} - (H-\frac{1}{2}) \int_v^1 w^{H-\frac{3}{2}}(w-v)^{H-\frac{1}{2}} dw \right]^p dv$.

Let $H \in (0, 1)$, and consider a fractional Lévy process $G_H = \{G_H(t)\}_{t \geq 0}$, which is $G_H(t) = \int_0^t K_H(t, u) dG(u)$, where $\{G(t)\}_{t \geq 0}$ is the Generalized Inverse Gaussian process and K_H is the Volterra kernel defined in Equation (2). The process G_H is referred to as the fractional Generalized Inverse Gaussian process with parameters $(H, \lambda, \delta, \gamma)$. We first describe the covariance structure of fractional Generalized Inverse Gaussian process as the Proposition 3.1 without proof.²

Proposition 3.1. *Let $t, s \geq 0$. The covariance between $G_H(s)$ and $G_H(t)$ is given by*

$$\text{Cov}(G_H(s), G_H(t)) = \frac{1}{2} \frac{\delta^2 N_{\lambda+2}(\delta\gamma)}{\gamma^2 N_{\lambda}(\delta\gamma)} (t^{2H} + s^{2H} - |t - s|^{2H}). \tag{3}$$

The characteristic function of fractional Generalized Inverse Gaussian process and cumulants are obtained by the Propositions 3.2 and 3.3, respectively.

Proposition 3.2. *With $H \in (0, 1)$, the characteristic function of $G_H(t)$ is given by*

$$\phi_{G_H(t)}(z) = E[\exp(izG_H(t))] = \exp \left(\int_0^t \psi_G(zK_H(t, u)) du \right), \quad \text{where} \quad \psi_G(u) = \log \phi_{G(1)}(u).$$

Proof. For $0 = t_0 < t_1 < \dots < t_M = t$ and $\Delta t = t_1 - t_0 = t_2 - t_1 = \dots = t_M - t_{M-1}$,

$$E[\exp(izG_H(t))] = E \left[\exp \left(iz \int_0^t K_H(t, u) dG(u) \right) \right] = \lim_{\Delta t \rightarrow 0} \prod_{j=0}^{M-1} E[\exp(izK(t, t_j)\Delta G(t_j))]. \tag{4}$$

For all $t \geq 0$, $\Delta G(t_j)$ can be represented as $\Delta G(t_j) \stackrel{d}{=} \Delta t G(1)$. Therefore, from Equation (4) the characteristic function of $G_H(t)$ is

$$\phi_{G_H(t)}(z) = \lim_{\Delta t \rightarrow 0} \prod_{j=0}^{M-1} \exp(\Delta t \psi_G(zK_H(t, t_j))) = \exp \left(\int_0^t \psi_G(zK_H(t, u)) du \right).$$

□

Proposition 3.3. *Let $n \in \mathbb{N}$. The cumulant $c_n(G_H(t))$ of $G_H(t)$ is given by*

$$c_n(G_H(t)) = \frac{1}{i^n} \frac{\partial^n}{\partial z^n} \log \phi_{G_H(t)}(z)|_{z=0} = \frac{1}{i^n} \psi_G^{(n)}(0) \int_0^t (K_H(t, u))^n du.$$

Therefore, by the cumulant $c_n(G_H(t))$ from Proposition 3.3, the mean and variance of $G_H(t)$ can be obtained, respectively, by

$$E[G_H(t)] = \frac{1}{i} \psi'_G(0) \int_0^t K_H(t, u) du = E[G] C_{H,1} t^{H+\frac{1}{2}} = \frac{\delta N_{\lambda+1}(\delta\gamma)}{\gamma N_{\lambda}(\delta\gamma)} C_{H,1} t^{H+\frac{1}{2}} \quad \text{and}$$

² Refer to Proposition 3.1 in Houdre and Kawai [16].

$$\text{Var}(G_H(t)) = -\psi_G''(0) \int_0^t (K_H(t, u))^2 du = \text{Var}(G) C_{H,2} t^{2H} = \frac{\delta^2}{\gamma^2} \left[\frac{N_{\lambda+2}(\delta\gamma)}{N_\lambda(\delta\gamma)} - \frac{N_{\lambda+1}^2(\delta\gamma)}{N_\lambda^2(\delta\gamma)} \right] C_{H,2} t^{2H}.$$

4. FRACTIONAL UNIVARIATE GENERALIZED HYPERBOLIC PROCESS

Assume that $\{B_H(t)\}_{t \geq 0}$ the univariate fractional Brownian motion with Hurst parameter $H \in (0, 1)$ is given by $B_H(t) = \int_0^t K_H(t, s) dB(s)$, where $\{B(t)\}_{t \geq 0}$ is a standard Brownian motion, and K_H is a Volterra kernel defined in Equation (2). Let $\{B_{H_1}(t)\}_{t \geq 0}$ be the fractional Brownian motion with Hurst parameter $H_1 \in (0, 1)$ and $\{G_{H_2}(t)\}_{t \geq 0}$ be the univariate fractional Generalized Inverse Gaussian process with parameters $(H_2, \lambda, \delta, \gamma)$. Suppose that $\{B_{H_1}(t)\}_{t \geq 0}$ and $\{G_{H_2}(t)\}_{t \geq 0}$ are independent. A process $X = \{X(t)\}_{t \geq 0}$ defined by $X(t) = \beta(G_{H_2}(t))^{2H_1} + B_{H_1}(G_{H_2}(t))$, where $\beta \in \mathbb{R}$, is referred to as the fractional univariate Generalized Hyperbolic process. The characteristic function of $X(t)$ is $\phi_{X(t)}(z) = \phi_{(G_{H_2}(t))^{2H_1}} \left(\beta z + \frac{iz^2}{2} \right)$, where $\phi_{(G_{H_2}(t))^{2H_1}}$ is the characteristic function of $(G_{H_2}(t))^{2H_1}$. Since $\phi_{(G_{H_2}(t))^{2H_1}}$ does not have a general closed form for all $H_1 \in (0, 1)$, we consider $H_1 = \frac{1}{2}$. Then, we have

$$\phi_{X(t)}(z) = \exp \left(\int_0^t \psi_G \left(\left(\beta z + \frac{iz^2}{2} \right) K_{H_2}(t, u) \right) du \right). \tag{5}$$

The mean of $X(t)$, $E[X(t)] = \beta E[(G_{H_2}(t))^{2H_1}]$. If $H_1 = \frac{1}{2}$, $E[X(t)] = \beta E[G_{H_2}(t)] = \frac{\beta \delta}{\gamma} \frac{N_{\lambda+1}(\delta\gamma)}{N_\lambda(\delta\gamma)} C_{H_2,1} t^{H_2 + \frac{1}{2}}$.

For $0 \leq s \leq t$, we have

$$E[B_{H_1}(G_{H_2}(s))B_{H_1}(G_{H_2}(t))] = \frac{1}{2} E \left[(G_{H_2}(t))^{2H_1} + (G_{H_2}(s))^{2H_1} - (G_{H_2}(t) - G_{H_2}(s))^{2H_1} \right],$$

so that the covariance between $X(s)$ and $X(t)$ is given by

$$\text{Cov}(X(s), X(t)) = \beta^2 \text{Cov}((G_{H_2}(s))^{2H_1}, (G_{H_2}(t))^{2H_1}) + \frac{1}{2} E \left[(G_{H_2}(t))^{2H_1} + (G_{H_2}(s))^{2H_1} - (G_{H_2}(t) - G_{H_2}(s))^{2H_1} \right].$$

In the case of $H_1 = \frac{1}{2}$, we obtain by Proposition (3.1)

$$\text{Cov}(X(s), X(t)) = \frac{1}{2} \frac{\beta^2 \delta^2}{\gamma^2} \frac{N_{\lambda+2}(\delta\gamma)}{N_\lambda(\delta\gamma)} \left(t^{2H_2} + s^{2H_2} - (t-s)^{2H_2} \right) + \frac{\delta}{\gamma} \frac{N_{\lambda+1}(\delta\gamma)}{N_\lambda(\delta\gamma)} C_{H_2,1} s^{H_2 + \frac{1}{2}}.$$

The fractional Generalized Hyperbolic process $X(t)$ defined as above can be applied to stock price process with the objective of valuing option. Suppose that under the risk neutral measure \mathbb{Q} , the stock price process $\{S(t)\}_{t \geq 0}$ is given by

$$S(t) = \frac{S(0) \exp(rt + X(t))}{E_{\mathbb{Q}}[\exp(X(t))]},$$

where r is the risk-free short rate. Then, by the inverse Fourier transform method in Carr and Madan [18] and Lewis [19], the call option price with time to maturity T and strike price K is

$$\frac{K^{1+\rho} e^{-rT}}{\pi S(0)^\rho} \text{Re} \int_0^\infty e^{-iu \log(K/S(0))} \frac{e^{(iu-\rho)T} \phi_{X(T)}(u + i\rho)}{(\rho - iu)(1 + \rho - iu)(\phi_{X(T)}(-i))^{iu-\rho}} du,$$

where ρ is real number such that $\rho < -1$, and the characteristic function $\phi_{X(T)}(\cdot)$ is defined on Equation (5). The put option price can be obtained by the same formula under the condition of $\rho > 0$.

5. FRACTIONAL MULTIVARIATE GENERALIZED HYPERBOLIC PROCESS

Consider a multivariate fractional Brownian motion $B_{H_1} = \{B_{H_1}(t)\}_{t \geq 0}$ such that $B_{H_1}(t) = (B_{H_1,1}(t), B_{H_1,2}(t), \dots, B_{H_1,N}(t))^T$, and suppose that

$$\text{Cov}(B_{H_1,m}(t), B_{H_1,n}(t)) = \sigma_{m,n} t^{2H_1}$$

for all $m, n \in \{1, 2, \dots, N\}$. Let Σ be the covariance matrix for $B_{H_1}(1)$, which is $\Sigma = [\sigma_{m,n}]_{m,n \in \{1,2,\dots,N\}}$, and $G_{H_2} = \{G_{H_2}(t)\}_{t \geq 0}$ be the univariate fractional Generalized Inverse Gaussian process with parameters $(H_2, \lambda, \delta, \gamma)$. Suppose that G_{H_2} is independent of B_{H_1} . Let $X = \{X(t)\}_{t \geq 0}$ with $X(t) = (X_1(t), X_2(t), \dots, X_N(t))^T$ be a process of the random vector defined by

$$X(t) = (G_{H_2}(t))^{2H_1} \beta + B_{H_1}(G_{H_2}(t)), \tag{6}$$

where $\beta = (\beta_1, \beta_2, \dots, \beta_N)^T \in \mathbb{R}^N$. Then, X is referred to as the fractional multivariate Generalized Hyperbolic process. The characteristic function of $X(t)$ is given by

$$\phi_{X(t)}(z) = E[\exp(iz^T X(t))] = \phi_{(G_{H_2}(t))^{2H_1}} \left(\beta^T z + \frac{i}{2} z^T \Sigma z \right),$$

where $z = (z_1, z_2, \dots, z_N)^T \in \mathbb{R}^N$ and $\phi_{(G_{H_2}(t))^{2H_1}}$ is the characteristic function of $(G_{H_2}(t))^{2H_1}$. Let $t \geq 0$, $m, n \in \{1, 2, \dots, N\}$, and $X_m(t)$ and $X_n(t)$ be the m -th and n -th elements of the vector $X(t)$, respectively. Then, the covariance between $X_m(t)$ and $X_n(t)$ is given by

$$\text{Cov}(X_m(t), X_n(t)) = \beta_m \beta_n \text{Var} \left((G_{H_2}(t))^{2H_1} \right) + \sigma_{m,n} E \left[(G_{H_2}(t))^{2H_1} \right].$$

If $H_1 = \frac{1}{2}$, we have

$$\phi_{X(t)}(z) = \exp \left(\int_0^t \psi_G \left(\left(\beta^T z + \frac{i}{2} z^T \Sigma z \right) K_{H_2}(t, u) \right) du \right) \quad \text{and}$$

$$\text{Cov}(X_m(t), X_n(t)) = \beta_m \beta_n \frac{\delta^2}{\gamma^2} \left[\frac{N_{\lambda+2}(\delta\gamma)}{N_\lambda(\delta\gamma)} - \frac{N_{\lambda+1}^2(\delta\gamma)}{N_\lambda^2(\delta\gamma)} \right] C_{H_2,2} t^{2H_2} + \sigma_{m,n} \frac{\delta}{\gamma} \frac{N_{\lambda+1}(\delta\gamma)}{N_\lambda(\delta\gamma)} C_{H_2,1} t^{H_2+\frac{1}{2}}.$$

6. SIMULATION

In this section, the sample paths of the fractional Generalized Hyperbolic processes are simulated by subordinating a discretized fractional Generalized Inverse Gaussian process with fractional Brownian motion on equally spaced intervals. We simulate $G_{H_2}(t)$ as follows:

1. Choose M fixed times in $[0, t]$: $t_0 = 0, t_1 = t/M, \dots, t_{M-1} = (M-1)t/M$, and $t_M = t$.
2. Generate M Generalized Inverse Gaussian variates $(G(t_1), G(t_2), \dots, G(t_M))$.
3. Generate $G_{H_2}(t)$ using $G_{H_2}(t) = \lim_{M \rightarrow \infty} \sum_{j=1}^M K_{H_2}(t, t_{j-1})(G(t_j) - G(t_{j-1}))$.

Let L_Σ be the lower triangular matrix obtained by the Cholesky decomposition for Σ with $\Sigma = L_\Sigma L_\Sigma^T$, where Σ is the correlation matrix in Equation (6). Then, we have $B_{H_1}(t) = L_\Sigma \bar{B}_{H_1}(t)$, where $\bar{B}_{H_1}(t) = (\bar{B}_{H_1,1}(t), \bar{B}_{H_1,2}(t), \dots, \bar{B}_{H_1,N}(t))^T$ is a mutually independent vector of fractional Brownian motions. For a given partition in 1) above and $t_j < t_k$ for $j < k$, we have

$$\bar{B}_{H_1,n}(G_{H_2}(t_j)) - \bar{B}_{H_1,n}(G_{H_2}(t_{j-1})) = (G_{H_2}(t_j) - G_{H_2}(t_{j-1}))^{H_1} (\bar{B}_{H_1,n}(t_j) - \bar{B}_{H_1,n}(t_{j-1})),$$

where $n \in \{1, 2, \dots, N\}$. Therefore,

$$X(t_k) = \beta (G_{H_2}(t_k))^{2H_1} + \sum_{j=1}^k (G_{H_2}(t_j) - G_{H_2}(t_{j-1}))^{H_1} L_\Sigma \tilde{B}_j, \tag{7}$$

where $\beta = (\beta_1, \beta_2, \dots, \beta_N)^T$ and $\tilde{B}_j = (\bar{B}_{H_1,1}(t_j) - \bar{B}_{H_1,1}(t_{j-1}), \bar{B}_{H_1,2}(t_j) - \bar{B}_{H_1,2}(t_{j-1}), \dots, \bar{B}_{H_1,N}(t_j) - \bar{B}_{H_1,N}(t_{j-1}))^T$.

Figure 1 illustrates the simulated sample paths from a Generalized Inverse Gaussian process with $M = 250$ and parameters $\lambda = -1.2$, $\delta = 0.1$, and $\gamma = 0.01$. The GIG process on panel (a) depicts a sample path from the Generalized Inverse Gaussian process. Then, obtained are the fGIGs, the fractional Generalized Inverse Gaussian processes, with $H = 0.70$ and 0.90 from the GIG process. Panel (b) gives the simulated sample paths of the univariate fractional Generalized Hyperbolic processes with $\beta = -0.05$, comparing with the path of the nonfractional Generalized Hyperbolic process. Notice that the sample paths for the fractional processes have less fluctuations than the path for nonfractional process, indicating the persistence of long-range dependence property of the fractional Generalized Hyperbolic processes. In panel (c), the two-dimensional plot for the bivariate fractional Generalized Hyperbolic process is presented using Equation (7) with $H_1 = 0.55, H_2 = 0.80, \beta = (-0.05, -0.03)^T$, and $\text{Cov}(B_{H_1,1}(1), B_{H_1,2}(1)) = 0.75$.

7. CONCLUDING REMARKS

In this paper, a fractional Generalized Hyperbolic process defined by the time-changed fractional Brownian motion with the fractional Generalized Inverse Gaussian process is presented. This process is featured by the capability to capture not only the endogenous long-range dependence by the fractional Brownian motion, but also the exogenous long-range dependence by the fractional Generalized Inverse Gaussian process. That is, the process could implement the long-range dependence in volatility as well as the long-range dependence in random process itself.




Figure 1 | Sample Paths from Simulations (a) Sample Paths for (Fractional) Generalized Inverse Gaussian Processes (b) Sample Paths for (Fractional) Generalized Hyperbolic Processes (c) Sample Path for Fractional Bivariate Generalized Hyperbolic Process.

CONFLICTS OF INTEREST

All authors report no conflicts of interest relevant to this article.

AUTHORS' CONTRIBUTIONS

We thank the anonymous reviewers for helpful comments on earlier drafts of the manuscript.

REFERENCES

1. B.B. Mandelbrot, J.W. Van Ness, *SIAM Rev.* 10 (1968), 422–437.
2. G. Samorodnitsky, M.S. Taqqu, *Stable Non-Gaussian Random Processes*, Chapman & Hall, CRC, New York, USA, 1994.
3. W. Willinger, M.S. Taqqu, V. Teverovsky, *Finance Stoch.* 3 (1999), 1–13.
4. A.W. Lo, *Econometrica.* 59 (1991), 1279–1313.
5. N.G. Cutland, P.E. Kopp, P.E. Willinger, in: E. Bolthausen, M. Dozzi, F. Russo (Eds.), *Seminar on Stochastic Analysis, Random Fields and Applications*, Progress in Probability, vol 36, Birkhäuser, Basel, Switzerland, 1995, pp. 327–351.
6. Y.S. Kim, *Appl. Math. Lett.* 25 (2012), 2396–2401.
7. Y.S. Kim, *Front. Appl. Math. Stat.* 1 (2015), 1–12.
8. S.T. Rachev, S. Mittnik, *Stable Paretian Models in Finance*, John Wiley & Sons, New York, NY, USA, 2000.
9. W. Sun, S.T. Rachev, F. Fabozzi, in: D. Seese, C. Weinhardt, F. Schlottmann (Eds.), *Handbook on Information Technology in Finance*, International Handbooks Information System, Springer, Berlin, Heidelberg, Germany, 2008.
10. Y.S. Kim, D. Jiang, S. Stoyanov, *J. Deriv.* 26 (2019), 71–88.
11. M.M. Meerschaert, T.J. Kozubowski, F.J. Molz, S. Lu, *Geophys. Res. Lett.* 31 (2004), 1–4.
12. T.J. Kozubowski, M.M. Meerschaert, K. Podgorski, *Adv. Appl. Probab.* 38 (2006), 451–464.
13. A. Kumar, M.M. Meerschaert, P. Vellaisamy, *Stat. Probab. Lett.* 81 (2011), 146–152.
14. A. Kumar, P. Vellaisamy, *Methodol. Comput. Appl. Probab.* 14 (2012), 263–283.

15. B. Jørgensen, *Statistical Properties of the Generalized Inverse Gaussian Distribution*, Lecture Notes in Statistics, Springer, New York, NY, USA, 1982.
16. C. Houdre, R. Kawai, *Stoch. Process Their Appl.* 116 (2006), 1161–1184.
17. D. Nualart, *Contemp. Math.* 336 (2003), 3–39.
18. P. Carr, D. Madan, *J. Comput. Finance.* 2 (1999), 61–73.
19. A.L. Lewis, *A Simple Option Formula for General Jump-Diffusion and Other Exponential Lévy Processes*, 2001.
<http://www.optioncity.net/pubs/ExpLevy.pdf>