



Research Article

Factors Associated with Cardio Ankle Vascular Index (CAVI) and its Mathematically Corrected Formula (CAVI₀) in Community Dwelling Individuals

Li-Ju Lin¹, Nan-Chen Hsieh², Shu-Fang Vivienne Wu³, Tan-Hsu Tan⁴ , Azadeh Alizargar⁴,
Chyi-Huey Bai⁵, Javad Alizargar⁶ 

¹Department of Gerontological Health Care, National Taipei University of Nursing and Health Sciences, Taipei City 112, Taiwan

²Department of Information Management, National Taipei University of Nursing and Health Sciences, Taipei City 112, Taiwan

³College of Nursing, School of Nursing, National Taipei University of Nursing and Health Sciences, Taipei City 112, Taiwan

⁴Department of Electrical Engineering, National Taipei University of Technology, Taipei City 10608, Taiwan

⁵Department of Public Health, College of Medicine, Taipei Medical University, Taipei, Taiwan

⁶Research Center for Healthcare Industry Innovation, National Taipei University of Nursing and Health Sciences, Taipei City 112, Taiwan

ARTICLE INFO

Article History

Received 26 May 2020

Accepted 09 November 2020

Keywords

Atherosclerosis
cardiovascular diseases
cardio ankle vascular index
ankle brachial index
hypertension
atherosclerosis
risk factor

ABSTRACT

Cardio Ankle Vascular Index (CAVI) and mathematically corrected formula derived from it (CAVI₀) are indices for arterial stiffness and atherosclerosis. Role of different atherosclerotic risk factors on CAVI and CAVI₀ is not clear in the community dwelling individuals. This study aims to evaluate the association of different Cardiovascular Disease (CVD) risk factors on CAVI and CAVI₀. Participants from a prospective cohort study have been recruited for CAVI measurement. Known risk factors for atherosclerosis were assessed in the individuals. CAVI and CAVI₀ is highly correlated (Pearson's $r = 0.95$ and $p < 0.001$). Further correlation analysis of the study variables with CAVI shows that CAVI is significantly correlated with age, Systolic Blood Pressure (SBP), Diastolic Blood Pressure (DBP), Mean Blood Pressure (MBP). CAVI₀ is also significantly correlated with age, SBP and MBP. CAVI₀ is not correlated with DBP (Pearson's $r = 0.05$ and $p = 0.46$). Among laboratory tests, CAVI and CAVI₀ are correlated with Blood Urea Nitrogen (BUN) ($r = 0.20$, $p = 0.009$ and $r = 0.22$, $p = 0.004$ respectively) and also with HBA1c ($r = 0.21$, $p = 0.006$ and $r = 0.25$, $p = 0.001$ respectively). Multivariate analysis showed that only age is independent determinant of CAVI and CAVI₀. As SBP and DBP are not independently associated with CAVI and CAVI₀, these indices could be considered reliable in differentiating people with high risks of CVD, as it is not dependent on other risk factors. As CAVI₀ is not correlated with DBP, when evaluating the results of CAVI₀, we should consider that this index maybe not be reflective of DBP in the patient.

© 2020 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).

1. INTRODUCTION

Cardiovascular Disease (CVD) like Coronary Heart Disease (CHD) are leading cause of death in the world [1]. Screening to find sub-clinical damages due to CVD has been suggested by European Society of Hypertension and cardiology guidelines, because classical risk factors fail to predict it [2]. Arterial stiffness is one of these suggested parameters for screening subclinical organ damage [3]. Arterial stiffness can be calculated by different indexes. One of the most known indexes is carotid-femoral Pulse Wave Velocity (cfPWV), and that is the velocity of the arterial pulse moving along the vessel wall [4]. Although this method is the most frequently used method in evaluating arterial stiffness, it is dependent on current Blood Pressure (BP) and short term changes in BP [5]. A novel index has been introduced named Cardio Ankle Vascular Index (CAVI), solving this issue as it is not dependent on the current BP [6,7].

*Corresponding author. Email: jaz.tmu@gmail.com

Peer review under responsibility of the Association for Research into Arterial Structure and Physiology

Data availability statement: The data that support the findings of this study are available from the corresponding author [JA], upon reasonable request.

Although it is said that blood pressure has no independent effect on CAVI values, some studies report contradicting results [8]. Aside from blood pressure, other factors such as age, body compositions, history of CVD and Ischemic Heart Disease (IHD), Type 2 Diabetes Mellitus (T2DM), smoking, alcohol drinking, exercise and anxiety have been evaluated in this study. These factors have been suggested to be associated with arterial stiffness and atherosclerosis [9–12]. Lipid profile and fasting blood sugar also have been shown to influence CAVI and yet have not been agreed upon [13]. Some laboratory tests like high sensitivity C-reactive protein (hs-CRP), Blood Urea Nitrogen (BUN), Creatinine (Cr), serum Uric Acid (UA) and Fasting Blood Sugar (FBS) can also have some effects on CAVI measurement [14]. The question of independent association of these factors with CAVI has not been addressed in details so far.

A mathematically corrected formula derived from CAVI was proposed by Spronck et al. [15,16] they also provided a tool to easily convert CAVI into CAVI₀ values. They claimed that CAVI₀ is less dependent on blood pressure changes. Studies to compare CAVI and CAVI₀ at the same time to observe the difference between associating factors can shed light on this matter.

This study primarily tries to answer the question of which one of the atherosclerosis and arterial stiffness risk factors has the most independent effect on CAVI and $CAVI_0$ measurement. Our secondary objective is to try to find out the factors that should be considered if we are dealing with using CAVI and $CAVI_0$ in clinical practice. We also try to understand what happens if we use $CAVI_0$ as our primary measure instead of CAVI. Sex differences in association of serum levels of study variables with CAVI is also considered in this study. A study on community dwelling individuals without high risk of CVD may help us better understand the independent role of mentioned risk factors and if we find an appropriate model, distinguish between the people who may benefit the measurement of CAVI and $CAVI_0$ in clinical practice.

2. MATERIALS AND METHODS

Participants were recruited as a part of annual checks of a community-based prospective cohort study on the community dwelling individuals of six villages in Shihlin District, and six villages in Wenshan District, Taipei, Taiwan (the coverage of Shin Kong Wu Ho-Su Memorial Hospital and Wan Fang Hospital respectively) that was designed to evaluate the CV and cerebrovascular risk factors in individuals. They were excluded if age ≤ 30 , had an incomplete questionnaire, prior history of cancer, chronic kidney disease, refused to blood draw or perform CAVI measurement. Data of 198 individuals were collected. All subjects gave their informed consent for inclusion before they participated in the study. The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Ethics Committee of Taipei Medical University with the Institute Review Board (reference numbers 94E-183, 94E-198 and 96E-004). We excluded one individual that had prior history of admission due to stroke and nine that had admission due to CVD. Twenty two individuals that had positive hepatitis B infection and seven that had hepatitis C infection were also excluded. To exclude the people with high risk of atherosclerosis, we removed two cases with Ankle Brachial Index (ABI) > 1.3 . Final data of 163 individuals were analyzed.

All subjects' demographic data were gathered on age, sex, body height and weight, history of IHD, T2DM, Family History of heart disease (FH) and Hypertension (HTN). IHD and T2DM was based on the physician diagnosis. FH is positive if there is a history of heart disease in brother or father < 60 years old and in sister or mother before 65 years of age. Smoking history was also gathered, whether they ever smoked or not, and if so the number of cigarettes per day and the duration of smoking. Pack-years of smoking was calculated by multiplying the packs of cigarettes per day by years of smoking.

Habit of alcohol drinking and the volume (cc) of alcohol drinks per week was asked. Participants were asked whether they have the habit of exercising and the frequency of exercising per week was also questioned. Fasting blood was taken and sent to the laboratory for determination of serum levels of BUN, Cr, AST, ALT, FBS, UA, hs-CRP, High Density Lipoprotein (HDL), Low Density Lipoprotein (LDL) and Triglyceride (TGL) using standard methods (using an X-1500-Sysmex, Germany; Beckman AU5800, USA; and Tosoh HLC-723G8 automated glycol-hemoglobin analyzer, Japan). Evaluation of anxiety was carried out using the standard General Anxiety Disorder 7 questionnaire (GAD-7). Every question scores [range 0 (not at all) to 3 (Nearly every day)]. Overall score of 5, 10 and 15 were used to categorize mild, moderate and severe cases of anxiety.

Participants were asked to lie down in supine position for 10 min and electrocardiogram (ECG) and phonocardiogram (PCG) were performed on them. Pulse wave velocity (PWV) was measured by VaSera VS-1000 (Fukuda Denshi, Tokyo, Japan) [17]. All the CAVI measurement was done intrinsically by VS-1000. CAVI is measured by the following formula:

$$CAVI = a \left[\left(\frac{2\rho}{\Delta P} \right) * \ln \left(\frac{SBP}{DBP} \right) PWV^2 \right] + b$$

$CAVI_0$ is calculated by the conversion tool that Spronck et al. [16] suggested from CAVI. The $CAVI_0$ has been obtained from the following formula:

$$CAVI_0 = 2\rho \times \left(\frac{PWV^2}{DBP} \right) - \ln \left(\frac{DBP}{P_0} \right)$$

SBP and DBP are systolic and diastolic blood pressure. $\Delta P = SBP - DBP$ and ρ is the blood density. a and b are the constants automatically measured by the device to match aortic PWV. P_0 is the reference pressure (100 mmHg).

All study variables were analyzed by ANOVA and Fisher's exact test stratified by CAVI levels. CAVI was analyzed for normal distribution test (Shapiro–Wilk test) before performing a multiple regression analysis. Pearson's correlation analysis performed on the study variables and 95% prediction eclipse of this study variables was drawn based on the study objectives to show the relationship between the variables. Linear regression model was used to evaluate the independency of associations of selected correlated factors with CAVI and $CAVI_0$. CAVI and $CAVI_0$ were served as the dependent variable in the linear regression model. All of the parameters with significant correlation with CAVI and $CAVI_0$ was selected to be put in the models and different models were selected base on the objectives of the study. Logistic regression models were used to determine high CAVI and $CAVI_0$. Odds ratio and Confidence Interval (CI) was calculated afterwards. All statistical analysis was performed using SAS 9.4. (SAS, Cary, NC, USA).

3. RESULTS

Data of 163 community individuals were analyzed. The mean age \pm Standard Deviation (SD) was 63.06 ± 9.40 . Sixty five (39.98%) were male. Table 1 summarized the study parameters stratified by CAVI level. It also contains the Pearson correlation coefficient (r) and p -value of different variables with CAVI measures. Study parameters stratified by CAVI level (using median and 75 percentile of CAVI as the cut point) can be found in Supplementary Materials (Tables 1–4).


CAVI and $CAVI_0$ is highly correlated (Pearson's $r = 0.95$ and $p < 0.001$). Further correlation analysis of the study variables with CAVI shows that CAVI is significantly correlated with age, SBP, DBP, MBP. $CAVI_0$ is also significantly correlated with age, SBP and MBP. $CAVI_0$ is not correlated with DBP (Pearson's $r = 0.05$ and $p = 0.46$). Among laboratory tests, CAVI and $CAVI_0$ are correlated with BUN ($r = 0.20$, $p = 0.009$ and $r = 0.22$, $p = 0.004$ respectively) and also with HBA1c ($r = 0.21$, $p = 0.006$ and $r = 0.25$, $p = 0.001$ respectively). CAVI and $CAVI_0$ are not significantly correlated with other study variables (Table 1).

The Pearson correlation of CAVI and $CAVI_0$ with GAD-7 overall score was -0.15 ($p = 0.041$) and -0.18 ($p = 0.019$) respectively (GAD-7 results can be seen in Table S5 of the Supplementary Material). Prediction eclipse of Pearson's correlation analysis between CAVI and $CAVI_0$ with age, MBP, DBP and SBP can be seen in Figure 1.

Table 1 | Characteristics of study participants stratified by sex

Variables	Mean ± SD	Sex		<i>p</i>	Pearson correlation with CAVI		Pearson correlation with CAVI ₀	
	Number (%)	Male	Female		<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>
CAVI	8.64 ± 1.08	8.69 ± 1.15	8.60 ± 1.04	0.614	1	–	0.95	<0.001
CAVI ₀	13.33 ± 2.68	13.37 ± 2.88	13.30 ± 2.56	0.877	0.95	–	1	–
Age (years)	63.06 ± 9.40	63.36 ± 9.77	62.86 ± 9.19	0.739	0.57	<0.001	0.61	<0.0001
BMI (kg/m ²)	24.76 ± 3.60	25.24 ± 3.29	24.44 ± 3.77	0.165	–0.06	0.41	–0.06	0.451
WC (cm)	80.53 ± 9.90	85.78 ± 8.55	77.05 ± 9.20	<0.001	–0.03	0.673	–0.03	0.684
HC (cm)	93.92 ± 6.93	94.81 ± 5.93	93.33 ± 7.49	0.183	–0.1	0.18	–0.11	0.170
ABI	1.10 ± 0.06	1.12 ± 0.06	1.10 ± 0.05	0.021	0.08	0.296	–0.01	0.897
SBP (mmHg)	130.10 ± 17.37	134.09 ± 16.57	127.45 ± 17.47	0.016	0.3	<0.001	0.33	<0.0001
DBP (mmHg)	80.74 ± 10.40	84.16 ± 10.29	78.46 ± 9.89	<0.001	0.18	0.016	0.06	0.461
HR (bpm)	69.01 ± 9.38	69 ± 9.04	69.02 ± 9.65	0.989	–0.09	0.216	–0.13	0.092
MBP (mmHg)	97.19 ± 12.04	100.81 ± 11.82	94.79 ± 11.63	0.001	0.25	0.001	0.19	0.014
T2DM	25 (15.34)	10 (15.38)	15 (15.31)	0.989	–	–	–	–
HTN	56 (34.57)	26 (40)	30 (30.93)	0.234	–	–	–	–
IHD	14 (8.59)	9 (13.85)	5 (5.10)	0.083	–	–	–	–
FH	45 (27.95)	19 (30.16)	26 (26.53)	0.719	–	–	–	–
SMK	34 (20.86)	32 (49.23)	2 (2.04)	<0.001	–	–	–	–
Pack-year	0.86 ± 4.76	2.17 ± 7.38	0	0.004	0.11	0.129	0.14	0.073
Habit Alc.	103 (63.19)	26 (40)	77 (78.57)	<0.001	–	–	–	–
Alc. vol. (cc)	51.83 ± 153.65	117.29 ± 235.74	17.30 ± 60.75	<0.001	0.02	0.726	0.05	0.595
Hab. Exe.	108 (66.26)	44 (67.69)	64 (65.31)	0.453	–	–	–	–
Exe. Freq. per week	3.28 ± 3.34	3.57 ± 3.69	3.09 ± 3.09	0.371	0.08	0.259	0.11	0.151
GAD-7	1.55 ± 2.49	1.20 ± 1.98	1.75 ± 2.76	0.135	–0.15	0.041	–0.18	0.019

BMI, body mass index; WC, waist circumference; HC, hip circumference; HR, heart rate; SMK, ever smoking cigarette; Pack-year, pack-years of smoking; Habit Alc., habit of alcohol drinking; Alc. Vol., alcohol volume consumption in a week; Hab. Exe., habit of doing exercise; Exe. Freq. per week, frequency of exercise per week; GAD-7, generalized anxiety disorder index (*p*-values of Fisher exact and ANOVA tests).

**Figure 1** | Scatter plots and 95% prediction eclipse of CAVI and CAVI₀ with systolic blood pressure, diastolic blood pressure, mean blood pressure and age.

Results of the multiple linear regression models can be found in [Table 2](#).

Median CAVI and CAVI₀ was set as the cut-off point for the logistic regression analysis. Results of these models can be found in [Table 2](#). The odds ratio and confidence interval of having a CAVI and CAVI₀ greater than the median cut-off point is shown in [Table 3](#).

4. DISCUSSION

This study performed on 163 community dwelling individuals suggesting that CAVI is significantly correlated with age, SBP, DBP, MBP and CAVI₀ is significantly correlated with age, SBP and MBP. CAVI₀ is

not correlated with DBP. CAVI and CAVI₀ are significantly correlated with BUN and also with HBA1c. None of blood pressure indices were associated with high CAVI and high CAVI₀ independent of age.

Results of this study confirms that CAVI and CAVI₀ are associated significantly and independently only to age. The fact that only age has independent effects on these indices can help to build reliable models to define individuals at the risk of atherosclerosis and CVD events, cfpWV although other studies had shown non consistent results. Bayes et al. [17] showed that CAVI is significantly different in men and women, also cardiovascular risk factors are associated with CAVI. Our results are not consistent with their study. In the study of Tabara et al. [18] the correlation of CAVI and age, SBP and BMI was 0.58, 0.17 and –0.20 respectively. These results are

consistent with our study results except our study found no significant correction between CAVI and BMI. They also found CAVI higher in males but we found this difference not significant. It might be due to the smaller sample size of our study.

Our study showed a significant correction between CAVI and CAVI₀ with BUN and HBA1c. Gomez-Marcos et al. [19] found out that CAVI is correlated positively with age, HBA1c, SBP, and DBP, and negatively correlated with waist circumference and BMI. Although they found out that there are more diabetic patients in the high CAVI group, HBA1c was not assessed in their study. However, the significant correlation between CAVI and HBA1c was also observed and confirmed by our study. About BUN serum levels, we found it significantly correlated to CAVI ($r = 0.20, p = 0.009$) and CAVI₀ ($r = 0.22, p = 0.004$). The Literature is ambiguous regarding this relationship, Park et al. [20] found no difference between BUN serum levels and different levels of CAVI, on the other hand Liu et al. [14] found them significantly correlated ($r = 0.231, p < 0.05$). The relationship also might be because of other factors in the study.

Although the effects of body weight reduction and exercise have been shown to have an important effect on reduction of arterial stiffness [21] our study just confirmed a negative nonsignificant relationship between BMI and CAVI. Beneficial effects of exercise on

Table 2 | Linear multiple regression for CAVI and CAVI₀

			Model 1	Model 2	Model 3	Model 4
CAVI	Age (years)	B	0.066	0.061	0.066	0.064
		p	<0.001	<0.001	<0.001	<0.001
	SBP (mmHg)	B	–	0.011	–	–
		p	–	0.005	–	–
	DBP (mmHg)	B	–	–	0.019	–
		p	–	–	0.003	–
MBP (mmHg)	B	–	–	–	0.017	
	p	–	–	–	0.002	
CAVI ₀	Age (years)	B	0.17	0.16	0.17	–
		p	<0.001	<0.001	<0.001	–
	SBP (mmHg)	B	–	0.031	–	–
		p	–	0.001	–	–
	MBP (mmHg)	B	–	–	0.028	–
		p	–	–	0.042	–

CAVI: Model 1 = age; Model 2 = age, SBP; Model 3 = age, DBP; Model 4 = age, MBP;
CAVI₀: Model 1 = age; Model 2 = age, SBP; Model 3 = age, MBP.

anti-oxidation state, blood glucose level, angiogenesis and endothelial functions can be a reason for the negative relationship between these measures [22].

Although anxiety has been shown to have an important effect on CVD mortality [23], its role in the arterial stiffness and atherosclerosis is controversial [19]. Our study showed the significant and negative correlation ($-0.15, p = 0.041$) between GAD and CAVI and a negative correlation ($-0.18, p = 0.019$) between GAD and CAVI₀. Yeragani et al. [24] showed that arterial stiffness is significantly higher in the patients with high anxiety. Our study results are in contrast with their findings. However, after controlling for the other risk factors in the regression models, anxiety showed no significant independent association with CAVI.

There are some differences between how CAVI₀ is calculated and the most important one is that CAVI₀ do not use SBP in its formula. It is worth noting that CAVI₀ has been doubted because it has high cross-sectional dependency on the DBP [25]. This dependency, however, can potentially be explained from the typical increase in pulse pressure that is associated with an increase in arterial stiffness [26]. This might explain the fact that CAVI₀ does not have a significant correlation with SBP in our study. In a study Shirai et al. [27] observed a negative correlation between DBP and CAVI₀ which could be explainable as the subjects of their study with stiffer arteries had normal SBP, thus should have lower DBP because of the fact that people with stiffer arteries have larger pulse pressure [26]. This cannot be applicable for our study, therefore there is no significant correlation between DBP and CAVI₀ in our study. The other fact that should be taken into account is that SBP and DBP are both important in the evaluation of the risk assessment of hypertension complications. As uncontrolled DBP is associated with a greater risk of CVD events and uncontrolled SBP is more associated with poorer prognosis [28], when evaluating the results of CAVI₀, we should consider that this index maybe not reflective of DBP in the patient.

5. CONCLUSION

Among all the factors that have some association with CAVI in community dwelling individual, only age could be served as independent factors influencing CAVI and CAVI₀. As SBP and DBP are

Table 3 | Logistic regression analysis of CAVI and CAVI₀ greater than median cut-off point

			Model 1	Model 2	Model 3	Model 4
CAVI	Age (years)	OR (CI)	1.14 (1.09 – 1.19)	1.13 (1.08 – 1.19)	1.14 (1.09 – 1.20)	1.14 (1.09 – 1.19)
		p	<0.001	<0.0001	<0.0001	<0.0001
	SBP (mmHg)	OR (CI)	–	1.01 (0.99 – 1.03)	–	–
		p	–	0.1368	–	–
	DBP (mmHg)	OR (CI)	–	–	1.09 (0.99 – 1.06)	–
		p	–	–	0.1540	–
MBP (mmHg)	OR (CI)	–	–	–	1.02 (0.99 – 1.05)	
	p	–	–	–	0.1260	
CAVI ₀	Age (years)	OR (CI)	1.17 (1.11 – 1.23)	1.16 (1.10 – 1.22)	1.16 (1.11 – 1.23)	–
		p	<0.0001	<0.0001	<0.0001	–
	SBP (mmHg)	OR (CI)	–	1.01 (0.99 – 1.03)	–	–
		p	–	0.1517	–	–
	MBP (mmHg)	OR (CI)	–	–	1.007 (0.97 – 1.03)	–
		p	–	–	0.6457	–

CAVI: Model 1 = age; Model 2 = age, SBP; Model 3 = age, DBP; Model 4 = age, MBP; CAVI₀: Model 1 = age; Model 2 = age, SBP; Model 3 = age, MBP.

not independently associated with CAVI and $CAVI_0$, these indices could be considered reliable in differentiating people with high risks of CVD, as it is not dependent on other risk factors. As $CAVI_0$ is not correlated with DBP, when evaluating the results of $CAVI_0$, we should consider that this index maybe not be reflective of DBP in the patient.

CONFLICTS OF INTEREST

The authors declare they have no conflicts of interest.

AUTHORS' CONTRIBUTION

CB and LL contributed in conceptualization. JA and AA contributed in data curation and writing the original draft. JA contributed in formal analysis. CB and TT contributed in investigation. JA and CB contributed in methodology. SVW contributed in resources and validation. JA and TT contributed in writing the review and editing. SW contributed in visualization. NH and CB contributed in funding acquisition. JA and NH contributed in software. CB contributed in supervision and project administration.

FUNDING

No financial support was provided.

SUPPLEMENTARY MATERIALS

Supplementary data related to this article can be found at <https://doi.org/10.2991/artres.k.201124.002>.

REFERENCES

- [1] Namekata T, Suzuki K, Ishizuka N, Shirai K. Establishing baseline criteria of cardio-ankle vascular index as a new indicator of arteriosclerosis: a cross-sectional study. *BMC Cardiovasc Disord* 2011;11:51.
- [2] Laurent S, Boutouyrie P, Asmar R, Gautier I, Laloux B, Guize L, et al. Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension* 2001;37:1236–41.
- [3] Mancia G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Böhm M, et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2013;31:1281–357.
- [4] Zhong Q, Hu MJ, Cui YJ, Liang L, Zhou MM, Yang YW, et al. Carotid-femoral pulse wave velocity in the prediction of cardiovascular events and mortality: an updated systematic review and meta-analysis. *Angiology* 2018;69:617–29.
- [5] Wohlfahrt P, Cifková R, Movsisyan N, Kunzová Š, Lešovský J, Homolka M, et al. Reference values of cardio-ankle vascular index in a random sample of a white population. *J Hypertens* 2017;35:2238–44.
- [6] Shirai K, Utino J, Otsuka K, Takata M. A novel blood pressure-independent arterial wall stiffness parameter; cardio-ankle vascular index (CAVI). *J Atheroscler Thromb* 2006;13:101–7.
- [7] Alizargar J, Bai CH, Hsieh NC, Wu SV, Weng SY, Wu JP. Association of kidney function tests with a cardio-ankle vascular index in community-dwelling individuals with a normal or mildly decreased estimated glomerular filtration rate. *Medicina (Kaunas)* 2019;55:657.
- [8] Shah BM, Shah S, Kandula NR, Gadgil MD, Kanaya AM. Psychosocial factors associated with subclinical atherosclerosis in South Asians: the MASALA study. *J Immigr Minor Health* 2016;18:1317–27.
- [9] da Luz PL, Coimbra SR. Wine, alcohol and atherosclerosis: clinical evidences and mechanisms. *Braz J Med Biol Res* 2004;37:1275–95.
- [10] Siasos G, Tsigkou V, Kokkou E, Oikonomou E, Vavuranakis M, Vlachopoulos C, et al. Smoking and atherosclerosis: mechanisms of disease and new therapeutic approaches. *Curr Med Chem* 2014;21:3936–48.
- [11] Al-Mamari A. Atherosclerosis and physical activity. *Oman Med J* 2009;24:173–8.
- [12] Alizargar J, Bai CH. Factors associated with carotid Intima media thickness and carotid plaque score in community-dwelling and non-diabetic individuals. *BMC Cardiovasc Disord* 2018;18:21.
- [13] Wang H, Liu J, Zhao H, Zhao X, Li L, Shi H, et al. Relationship between cardio-ankle vascular index and plasma lipids in hypertension subjects. *J Hum Hypertens* 2015;29:105–8.
- [14] Liu H, Hong Y, Wu X, Zhang X, Huang M, Wang H. Evaluation of cardio-ankle vascular index and influencing factors in natural population of she minority in China. *Intern Med Open J* 2016;6:1–10.
- [15] Spronck B, Mestanik M, Tonhajzerova I, Jurko A, Jurko T, Avolio AP, et al. Direct means of obtaining $CAVI_0$ —a corrected cardio-ankle vascular stiffness index (CAVI)—from conventional CAVI measurements or their underlying variables. *Physiol Measure* 2017;38:N128–N37.
- [16] Spronck B, Mestanik M, Tonhajzerova I, Jurko A, Tan I, Butlin M, et al. Easy conversion of cardio-ankle vascular index into $CAVI_0$: influence of scale coefficients. *J Hypertens* 2019;37:1913–14.
- [17] Elosua-Bayés M, Martí-Lluch R, García-Gil MDM, Camós L, Comas-Cufí M, Blanch J, et al. Association of classic cardiovascular risk factors and lifestyles with the cardio-ankle vascular index in a general mediterranean population. *Rev Esp Cardiol (Engl Ed)* 2018;71:458–65 [Article in En, Spanish].
- [18] Tabara Y, Setoh K, Kawaguchi T, Takahashi Y, Kosugi S, Nakayama T, et al. Factors affecting longitudinal changes in cardio-ankle vascular index in a large general population: the Nagahama study. *J Hypertens* 2018;36:1147–53.
- [19] Gómez-Marcos M, Recio-Rodríguez JI, Patino-Alonso MC, Agudo-Conde C, Gómez-Sánchez L, Gomez-Sanchez M, et al. Cardio-ankle vascular index is associated with cardiovascular target organ damage and vascular structure and function in patients with diabetes or metabolic syndrome, LOD-DIABETES study: a case series report. *Cardiovasc Diabetol* 2015;14:7.
- [20] Park SY, Chin SO, Rhee SY, Oh S, Woo JT, Kim SW, et al. Cardio-ankle vascular index as a surrogate marker of early atherosclerotic cardiovascular disease in Koreans with type 2 diabetes mellitus. *Diabetes Metab J* 2018;42:285–95.

- [21] Satoh N, Shimatsu A, Kato Y, Araki R, Koyama K, Okajima T, et al. Evaluation of the cardio-ankle vascular index, a new indicator of arterial stiffness independent of blood pressure, in obesity and metabolic syndrome. *Hypertens Res* 2008;31:1921–30.
- [22] Thitiwuthikiat P, Imerbtham T, Weeraphan O, Siri Wittayawan D. Improvement of cardio-ankle vascular index by arm-swing exercise in older adults. *J Health Sci Med Res* 2018;36:53–60.
- [23] Janszky I, Ahnve S, Lundberg I, Hemmingsson T. Early-onset depression, anxiety, and risk of subsequent coronary heart disease: 37-year follow-up of 49,321 young Swedish men. *J Am Coll Cardiol* 2010;56:31–7.
- [24] Yeragani VK, Tancer M, Seema KP, Josyulab K, Desai N. Increased pulse-wave velocity in patients with anxiety: implications for autonomic dysfunction. *J Psychosom Res* 2006; 61:25–31.
- [25] Alizargar J, Hsieh NC, Wu SFV, Weng SY. Using the cardio-ankle vascular index (CAVI) or the mathematical correction form (CAVI₀) in clinical practice. *Int J Mol Sci* 2020;21:2410.
- [26] Spronck B, Jurko A, Mestanik M, Avolio AP, Tonhajzerova I. Reply to comments: using the cardio-ankle vascular index (CAVI) or the mathematical correction form (CAVI₀) in clinical practice. *Int J Mol Sci* 2020;21:2647.
- [27] Shirai K, Suzuki K, Tsuda S, Shimizu K, Takata M, Yamamoto T, et al. Comparison of cardio-ankle vascular index (CAVI) and CAVI₀ in large healthy and hypertensive populations. *J Atheroscler Thromb* 2019;26:603–15.
- [28] Konstantinidis D, Tsioufis C, Kasiakogias A, Iliakis P, Kakouri N, Andrikou E, et al. Systolic versus diastolic blood pressure as predictors of cardiovascular events among treated hypertensive patients: a 6-year prospective study. 2019;40:ehz746.0422.