



## Conference Abstract

# P.19 Intradialytic Changes in Cerebral Blood Flow and Regional Changes in Arterial Stiffness

Mathilde Paré<sup>1,2,3,4,\*</sup>, Hasan Obeid<sup>1,2,5,6</sup>, Lawrence Labrecque<sup>3,4</sup>, Audrey Drapeau<sup>3,4</sup>, Karine Marquis<sup>1,2</sup>, Patrice Brassard<sup>3,4</sup>, Mohsen Agharazii<sup>1,2</sup>

<sup>1</sup>CHU de Québec Research Center, L'Hôtel-Dieu de Québec

<sup>2</sup>Division of Nephrology, Faculty of Medicine, Université Laval

<sup>3</sup>Research Center of the Institut Universitaire de Cardiologie et de Pneumologie de Québec

<sup>4</sup>Department of Kinesiology, Faculty of Medicine, Université Laval

<sup>5</sup>INSERM, UMR-970, Paris Cardiovascular Research Center, 75015

<sup>6</sup>AP-HP, Pharmacology Unit, Hôpital Européen Georges Pompidou, Université de Paris

### Keywords

Arterial stiffness  
flow pulsatility

### ABSTRACT

**Purpose/Background/Objective:** Cognitive decline is highly prevalent amongst end-stage renal disease (ESRD) patients and is accelerated upon initiation of hemodialysis (HD) [1]. ESRD increases aortic stiffness and blood flow pulsatility, which may damage small vessels of target organs like the brain [2]. In this pilot study, we aimed to evaluate the acute effect of HD on cerebral blood flow and its relation to arterial stiffness.

**Methods:** Before, every hour during and after HD (T0–T4), we measured cerebral flow velocity (FV) using transcranial Doppler, blood pressure (BP) via digital finger cuff (Nexfin), cardiac activity using ECG and aortic pulse wave velocity (PWV) with Mobile-O-Graph. FV pulsatility index (PI) and transit times between ECG peak and the foot of both FV and BP waveforms (cerebral dT; digital dT) were computed using in house MATLAB-based analysis. Changes during HD were evaluated with Generalized Estimating Equation models adjusting for multiple comparisons in SPSS 26.0.

**Results:** In eight participants aged  $63 \pm 17$  y. old (4 diabetics, 3 women), peak FV decreased from baseline at T1 and T2 ( $-11.2$  cm/s,  $p = 0.007$ ;  $-12.2$  cm/s,  $p < 0.001$ ), PI decreased at T1 ( $0.81$  to  $0.77$ ,  $p = 0.005$ ), whilst minimum FV, mean BP and partial pressure of CO<sub>2</sub> remained unchanged. Digital dT increased at T3 ( $0.19$  to  $0.22$ ,  $p < 0.001$ ) and cerebral dT increased throughout HD (T1–T4,  $p < 0.005$ ), whereas aortic PWV did not change.

**Conclusions:** During hemodialysis, cerebral and digital transit times increased, suggesting decreased stiffness of small peripheral vessels, without significant changes in aortic stiffness. Reduced stiffness of cerebral arteries may partially explain decreased cerebral flow pulsatility.

### REFERENCES

- [1] Zammit AR, Katz MJ, Bitzer M, Lipton RB. Cognitive impairment and dementia in older adults with chronic kidney disease: a review. *Alzheimer Dis Assoc Disord* 2016;30:357–66.
- [2] Toth P, Tarantini S, Csiszar A, Ungvari Z. Functional vascular contributions to cognitive impairment and dementia: mechanisms and consequences of cerebral autoregulatory dysfunction, endothelial impairment, and neurovascular uncoupling in aging. *Am J Physiol Heart Circ Physiol* 2017;312:H1–H20.

© 2020 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).