

Artery Research Vol. **26(S1)**; 2020, *p*. S50 DOI: https://doi.org/10.2991/artres.k.201209.040; ISSN 1872-9312; eISSN 1876-4401 https://www.atlantis-press.com/journals/artres



Conference Abstract P.27 Mechanisms of NADPH Oxidase Participation in the Regulation of Diaphragm Artery Contractile Responses

Anna Borzykh^{1,*}, Ilya Kuzmin², Olga Vinogradova^{1,2}, Olga Tarasova^{1,2}

¹SRC RF – Institute for Biomedical Problems RAS ²M.V. Lomonosov Moscow State University

Keywords

Artery diaphragm NADPH oxidase

ABSTRACT

Reactive oxygen species (ROS) produced by NADPH-oxidase (NOX) participate in vascular tone control, but their effects in the arteries of respiratory muscles is poorly understood. Possible targets of vasoregulatory ROS influence are NO-pathway in the endothelium and Rho-kinase pathway in smooth muscle cells. Therefore, the aim of this study was to evaluate the interaction of NOX-dependent control with NO- and Rho-kinase signaling pathways in rat diaphragm arteries (DA).

Methods: The segments of DA were isolated from male Wistar rats and mounted in wire myograph (DMT A/S). We studied the effects of NOX inhibitor VAS2870 (1 μ M) on contractile responses to α_1 -adrenergic agonist methoxamine in the absence and in the presence of NO synthase (L-NNA 100 μ M) or Rho-kinase (Y27632, 3 μ M) inhibitors as well as in the presence of NO donor DEA/NO.

Results: VAS2879 prominently attenuated the contractile responses of DA to methoxamine (30% decrease of the area under the concentration-response curve). L-NNA and Y27632 increased and decreased methoxamine-induced contraction of DA, respectively. L-NNA did not change the effects of VAS2870 and the sensitivity to DEA/NO did not differ in arteries with active and inhibited NOX. Along with that Y27632 eliminated the effects of VAS2879 on DA contractile responses to methoxamine.

Conclusions: We showed that NOX-produced ROS potentiate contractile responses of DA. ROS did not affect the activity of NO-pathway in either endothelial or smooth muscle cells of DA. However, ROS modulate the activity of the Rho-kinase pathway in DA smooth muscle cells. Supported by RSF (project No 19-75-00060).

© 2020 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (http://creativecommons.org/licenses/by-nc/4.0/).

^{*}Corresponding author. Email: borzykh.anna@gmail.com