



Journal of Nonlinear Mathematical Physics

ISSN (Online): 1776-0852

ISSN (Print): 1402-9251

Journal Home Page: <https://www.atlantis-press.com/journals/jnmp>

No periodic orbits for the type A Bianchi's systems

Claudio A. Buzzi, Jaume Llibre

To cite this article: Claudio A. Buzzi, Jaume Llibre (2015) No periodic orbits for the type A Bianchi's systems, Journal of Nonlinear Mathematical Physics 22:2, 170–179, DOI: <https://doi.org/10.1080/14029251.2015.1023561>

To link to this article: <https://doi.org/10.1080/14029251.2015.1023561>

Published online: 04 January 2021

No periodic orbits for the type A Bianchi's systems

Claudio A. Buzzi

*Departamento de Matemática – IBILCE–UNESP,
Rua C. Colombo, 2265, CEP 15054–000 S. J. Rio Preto, São Paulo, Brazil
buzzi@ibilce.unesp.br*

Jaume Llibre

*Departament de Matemàtiques – Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain
jllibre@mat.uab.cat*

Received 6 November 2014

Accepted 7 January 2015

It is known that the 6 models of Bianchi class A have no periodic solutions. In this article we provide a new, direct, unified and easier proof of this result.

Keywords: Bianchi cosmological models; periodic orbits.

2000 Mathematics Subject Classification: 34C05, 34C25, 34C60

1. Introduction

This paper deals with the Bianchi's cosmological models. These models require a three dimensional Lie algebra and Bianchi [1,2] was the first to solve the problem of classifying three dimensional Lie algebras. There are nine types of models according with the dimension n of the algebra.

- a) $n = 0$ Type *I*;
- b) $n = 1$ Type *II*, *III*;
- c) $n = 2$ Type *IV*, *V*, *VI*, *VII*;
- d) $n = 3$ Type *VIII*, *IX*.

Let $\{X_1, X_2, X_3\}$ be an appropriate basis of the three dimensional Lie Algebra. The classification depends on a scalar $a \in \mathbb{R}$ and a vector (n_1, n_2, n_3) with $n_i \in \{+1, -1, 0\}$ such that

$$[X_1, X_2] = n_3 X_3, \quad [X_2, X_3] = n_1 X_1 - a X_2, \quad [X_3, X_1] = n_2 X_2 + a X_1,$$

where $[\cdot, \cdot]$ is the Lie bracket. In particular, for $a = 0$ we obtain models of type *A* and for $a \neq 0$ we obtain models of type *B*. For more details see Bogoyavlensky [3].

According with [3] all cases of type *A* are Hamiltonian systems in the phase space p_i, q_i for $i = 1, 2, 3$ with the Hamiltonian function

$$H = \frac{1}{(q_1 q_2 q_3)^{\frac{1-k}{2}}} \left(T + \frac{1}{4} V_G \right),$$

Type	I	II	VI ₀	VII ₀	VIII	IX
n_1	0	1	1	1	1	1
n_2	0	0	-1	1	1	1
n_3	0	0	0	0	-1	1

Table 1. Cosmologies of types A

with $0 \leq k \leq 1$ and where

$$T = 2 \sum_{i < j}^3 p_i p_j q_i q_j - \sum_{i=1}^3 p_i^2 q_i^2, \text{ and } V_G = 2 \sum_{i < j}^3 n_i n_j q_i q_j - \sum_{i=1}^3 n_i^2 q_i^2.$$

If we rescale the time by τ defined by $d\tau = (q_1 q_2 q_3)^{-k/2} dt$ then the system becomes

$$\begin{aligned} \dot{q}_1 &= \frac{2q_1}{(q_1 q_2 q_3)^{(1-k)/2}} (p_2 q_2 + p_3 q_3 - p_1 q_1), \\ \dot{q}_2 &= \frac{2q_2}{(q_1 q_2 q_3)^{(1-k)/2}} (p_3 q_3 + p_1 q_1 - p_2 q_2), \\ \dot{q}_3 &= \frac{2q_3}{(q_1 q_2 q_3)^{(1-k)/2}} (p_1 q_1 + p_2 q_2 - p_3 q_3), \\ \dot{p}_1 &= -\frac{1}{(q_1 q_2 q_3)^{(1-k)/2}} (2p_1(p_2 q_2 + p_3 q_3 - p_1 q_1) + \\ &\quad \frac{1}{2} n_1(n_2 q_2 + n_3 q_3 - n_1 q_1)) + \frac{1-k}{2} \frac{\bar{H}}{q_1}, \\ \dot{p}_2 &= -\frac{1}{(q_1 q_2 q_3)^{(1-k)/2}} (2p_2(p_3 q_3 + p_1 q_1 - p_2 q_2) + \\ &\quad \frac{1}{2} n_2(n_3 q_3 + n_1 q_1 - n_2 q_2)) + \frac{1-k}{2} \frac{\bar{H}}{q_2}, \\ \dot{p}_3 &= -\frac{1}{(q_1 q_2 q_3)^{(1-k)/2}} (2p_3(p_1 q_1 + p_2 q_2 - p_3 q_3) + \\ &\quad \frac{1}{2} n_3(n_1 q_1 + n_2 q_2 - n_3 q_3)) + \frac{1-k}{2} \frac{\bar{H}}{q_3}, \end{aligned}$$

with $\bar{H} = T + V_G/4$. The constants n_1, n_2, n_3 determine the type of the model according with Table 1. Performing the change of coordinates $ds = (q_1 q_2 q_3)^{\frac{1-k}{2}} d\tau$ and $q_i = x_i$ and $p_i = x_{i+3}/x_i$, $i = 1, 2, 3$,

we obtain the system

$$\begin{aligned}\dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= n_1x_1(n_1x_1 - n_2x_2 - n_3x_3) + \frac{1}{4}(k-1)F_2, \\ \dot{x}_5 &= n_2x_1(-n_1x_1 + n_2x_2 - n_3x_3) + \frac{1}{4}(k-1)F_2, \\ \dot{x}_6 &= n_3x_3(-n_1x_1 - n_2x_2 + n_3x_3) + \frac{1}{4}(k-1)F_2,\end{aligned}\tag{1.1}$$

with

$$\begin{aligned}F_2 &= n_1^2x_1^2 + n_2^2x_2^2 + n_3^2x_3^2 - 2n_1n_2x_1x_2 - 2n_1n_3x_1x_3 - 2n_2n_3x_2x_3 \\ &\quad + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6.\end{aligned}$$

Note that system (1.1) is a homogeneous system of degree 2, and the first integral given by the Hamiltonian H becomes

$$\begin{aligned}\mathcal{H} &= (x_1x_2x_3)^{\frac{k-1}{2}} (n_1^2x_1^2 + n_2^2x_2^2 + n_3^2x_3^2 - 2n_1n_2x_1x_2 - 2n_1n_3x_1x_3 \\ &\quad - 2n_2n_3x_2x_3 + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6).\end{aligned}$$

We note that the class of systems (1.1), being a homogeneous system of degree 2, can be modified so that it becomes isochronous by adding in the right-hand side of each of its six equations of motion the term $i\omega x_n$ (where i is the imaginary unit, ω an arbitrary real constant, and of course n going from 1 to 6 numbers the six equations of motions). For a proof, see for instance Calogero [5]. Of course such a modification entails a doubling of the number of real dependent variables (from 6 to 12), and the physical interpretation of the resulting system is moot.

It is known that all the Bianchi class A models do not have periodic orbits. We note that in this paper always that we talk about a periodic orbit or solution we are talking on a bounded periodic orbit or solution. This has been proved using evolutions equations associated to these models, and showing that such equations always have some monotone function evaluated on the orbits. Consequently these models cannot exhibit periodic motion. For more details, see chapter 6 of the book by Wainwright and Ellis [17]. Additionally Starkov in [16] provided another proof that Bianchi VIII and IX systems has no periodic solutions. The Bianchi systems continue to be very interesting for the research, see for instance [13, 14] and the references quoted there.

In this article we provide a new, direct and easier proof on the non-existence of periodic orbits for the 6 models of Bianchi class A. For some additional dynamical properties of the Bianchi models see [4, 6–8] and the references quoted there.

2. The Bianchi I system

In this section we consider the Bianchi I system. According with Table 1 we have $n_1 = n_2 = n_3 = 0$. So system (1.1) becomes

$$\begin{aligned}\dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= \frac{k-1}{4}(x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6), \\ \dot{x}_5 &= \frac{k-1}{4}(x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6), \\ \dot{x}_6 &= \frac{k-1}{4}(x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6).\end{aligned}\tag{2.1}$$

Proposition 2.1. *The Bianchi I system, given by (2.1), does not have periodic solutions.*

Proof. System (2.1) has the first integrals $F_1 = x_4 - x_5$ and $F_2 = x_4 - x_6$. Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic solution of (2.1). So, there exist two constants a and b such that $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ for all t . We have that $x_4(t)$ is a periodic solution of the equation

$$\dot{x}_4 = \frac{k-1}{4}(-3x_4^2 - 2(a+b)x_4 + (a-b)^2). \quad (2.2)$$

Observe that the discriminant of the equation $-3x_4^2 - 2(a+b)x_4 + (a-b)^2 = 0$ is $\Delta = 4(a+b)^2 + 12(a-b)^2$. So $\Delta \geq 0$, if $\Delta > 0$ then equation (2.2) has two equilibrium points. One of them is an attractor and the other one is a repeller (see Figure 1(a)). If $\Delta = 0$ then we have just one equilibrium point which is semi-stable (see Figure 1(b)). In both cases the unique possibility in order to $x_4(t)$ be periodic is that $x_4(t) = c$ constant for all t being c an equilibrium of (2.2). Substituting $x_4(t) = c$, $x_5(t) = a + c$ and $x_6(t) = b + c$ in the first three equations of (2.1), and using that $x_1(t)$, $x_2(t)$ and $x_3(t)$ are periodic, we get that $x_1(t)$, $x_2(t)$ and $x_3(t)$ must be constant for all t . So $\Gamma(t)$ is constant, i.e. $\Gamma(t)$ is an equilibrium point of (2.1), and (2.1) do not have periodic solutions. \square



Fig. 1. Phase portrait of the differential equation (2.2).

3. The Bianchi II system

In this section we consider the Bianchi II system. According with Table 1 we have $n_1 = 1$ and $n_2 = n_3 = 0$. So system (1.1) becomes

$$\begin{aligned} \dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= x_1^2 + \frac{k-1}{4}F, \\ \dot{x}_5 &= \frac{k-1}{4}F, \\ \dot{x}_6 &= \frac{k-1}{4}F, \end{aligned} \quad (3.1)$$

where $F = x_1^2 + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6$.

Proposition 3.1. *The Bianchi II system, given by (3.1), does not have periodic solutions.*

Proof. Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic solution of (3.1). The real function $x_4(t) - x_5(t)$ is periodic. So, it is bounded. If it is not constant then there exists t_0 such that $\dot{x}_4(t_0) - \dot{x}_5(t_0) < 0$, but from (3.1) we have that $\dot{x}_4(t_0) - \dot{x}_5(t_0) = (x_1(t_0))^2$. It implies that there exists a constant $a \in \mathbb{R}$ such that $x_5(t) = x_4(t) + a$ and $x_1(t) = 0$ for all t . By using the same argument we have that there exists another constant $b \in \mathbb{R}$ such that $x_6(t) = x_4(t) + b$ for all t . Next step is substitute $x_1(t) = 0$, $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ in the equation $\dot{x}_4 = x_1^2 + \frac{k-1}{4}F$, we get $\dot{x}_4 = -3x_4^2 - (a+b)x_4 + (a-b)^2$. By using the same argument of the proof of Proposition 2.1 we conclude the proof of this proposition. \square

4. The Bianchi VI_0 system

In this section we consider the Bianchi VI_0 system. According with Table 1 we have $n_1 = 1$, $n_2 = -1$ and $n_3 = 0$. So system (1.1) becomes

$$\begin{aligned}\dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= x_1(x_1 + x_2) + \frac{k-1}{4}F, \\ \dot{x}_5 &= x_2(x_1 + x_2) + \frac{k-1}{4}F, \\ \dot{x}_6 &= \frac{k-1}{4}F,\end{aligned}\tag{4.1}$$

where $F = x_1^2 + x_2^2 + 2x_1x_2 + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6$.

Proposition 4.1. *The Bianchi VI_0 system, given by (4.1), does not have periodic solutions.*

Proof. Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic solution of (4.1). The real function $x_4(t) + x_5(t) - 2x_6(t)$ is periodic. So, it is bounded. If it is not constant then there exists t_0 such that $\dot{x}_4(t_0) + \dot{x}_5(t_0) - 2\dot{x}_6(t_0) < 0$, but from (4.1) we have that $\dot{x}_4(t_0) + \dot{x}_5(t_0) - 2\dot{x}_6(t_0) = (x_1(t_0) + x_2(t_0))^2$. It implies that $\dot{x}_4(t) + \dot{x}_5(t) - 2\dot{x}_6(t) = 0 = (x_1(t) + x_2(t))^2$ for all t . Substituting $x_1(t) = -x_2(t)$ in (4.1) we have that $\dot{x}_4(t) - \dot{x}_5(t) = 0$ and $\dot{x}_4(t) - \dot{x}_6(t) = 0$ for all t . There exist constants $a, b \in \mathbb{R}$ such that $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ for all t . Next step is substitute $x_1(t) = -x_2(t)$, $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ in the equation $\dot{x}_4 = x_1(x_1 + x_2) + \frac{k-1}{4}F$. By using the same argument of the proof of Proposition 2.1 we conclude the proof of this proposition. \square

5. The Bianchi VII_0 system

In this section we consider the Bianchi VII_0 system. According with Table 1 we have $n_1 = 1$ and $n_2 = 1$ and $n_3 = 0$. So system (1.1) becomes

$$\begin{aligned}\dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= x_1(x_1 - x_2) + \frac{k-1}{4}F, \\ \dot{x}_5 &= x_2(-x_1 + x_2) + \frac{k-1}{4}F, \\ \dot{x}_6 &= \frac{k-1}{4}F,\end{aligned}\tag{5.1}$$

where $F = x_1^2 + x_2^2 - 2x_1x_2 + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6$.

Proposition 5.1. *The Bianchi VII_0 system, given by (5.1), does not have periodic solutions.*

Proof. Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic solution of (5.1). The real function $x_4(t) + x_5(t) - 2x_6(t)$ is periodic. So, it is bounded. If it is not constant then there exists t_0 such that $\dot{x}_4(t_0) + \dot{x}_5(t_0) - 2\dot{x}_6(t_0) < 0$, but from (5.1) we have that $\dot{x}_4(t_0) + \dot{x}_5(t_0) - 2\dot{x}_6(t_0) = (x_1(t_0) - x_2(t_0))^2$. It implies that $\dot{x}_4(t) + \dot{x}_5(t) - 2\dot{x}_6(t) = 0 = (x_1(t) - x_2(t))^2$ for all t . Substituting $x_1(t) = x_2(t)$ in (5.1) we have that $\dot{x}_4(t) - \dot{x}_5(t) = 0$ and $\dot{x}_4(t) - \dot{x}_6(t) = 0$ for all t . There exist constants $a, b \in \mathbb{R}$ such that $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ for all t . Next step is substitute $x_1(t) = x_2(t)$, $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ in the equation $\dot{x}_4 = x_1(x_1 - x_2) + \frac{k-1}{4}F$. By using the same argument of the proof of Proposition 2.1 we conclude the proof of this proposition. \square

6. The Bianchi VIII system

In this section we consider the Bianchi VIII system. According with Table 1 we have $n_1 = 1$ and $n_2 = 1$ and $n_3 = -1$. So system (1.1) becomes

$$\begin{aligned}\dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= x_1(x_1 - x_2 + x_3) + \frac{k-1}{4}F, \\ \dot{x}_5 &= x_2(-x_1 + x_2 + x_3) + \frac{k-1}{4}F, \\ \dot{x}_6 &= x_3(x_1 + x_2 + x_3) + \frac{k-1}{4}F,\end{aligned}\tag{6.1}$$

where

$$F = x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 + 2x_1x_3 + 2x_2x_3 + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6.\tag{6.2}$$

Lemma 6.1. *The hyperplanes $\{x_1 = 0\}$, $\{x_2 = 0\}$, and $\{x_3 = 0\}$ are invariant manifolds for system (6.1) and there is no periodic orbits in these hyperplanes.*

Proof. Clearly the hyperplanes $\{x_1 = 0\}$, $\{x_2 = 0\}$, and $\{x_3 = 0\}$ are invariant manifolds for system (6.1), i.e. if a solution of (6.1) has a point in $\{x_i = 0\}$ then the whole solution is contained in $\{x_i = 0\}$. Now we prove that in the hyperplanes $\{x_1 = 0\}$, $\{x_2 = 0\}$, and $\{x_3 = 0\}$ there are no periodic orbits. Let $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ be a periodic solution of (6.1). Suppose that $\Gamma(t)$ is in $\{x_1 = 0\}$. From (6.1) we have $\dot{x}_5 + \dot{x}_6 - 2\dot{x}_4 = (x_2 + x_3)^2$. We get that $x_1(t) = 0$ and $x_2(t) = -x_3(t)$ for all t . Substituting these conditions in the equations of (6.1) we have that $\dot{x}_4 = \dot{x}_5 = \dot{x}_6$, and so there exist constants a and b such that $x_5(t) = x_4(t) + a$ and $x_6(t) = x_4(t) + b$ for all t . Substituting all these conditions in the fourth equation of (6.1) we obtain again the equation (2.2). So, in order that Γ be periodic, $x_4(t)$ is constant. Now from the second and third equations of (6.1) we have that $x_2(t)$ and $x_3(t)$ are constants, and Γ is an equilibrium point instead of a periodic orbit. In the same way we can prove that there are no periodic orbits in $\{x_2 = 0\}$ and in $\{x_3 = 0\}$. \square

Consider the three sets

$$\begin{aligned}F^+ &= \{x \in \mathbb{R}^6 : F(x) > 0\}, \\ F^0 &= \{x \in \mathbb{R}^6 : F(x) = 0\} \text{ and} \\ F^- &= \{x \in \mathbb{R}^6 : F(x) < 0\},\end{aligned}$$

where F is given in (6.2).

Lemma 6.2. *The sets F^+ , F^0 and F^- are invariant by system (6.1) and there are no periodic orbits in the set F^- .*

Proof. First of all observe that if we call X the vector field associated to the system (6.1) then we have that

$$XF = \langle X, \nabla F \rangle = -\frac{1}{2}(k-1)(x_4 + x_5 + x_6)F,\tag{6.3}$$

where $\langle \cdot, \cdot \rangle$ denotes the standard inner product in \mathbb{R}^6 and ∇F is the gradient of F . From (6.3) we get that F^0 is an invariant set to (6.1), and consequently F^+ and F^- also are invariant.

Now we prove that there are not periodic orbit in F^- . Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic orbit of (6.1) and that it is in the set F^- . By Lemma 6.1 we have that $x_1(t) \neq 0$, $x_2(t) \neq 0$ and $x_3(t) \neq 0$ for all t . Consider the function

$$h(x) = \frac{x_4 + x_5 + x_6}{x_1 x_2 x_3}.$$

Observe that $h \circ \Gamma$ is defined for all t and it is a periodic function. So there exists at least a point $t = t_0$ such that $(h \circ \Gamma)(t_0) = 0$. We have

$$(h \circ \Gamma)(t) = \langle \nabla h, \dot{\Gamma}(t) \rangle = Xh(\Gamma(t)).$$

But

$$Xh(x) = (1 + 3k)F(x) - 8(x_4^2 + x_5^2 + x_6^2), \quad (6.4)$$

which is always negative in the set F^- . So the periodic orbit $\Gamma(t)$ cannot be contained in F^- . \square

Let U be a subset of \mathbb{R}^6 . Let $h : U \rightarrow \mathbb{R}$ be a C^1 function. By $S(h)$ we denote the set $\{x \in \mathbb{R}^6 : Xh(x) = 0\}$. Suppose that we are interested in the localization of the periodic orbits of system $\dot{x} = X(x)$ located in the set U . We define $h_{inf} = \inf\{h(x) : x \in U \cap S(h)\}$, $h_{sup} = \sup\{h(x) : x \in U \cap S(h)\}$.

The following two propositions are inspired by the formulation and proofs of localization theorems which were proposed and developed by Krishchenko and Starkov, firstly, for periodic orbits in [9, 10], and later for compact invariant sets in [11], see also [12]. Particular cases of these results can be found in [15].

Proposition 6.1. *All the periodic orbits of system $\dot{x} = X(x)$ located in U are contained in the set $\{x \in U : h_{inf} \leq h(x) \leq h_{sup}\}$.*

Proof. Let $\Gamma(t)$ be a periodic orbit of system $\dot{x} = X(x)$ contained in the set U . Denote by $\gamma = \{\Gamma(t) : t \in \mathbb{R}\}$. The set γ is compact and so the C^1 function h , restricted to the set γ , has a maximum M and a minimum m . In particular $\gamma \subset \{x \in U : m \leq h(x) \leq M\}$. For all points $t = t_1$ such that $h(\Gamma(t_1)) = m$ we have that $\Gamma(t_1) \in S(h)$. It implies that $m = \inf\{h(x) : x \in \gamma \cap S(h)\}$. On the other hand we have that $\gamma \cap S(h) \subset U \cap S(h)$ implies $m \geq h_{inf}$. Analogously we have $h_{sup} \leq M$. So we have

$$\gamma \subset \{x \in U : m \leq h(x) \leq M\} \subset \{x \in U : h_{inf} \leq h(x) \leq h_{sup}\}.$$

\square

Proposition 6.2. *Let U be a set in \mathbb{R}^6 . If $S(h) \cap U = \emptyset$ then system $\dot{x} = X(x)$ has no periodic orbits contained in U .*

Proof. Suppose that $\Gamma(t)$ is a periodic orbit of system $\dot{x} = X(x)$ contained in the set U . As we saw in the proof of Proposition 6.1, for all points $t = t_1$ such that $h(\Gamma(t_1)) = m$ we have that $\Gamma(t_1) \in S(h) \cap U$. And so $S(h) \cap U \neq \emptyset$, which is a contradiction. \square

Lemma 6.3. *There are no periodic orbits of system (6.1) located in F^0 .*

Proof. Consider the set $U = F^0 \cap \{x_1 \neq 0\} \cap \{x_2 \neq 0\} \cap \{x_3 \neq 0\}$ and the function

$$h(x) = \frac{x_4 + x_5 + x_6}{x_1 x_2 x_3}.$$

Accordingly to (6.4) we have that $S(h) = \{x \in \mathbb{R}^6 : x_4 = x_5 = x_6 = 0\}$. So $h_{inf} = h_{sup} = 0$. From Proposition 6.1 all compact invariant sets of (6.1) located on U are contained in $B = \{x_4 + x_5 + x_6 = 0\}$.

Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic orbit of (6.1) and it is in the set F^0 . By using the fact that this orbit is contained in B we have that $\dot{x}_4 + \dot{x}_5 + \dot{x}_6 = 0$. From system (6.1) and the fact that Γ are in F^0 we have that $x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 + 2x_1x_3 + 2x_2x_3 = 0$, and consequently $x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_4x_6 - 2x_5x_6 = 0$. Substituting $x_6 = -x_4 - x_5$ in the last equation we get

$$2x_4^2 + 2x_5^2 + 2(x_4 + x_5)^2 = 0,$$

and so $x_4(t) = x_5(t) = x_6(t) = 0$ for all t . Now substituting these values in system (6.1) we get that $x_1(t)$, $x_2(t)$ and $x_3(t)$ are constant functions. So Γ is not a periodic orbit. \square

Lemma 6.4. *If there exists a periodic orbit for system (6.1), then it intersects the set $\{x \in \mathbb{R}^6 : x_4 + x_5 + x_6 = 0\}$.*

Proof. Consider the set $B = \{x \in \mathbb{R}^6 : x_1 \neq 0, x_2 \neq 0, x_3 \neq 0 \text{ and } x_4 + x_5 + x_6 \neq 0\}$ and the function $h(x) = x_1 x_2 x_3$. We have that

$$Xh(x) = x_1 x_2 x_3 (x_4 + x_5 + x_6),$$

and so $S(h) \cap B = \emptyset$. According with Proposition 6.2 system (6.1) has no periodic orbits in B . If system (6.1) has a periodic orbit then it intersects $\{x \in \mathbb{R}^6 : x_4 + x_5 + x_6 = 0\}$. \square

Lemma 6.5. *There are no periodic orbits for system (6.1) located in F^+ .*

Proof. Suppose that $\Gamma(t) = (x_1(t), x_2(t), x_3(t), x_4(t), x_5(t), x_6(t))$ is a periodic orbit of system (6.1) and that it is in the set F^+ . By Lemma 6.4 we have that Γ intersects the set $\{x_4 + x_5 + x_6 = 0\}$. Consider the function $h(x) = x_1 x_2 x_3 (x_4 + x_5 + x_6)$. By Lemma 6.1 we have that $x_1(t) \neq 0$, $x_2(t) \neq 0$, and $x_3(t) \neq 0$ for all t . Observe that the zeroes of the function $h \circ \Gamma$ occur for the values t such that the orbit Γ intersects the set $\{x_4 + x_5 + x_6 = 0\}$. Computing the derivative of $h \circ \Gamma$ we have $(h \circ \Gamma)(t) = \langle \nabla h, \dot{\Gamma}(t) \rangle = Xh(\Gamma(t))$ where

$$Xh(x) = (1 + 3k)F(x) + 16(x_4x_5 + x_4x_6 + x_5x_6).$$

We observe that in the set $\{x_4 + x_5 + x_6 = 0\}$ we have $x_4x_5 + x_4x_6 + x_5x_6 = 0$. We get that in all zeroes of the real periodic function $h \circ \Gamma$ its derivative is positive. This is a contradiction, because we cannot have a periodic real function with positive derivative in all of its zeroes. \square

Proposition 6.3. *The Bianchi VIII system, given by (6.1), does not have periodic solutions.*

Proof. It follows from lemmas 6.2, 6.3, and 6.5. \square

7. The Bianchi IX system

In this section we consider the Bianchi IX system. According with Table 1 we have $n_1 = 1$ and $n_2 = 1$ and $n_3 = 1$. So system (1.1) becomes

$$\begin{aligned}\dot{x}_1 &= x_1(-x_4 + x_5 + x_6), \\ \dot{x}_2 &= x_2(x_4 - x_5 + x_6), \\ \dot{x}_3 &= x_3(x_4 + x_5 - x_6), \\ \dot{x}_4 &= x_1(x_1 - x_2 - x_3) + \frac{k-1}{4}F, \\ \dot{x}_5 &= x_2(-x_1 + x_2 - x_3) + \frac{k-1}{4}F, \\ \dot{x}_6 &= x_3(x_1 + x_2 - x_3) + \frac{k-1}{4}F,\end{aligned}\tag{7.1}$$

where

$$F = x_1^2 + x_2^2 + x_3^2 - 2x_1x_2 - 2x_1x_3 - 2x_2x_3 + x_4^2 + x_5^2 + x_6^2 - 2x_4x_5 - 2x_5x_6 - 2x_4x_6.\tag{7.2}$$

Lemma 7.1. *The hyperplanes $\{x_1 = 0\}$, $\{x_2 = 0\}$, and $\{x_3 = 0\}$ are invariant manifolds for system (7.1) and there are no periodic orbits in these hyperplanes.*

Proof. The proof is very similar to the proof of Lemma 6.1. □

Consider the three sets

$$\begin{aligned}F^+ &= \{x \in \mathbb{R}^6 : F(x) > 0\}, \\ F^0 &= \{x \in \mathbb{R}^6 : F(x) = 0\} \text{ and} \\ F^- &= \{x \in \mathbb{R}^6 : F(x) < 0\},\end{aligned}$$

where F is given in (7.2).

Lemma 7.2. *The sets F^+ , F^0 and F^- are invariant by system (7.1) and there are no periodic orbit in the set F^- .*

Proof. The proof is very similar to the proof of Lemma 6.2. □

Lemma 7.3. *There are no periodic orbits for system (7.1) located in F^0 .*

Proof. The proof is very similar to the proof of Lemma 6.3. □

Lemma 7.4. *If there exists a periodic orbit for system (7.1) then it intersects the set $\{x \in \mathbb{R}^6 : x_4 + x_5 + x_6 = 0\}$.*

Proof. It is the same proof of Lemma 6.4 □

Lemma 7.5. *There are no periodic orbits for system (7.1) located in F^+ .*

Proof. The same construction in the proof of Lemma 6.5 works in this case. □

Proposition 7.1. *The Bianchi IX system, given by (7.1), does not have periodic solutions.*

Proof. It follows from lemmas 7.2, 7.3, and 7.5. □

Acknowledgments

The first author is supported by FAPESP-BRAZIL grant 2013/24541-0. The second author is partially supported by a MINECO/FEDER grant MTM2008-03437 and MTM2013-40998-P, an AGAUR grant number 2014SGR568, an ICREA Academia, the grants FP7-PEOPLE-2012-IRSES 318999 and 316338, FEDER-UNAB-10-4E-378, and a CAPES grant number 88881.030454/2013-01 from the program CSF-PVE. The first author would like to thanks for the hospitality at the Departament de Matemàtiques of Universitat Autònoma de Barcelona where this work was done.

References

- [1] L. Bianchi, Sugli spazi a tre dimensioni che ammettono un gruppo continuo di movimenti (On the spaces of three dimensions that admit a continuous group of movements), *Soc. Ital. Sci. Mem. di Mat.* **11** (1898), 267–352.
- [2] L. Bianchi, *Lezioni sulla teoria dei gruppi continui finiti di trasformazioni (Lectures on the theory of finite continuous transformation groups)*, Spoerri, Pisa (1918), 550–557.
- [3] O.I. Bogoyavlensky, *Methods in the qualitative theory of dynamical systems in astrophysics and gas dynamics*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1985.
- [4] C.A. Buzzi, J. Llibre and P.R. da Silva, *On the dynamics of the Bianchi IX System*, *J. Phys. A: Math. Gen.* **40** (2007), 7187–7192.
- [5] F. CALOGERO, *Isochronous systems*, Corrected paperback edition of the 2008 original. Oxford University Press, Oxford, 2012.
- [6] J. CHAVARRIGA AND I.A. GARCÍA, *Integrability and explicit solutions in some Bianchi cosmological dynamical systems*, *J. Nonlinear Math. Phys.* **8** (2001), 96–105.
- [7] A. Ferragut, J. Llibre and C. Pantazi, *Analytic Integrability of Bianchi A cosmological models*, *J. of Geometry and Physics* **62** (2012), 381–386.
- [8] R.M. Gad and A.S. Alofi, Homothetic vector field in plane symmetric Bianchi type-I cosmological model in Lyra geometry, *Modern Phys. Lett. A* **29** (2014), no. 22, 1450116, 12 pp.
- [9] A.P. Krishchenko, Localization of limit cycles, *Differential Equations* **31** (1995), 1826–1833.
- [10] A.P. Krishchenko, Estimations of domains with cycles, *Computers and Mathematics with Applications* **34** (1997), 325–332.
- [11] A.P. Krishchenko, Localization of invariant compact sets of dynamical systems, *Differential Equations* **41** (2005), 1669–1676.
- [12] A.P. Krishchenko and K.E. Starkov, Localization of compact invariant sets of the Lorenz system, *Phys. Lett. A* **353** (2006), 383–388.
- [13] D. Müller and J.A. de Deus, Bianchi I solutions of effective quadratic gravity, *Internat. J. Modern Phys. D* **21** (2012), 1250037, 17 pp.
- [14] M.E. Rodrigues, I.G. Salako, M.J.S. Houndjo and J. Tossa, Locally rotationally symmetric Bianchi type-I cosmological model in $f(T)$ gravity: from early to dark energy dominated universe, *Internat. J. Modern Phys. D* **23** (2014), 1450004, 21 pp.
- [15] K.E. Starkov, Compact invariant sets of the static spherically symmetric Einstein-Yang-Mills equations, *Phys. Lett. A* **374** (2010), no. 15–16, 1728–1731.
- [16] K.E. Starkov, Compact invariant sets of the Bianchi VIII and Bianchi IX Hamiltonian systems, *Phys. Lett. A* **375** (2011), 3184–3187.
- [17] J. Wainwright and G.F.R. Ellis, *Dynamical systems in cosmology*, Cambridge University Press, Cambridge, 1997.