



Journal of Nonlinear Mathematical Physics

ISSN (Online): 1776-0852

ISSN (Print): 1402-9251

Journal Home Page: <https://www.atlantis-press.com/journals/jnmp>

Darboux Integrability of a Simplified Friedman–Robertson–Walker Hamiltonian System

Jaume Llibre, Clàudia Valls

To cite this article: Jaume Llibre, Clàudia Valls (2012) Darboux Integrability of a Simplified Friedman–Robertson–Walker Hamiltonian System, Journal of Nonlinear Mathematical Physics 19:4, 607–617, DOI:

<https://doi.org/10.1142/S1402925112500362>

To link to this article: <https://doi.org/10.1142/S1402925112500362>

Published online: 04 January 2021

Journal of Nonlinear Mathematical Physics, Vol. 19, No. 4 (2012) 1250036 (11 pages)

© J. Llibre and C. Valls

DOI: 10.1142/S1402925112500362

DARBOUX INTEGRABILITY OF A SIMPLIFIED FRIEDMAN–ROBERTSON–WALKER HAMILTONIAN SYSTEM

JAUME LLIBRE

*Departament de Matemàtiques
Universitat Autònoma de Barcelona
08193 Bellaterra, Barcelona, Catalonia, Spain
jllibre@mat.uab.cat*

CLÀUDIA VALLS

*Departamento de Matemática
Instituto Superior Técnico
Universidade Técnica de Lisboa
Av. Rovisco Pais 1049-001, Lisboa, Portugal
cvalls@math.ist.utl.pt*

Received 26 June 2012

Accepted 26 September 2012

Published 31 December 2012

We characterize the Darboux first integrals of a simplified Friedman–Robertson–Walker Hamiltonian system depending on one parameter.

Keywords: Darboux first integrals; Darboux polynomials; Friedman–Robertson–Walker Hamiltonian system.

PACS: Primary: 34A05, 34A34, 34C14

1. Introduction and Statement of the Main Result

Given a system of ordinary differential equations depending on parameters in general is very difficult to recognize for which values of the parameters the equations have first integrals because there are no satisfactory methods to answer this question.

In this paper we study the first integrals of the simplified Friedman–Robertson–Walker Hamiltonian differential system in \mathbb{R}^4

$$\begin{aligned}
 \dot{x} &= -p_x, \\
 \dot{y} &= p_y, \\
 \dot{p}_x &= x - bxy^2, \\
 \dot{p}_y &= -y - bx^2y,
 \end{aligned} \tag{1}$$

where $b \in \mathbb{R}$ is a parameter and the dot denotes derivative with respect to time t . The Hamiltonian of this system is

$$H_0 = \frac{1}{2}(p_y^2 - p_x^2) + \frac{1}{2}(y^2 - x^2) + \frac{b}{2}x^2y^2.$$

System (1) models a universe, filled by a conformally coupled but massive real scalar field, see [2] for more details where the authors present analytical and numerical evidence of the existence of chaotic motion. For a more general point of view on the Friedman–Robertson–Walker system see for instance the works [5, 7] and the references quoted therein.

The vector field associated to system (1) is

$$\mathcal{X} = -p_x \frac{\partial}{\partial x} + p_y \frac{\partial}{\partial y} + x(1 - by^2) \frac{\partial}{\partial p_x} - y(1 + bx^2) \frac{\partial}{\partial p_y}.$$

Let $U \subset \mathbb{C}^4$ be an open set. We say that the non-constant function $H: U \rightarrow \mathbb{C}$ is a *first integral* of the polynomial vector field \mathcal{X} on U if $H(x(t), y(t), p_x(t), p_y(t))$ is constant for all values of t for which the solution $(x(t), y(t), p_x(t), p_y(t))$ of \mathcal{X} is defined on U . Clearly H is a first integral of \mathcal{X} on U if and only if

$$\mathcal{X}H = -p_x \frac{\partial H}{\partial x} + p_y \frac{\partial H}{\partial y} + x(1 - by^2) \frac{\partial H}{\partial p_x} - y(1 + bx^2) \frac{\partial H}{\partial p_y} = 0$$

on U .

In this paper we want to study the so-called Darboux first integrals of the Friedman–Robertson–Walker polynomial differential systems (1), using the Darboux theory of integrability (originated in the papers [4]). For a present state of this theory see [6, Chapter 8], the paper [9], and the references quoted in them.

We emphasize that the study of the existence of first integrals is a classical problem in the theory of differential systems, because the knowledge of first integrals of a differential system can be very useful in order to understand and simplify the topological structure of their orbits. Thus, their existence or not can also be viewed as a measure of the complexity of a differential system.

We recall that a first integral is of *Darboux* type if it is of the form

$$f_1^{\lambda_1} \cdots f_p^{\lambda_p} F_1^{\mu_1} \cdots F_q^{\mu_q}, \tag{2}$$

where f_1, \dots, f_p are Darboux polynomials (see Sec. 2 for a definition), F_1, \dots, F_q are exponential factors (see Sec. 2 for a definition), and $\lambda_j, \mu_k \in \mathbb{C}$ for all j and k .

The functions of the form (2) are called *Darboux* functions, and they are the base of the Darboux theory of integrability, which looks when these functions are first integrals or integrating factors. In this last case, the first integrals associated to integrating factors given by Darboux functions are the Liouvillian first integrals, see for more details [6, 9].

The Darboux theory of integrability is essentially an algebraic theory of integrability based on the invariant algebraic hypersurfaces that a polynomial differential system has. In fact to every Darboux polynomial there is associated some invariant algebraic hypersurface (see again Sec. 2), and the exponential factors appear when an invariant algebraic surface has multiplicity larger than 1, for more details see [3, 6, 9]. As far as we know is the unique theory of integrability which is developed for studying the first integrals of polynomial

differential systems. In general the other theories of integrability do not need that the differential system be polynomial.

The main result of this paper is the following.

Theorem 1. *The unique first integrals of Darboux type of the Friedman–Robertson–Walker Hamiltonian system (1) with $b \neq 0$ are functions of Darboux type in the variable H_0 .*

We prove Theorem 1 in Sec. 3.

Since the Darboux theory of integrability of a polynomial differential system is based on the existence of Darboux polynomials and their multiplicity, the study of the existence or not of Darboux first integrals needs to look for the Darboux polynomials. These will be done in the next Theorems 7 and 8, whose results are the main steps for proving Theorem 1.

We must mention that any comment on the existence or non-existence of first integrals additional to the Hamiltonian itself appears in [2]. In that paper the authors are mainly concerned with the chaotic behavior of the Friedman–Robertson–Walker polynomial Hamiltonian system. Of course, roughly speaking the chaotic motion is against the existence of first integrals.

The Hamiltonian system (1) is *completely integrable* if it has two independent first integrals H_1 and H_2 in involution. That is, H_1 and H_2 are *independent* if their gradients are linearly independent over a set of full Lebesgue measure in \mathbb{C}^4 . Moreover, H_1 and H_2 are in *involution* if the Poisson bracket $\{H_i, H_0\} = 0$ for $i \in \{1, 2\}$.

Corollary 2. *The Friedman–Robertson–Walker Hamiltonian system (1) is completely integrable with first integrals of Darboux type if and only if $b = 0$.*

Corollary 2 is proved at the end of Sec. 3.

2. Basic Results

Let $h = h(x, y, p_x, p_y) \in \mathbb{C}[x, y, p_x, p_y] \setminus \mathbb{C}$. As usual $\mathbb{C}[x, y, p_x, p_y]$ denotes the ring of all complex polynomials in the variables x, y, p_x, p_y . We say that h is a *Darboux polynomial* of system (1) if it satisfies

$$\mathcal{X}h = Kh,$$

the polynomial $K = K(x, y, p_x, p_y) \in \mathbb{C}[x, y, p_x, p_y]$ is called *the cofactor* of h and has degree at most 2. Every Darboux polynomial h defines an *invariant algebraic hypersurface* $h = 0$, i.e. if a trajectory of system (1) has a point in $h = 0$, then the whole trajectory is contained in $h = 0$, see for more details [6]. When $K = 0$ the Darboux polynomial h is a polynomial first integral.

An *exponential factor* E of system (1) is a function of the form $E = \exp(g/h) \notin \mathbb{C}$ with $g, h \in \mathbb{C}[x, y, p_x, p_y]$ satisfying $(g, h) = 1$ and

$$\mathcal{X}E = LE,$$

for some polynomial $L = L(x, y, p_x, p_y)$ of degree at most 2, called the *cofactor* of E .

A geometrical meaning of the notion of exponential factor is given by the next result.

Proposition 3. *If $E = \exp(g/h)$ is an exponential factor for the polynomial differential system (1) and h is not a constant polynomial, then $h = 0$ is an invariant algebraic*

hypersurface, and eventually e^g can be exponential factors, coming from the multiplicity of the infinite invariant hyperplane.

The proof of Proposition 3 can be found in [3, 8]. We explain a little the last part of the statement of Proposition 3. If we extend to the projective space $\mathbb{P}\mathbb{R}^4$ the polynomial differential system (1) defined in the affine space \mathbb{R}^4 , then the hyperplane at infinity always is invariant by the flow of the extended differential system. Moreover, if this invariant hyperplane has multiplicity higher than 1, then it creates exponential factors of the form e^g , see for more details [8].

Theorem 4. *Suppose that the polynomial vector field \mathcal{X} of degree m defined in \mathbb{C}^4 admits p invariant algebraic hypersurfaces $f_i = 0$ with cofactors K_i , for $i = 1, \dots, p$ and q exponential factors $E_j = \exp(g_j/h_j)$ with cofactors L_j , for $j = 1, \dots, q$. Then there exists $\lambda_i, \mu_j \in \mathbb{C}$ not all zero such that*

$$\sum_{i=1}^p \lambda_i K_i + \sum_{j=1}^q \mu_j L_j = 0$$

if and only if the function of Darboux type

$$f_1^{\lambda_1} \dots f_p^{\lambda_p} E_1^{\mu_1} \dots E_q^{\mu_q}$$

is a first integral of \mathcal{X} .

Theorem 4 is proved in [6].

From Theorem 4 it follows easily the next well-known result.

Corollary 5. *The existence of a rational first integral for a polynomial differential system implies either the existence of a polynomial first integral or the existence of two Darboux polynomials with the same nonzero cofactor.*

The following result is well-known.

Lemma 6. *Assume that $\exp(g_1/h_1), \dots, \exp(g_r/h_r)$ are exponential factors of some polynomial differential system*

$$\begin{aligned} x' &= P(x, y, p_x, p_y), & y' &= Q(x, y, p_x, p_y), \\ p'_x &= R(x, y, p_x, p_y), & p'_y &= U(x, y, p_x, p_y) \end{aligned} \tag{3}$$

with $P, Q, R, U \in \mathbb{C}[x, y, p_x, p_y]$ with cofactors L_j for $j = 1, \dots, r$. Then

$$\exp(G) = \exp(g_1/h_1 + \dots + g_r/h_r)$$

is also an exponential factor of system (3) with cofactor $L = \sum_{j=1}^r L_j$.

3. Proof of Theorem 1

According with Sec. 2 for proving Theorem 1 we need to characterize the Darboux polynomials of system (1).

The following result characterizes the polynomial first integrals of system (1) with $b \neq 0$, i.e. it characterizes the Darboux polynomial with zero cofactor.

Theorem 7. *The unique polynomial first integrals of system (1) with $b \neq 0$ are polynomials in the variable H_0 .*

Proof. Doing the change of variables

$$z_1 = p_x - p_y, \quad z_2 = p_x + p_y, \tag{4}$$

system (1) becomes

$$\begin{aligned} \dot{x} &= -\frac{1}{2}(z_1 + z_2), \\ \dot{y} &= -\frac{1}{2}(z_1 - z_2), \\ \dot{z}_1 &= x + y + bxy(x - y), \\ \dot{z}_2 &= x - y - bx(x + y), \end{aligned} \tag{5}$$

and H_0 writes as

$$H_0 = \frac{1}{2}(y^2 - x^2 - z_1 z_2 + bx^2 y^2).$$

Now we restrict to $H_0 = 0$. We get that

$$z_2 = \frac{bx^2 y^2 + y^2 - x^2}{z_1}. \tag{6}$$

Then system (5) on $H_0 = 0$ becomes, after the rescaling by $d\tau = z_1 dt$:

$$\begin{aligned} x' &= -\frac{1}{2}z_1^2 - \frac{1}{2}(bx^2 y^2 + y^2 - x^2), \\ y' &= -\frac{1}{2}z_1^2 + \frac{1}{2}(bx^2 y^2 + y^2 - x^2), \\ z_1' &= z_1(x + y + bxy(x - y)), \end{aligned} \tag{7}$$

where the prime denotes the derivative with respect to the variable τ . If $f = f(x, y, p_x, p_y)$ is a polynomial first integral of system (1) then, if we denote by $g = g(x, y, z_1, z_2)$ the polynomial first integral f written in the new variables (x, y, z_1, z_2) , we have that g is also a polynomial first integral of the differential system (5). Furthermore, if we denote by $h = h(x, y, z_1)$ the polynomial first integral g restricted to the invariant hypersurface (6), then h is a rational first integral of the differential system (7) in the sense that it is the quotient of a polynomial in the variables (x, y, z_1) and a power of z_1 . Furthermore h satisfies

$$\begin{aligned} &\left(-\frac{1}{2}z_1^2 - \frac{1}{2}(bx^2 y^2 + y^2 - x^2)\right) \frac{\partial h}{\partial x} \\ &+ \left(-\frac{1}{2}z_1^2 + \frac{1}{2}(bx^2 y^2 + y^2 - x^2)\right) \frac{\partial h}{\partial y} + z_1(x + y + bxy(x - y)) \frac{\partial h}{\partial z_1} = 0. \end{aligned} \tag{8}$$

We write

$$h = \sum_{j=0}^n h_j(x, y, z_1), \tag{9}$$

where each h_j is the quotient of a polynomial in the variable (x, y, z_1) with degree j and the denominator is a power in the variable z_1 . Here we define the degree of h_j as j . Hence, computing the terms of degree $n + 4$ we have

$$\frac{1}{2}bx^2y^2 \left(\frac{\partial h_n}{\partial y} - \frac{\partial h_n}{\partial x} \right) + z_1bxy(x - y) \frac{\partial h_n}{\partial z_1} = 0.$$

We write it as

$$L[h_n] = 0, \quad \text{where } L = \frac{1}{2}bx^2y^2 \left(\frac{\partial}{\partial y} - \frac{\partial}{\partial x} \right) + z_1bxy(x - y) \frac{\partial}{\partial z_1}. \tag{10}$$

The characteristic equations associated to the linear partial differential operator L are

$$\frac{dx}{dy} = -1, \quad \frac{dy}{dz_1} = \frac{xy}{2z_1(x - y)}.$$

The system has the general solution

$$x + y = d_1, \quad \frac{y^{2x}}{e^{2y}z_1^x} = d_2,$$

where d_1 and d_2 are constants of integration. According to this, we make the change of variables

$$u = x + y, \quad v = \frac{y^{2x}}{e^{2y}z_1^x}, \quad w = y. \tag{11}$$

Its inverse transformation is

$$x = u - w, \quad z = y^2(e^{2w}v)^{1/(w-u)}, \quad y = w. \tag{12}$$

Under the change of variables (11) and (12) Eq. (10) becomes the following ordinary differential equation (for fixed u and v):

$$\frac{b}{2}w^2(u - w)^2 \frac{\partial \bar{h}_n}{\partial w} = 0,$$

where \bar{h}_n is h_n written in the variables u, v and w . Solving it we have

$$\bar{h}_n = \bar{h}_n(u, v) = h_n \left(x + y, \frac{y^{2x}}{e^{2y}z_1^x} \right).$$

Since the numerator of h_n is a polynomial of degree n we must have

$$h_n = \sum_{k=0}^n \alpha_k (x + y)^k, \quad \alpha_k \in \mathbb{R}. \tag{13}$$

Now using the transformations (11) and (12) and working in a similar manner to solve \bar{h}_n , computing the terms of degree $n + 3$, we get

$$h_{n-1} = \sum_{k=0}^{n-1} \beta_k (x + y)^k, \quad \beta_k \in \mathbb{R}. \tag{14}$$

Now, using the transformations (11) and (12) and working in a similar manner to solve \bar{h}_{n-2} , computing the terms of degree $n + 2$, we get

$$\frac{b}{2} w^2 (u - w)^2 \frac{\partial \bar{h}_{n-2}}{\partial w} = w^4 (e^{2wv})^{2/(w-u)} \sum_{k=0}^n k \alpha_k u^{k-1}.$$

Integrating this equation we obtain that $\bar{h}_{n-2} = \bar{h}_{n-2}(u, v, w)$ is

$$\begin{aligned} & \frac{1}{b} \left(\sum_{k=0}^n k \alpha_k u^{k-1} \right) \left(\frac{(e^{2wv})^{2/(w-u)} (5u^2 - 8wu + 4w^2 + 2(u - w) \log(e^{2wv}))}{-2u + 2w - \log(e^{2wv})} \right. \\ & \left. - 4e^4 \text{Ei} \left(\frac{2 \log(e^{2wv})}{w - u} - 4 \right) (3u - 2w + \log(e^{2wv})) \right) + k_{n-2}(u, v), \end{aligned}$$

where $\text{Ei}(z)$ is the exponential integral function, see for instance [1]. Going back to the variables x, y and z_1 we obtain that $h_{n-2}(x, y, z_1)$ is

$$\begin{aligned} & \frac{1}{b} \left(\sum_{k=0}^n k \alpha_k (x + y)^{k-1} \right) \left(- \frac{(y^{2x} z_1^{-x})^{-2/x} (5x^2 + 2yx + 2 \log(y^{2x} z_1^{-x}) x + y^2)}{2x + \log(y^{2x} z_1^{-x})} \right. \\ & \left. - 4e^4 \text{Ei} \left(- \frac{2 \log(y^{2x} z_1^{-x})}{x} - 4 \right) (3x + y + \log(y^{2x} z_1^{-x})) \right) + k_{n-2} \left(x + y, \frac{y^{2x}}{e^{2y} z_1^x} \right). \end{aligned}$$

Since the numerator of $h_{n-2}(x, y, z_1)$ is a polynomial of degree $n - 2$ we must have

$$\sum_{k=0}^n k \alpha_k (x + y)^{k-1} = 0.$$

So $\alpha_k = 0$ for $k = 1, \dots, n$. Then $h_n = \alpha_0$, which implies that $h = \alpha_0/z_1^m$ for some non-negative integer m . Then, from (8) we get

$$- \frac{m \alpha_0}{z_1^m} (x + y + bxy(x - y)) = 0.$$

Therefore $m \alpha_0 = 0$, and consequently $h = \text{constant}$, in contradiction with the fact that h is a first integral. □

Now we characterize the existence of Darboux polynomials with nonzero cofactor when $b \neq 0$.

Theorem 8. *System (1) with $b \neq 0$ has no Darboux polynomial with nonzero cofactor.*

Proof. Let f be a Darboux polynomial with a nonzero cofactor K . Since K is a polynomial of degree at most 2, we write it as

$$K = a_0 + a_1x + a_2y + a_3p_x + a_4p_y + a_5x^2 + a_6xy + a_7xp_x + a_8xp_y + a_9y^2 + a_{10}yp_x + a_{11}yp_y + a_{12}p_x^2 + a_{13}p_xp_y + a_{14}p_y^2.$$

Note that system (1) is invariant under the involution $\sigma: \mathbb{C}[x, y, p_x, p_y] \rightarrow \mathbb{C}[x, y, p_x, p_y]$ defined by

$$\sigma(x) = -x, \quad \sigma(y) = -y, \quad \sigma(p_x) = -p_x, \quad \sigma(p_y) = -p_y.$$

If f is a Darboux polynomial of system (1) then $g = f \cdot \sigma(f)$ is a Darboux polynomial of system (1) invariant by σ and with cofactor

$$K_1 = K + \sigma(K) = 2(a_0 + a_5x^2 + a_6xy + a_7xp_x + a_8xp_y + a_9y^2 + a_{10}yp_x + a_{11}yp_y + a_{12}p_x^2 + a_{13}p_xp_y + a_{14}p_y^2).$$

Let f be a Darboux polynomial of system (1) with nonzero cofactor K . If f is invariant by σ then $K = K_1$ and we take $g = f$. If f is not invariant by σ then we consider $g = f \cdot \sigma(f)$ a new Darboux polynomial of system (1) invariant by σ and with cofactor K_1 . We have

$$-p_x \frac{\partial g}{\partial x} + p_y \frac{\partial g}{\partial y} + x(1 - by^2) \frac{\partial g}{\partial p_x} - y(1 + bx^2) \frac{\partial g}{\partial p_y} = K_1g. \tag{15}$$

We write g as a polynomial in the variable p_x as

$$g = \sum_{i=0}^k g_i(x, y, p_y)p_x^i,$$

where each g_i is a polynomial in the variables x, y and p_y . Without loss of generality we can assume that $g_k(x, y, p_y) \neq 0$. Then computing the coefficient of p_x^{k+2} in (15) we get

$$2a_{12}g_k = 0 \quad \text{which yields } a_{12} = 0.$$

Now computing the coefficient of p_x^{k+1} in (15) we obtain

$$-\frac{\partial g_k}{\partial x} = 2(a_7x + a_{10}y + a_{13}p_y)g_k.$$

Solving it we get

$$g_k = K_k(y, p_y)e^{a_7x^2 + 2a_{10}yx + 2a_{13}p_yx}.$$

Since g_k must be a polynomial we get $a_7 = a_{10} = a_{13} = 0$.

Now we write g as a polynomial in the variable p_y as

$$g = \sum_{i=0}^m \bar{g}_i(x, y, p_x)p_y^i,$$

where each \bar{g}_i is a polynomial in the variables x, y and p_x . Without loss of generality we can assume that $\bar{g}_m(x, y, p_x) \neq 0$. Then computing the coefficient of p_y^{m+2} in (15) we get

$$2a_{14}\bar{g}_m = 0 \quad \text{which yields } a_{14} = 0.$$

Now computing the coefficient of p_y^{m+1} in (15) we obtain

$$\frac{\partial \bar{g}_m}{\partial y} = 2(a_8x + a_{11}y)\bar{g}_m.$$

Solving it we get

$$\bar{g}_m = K_m(x, p_x)e^{2a_8xy+a_{11}y^2}.$$

Since \bar{g}_m must be a polynomial we get $a_8 = a_{11} = 0$. Hence $K_1 = 2(a_0 + a_5x^2 + a_6xy + a_9y^2)$. Then

$$-p_x \frac{\partial g}{\partial x} + p_y \frac{\partial g}{\partial y} + x(1 - by^2) \frac{\partial g}{\partial p_x} - y(1 + bx^2) \frac{\partial g}{\partial p_y} = 2(a_0 + a_5x^2 + a_6xy + a_9y^2)g.$$

Now we write $g = \sum_{j=0}^{\ell} g_j(x, y, p_x, p_y)$ where each g_j is a homogeneous polynomial of degree j . Then computing the terms of degree $\ell + 2$ we get

$$-bxy \left(y \frac{\partial g_{\ell}}{\partial p_x} + x \frac{\partial g_{\ell}}{\partial p_y} \right) = 2(a_5x^2 + a_6xy + a_9y^2)g_{\ell}.$$

Solving it we get

$$g_{\ell} = K \left(x, y, \frac{-xp_x + yp_y}{y} \right) e^{-\frac{2px(a_5x^2+a_6xy+a_9y^2)}{by^2}}.$$

Since it must be a polynomial we get $a_5 = a_6 = a_9 = 0$. Then $K_1 = 2a_0$.

Now proceeding as in the proof of Theorem 7 introducing the change of variables (4) and the restriction $H_0 = 0$ (see (7)) we get that $g(x, y, p_x, p_y) = h(x, y, z_1)$ with

$$\begin{aligned} & \left(-\frac{1}{2}z_1^2 - \frac{1}{2}(bx^2y^2 + y^2 - x^2) \right) \frac{\partial h}{\partial x} + \left(-\frac{1}{2}z_1^2 + \frac{1}{2}(bx^2y^2 + y^2 - x^2) \right) \frac{\partial h}{\partial y} \\ & + z_1(x + y + bxy(x - y)) \frac{\partial h}{\partial z_1} = 2a_0z_1h. \end{aligned} \tag{16}$$

We write h as in (9) and we get h_n and h_{n-1} as in (13) and (14). Now, using the transformations (11) and (12) and working in a similar manner to solve \bar{h}_{n-2} , computing the terms of degree $n + 2$ in (16), we get

$$\begin{aligned} \frac{b}{2}w^2(u - w)^2 \frac{\partial \bar{h}_{n-2}}{\partial w} &= w^4(e^{2w}v)^{2/(w-u)} \sum_{k=0}^n k\alpha_k u^{k-1} \\ &+ 2a_0w^2(e^{2w}v)^{1/(w-u)} \sum_{k=0}^n \alpha_k u^k. \end{aligned}$$

Integrating this equation and going back to the variables x, y and z_1 we get that $h_{n-2}(x, y, z_1)$ is

$$\begin{aligned} & \frac{1}{b(-2y + 2(x + y) + \log(y^{2x} z_1^{-x}))} \left(-4a_0(y^{2x} z_1^{-x})^{-1/x} \sum_{k=0}^n \alpha_k(x + y)^k \right. \\ & - \left((4y^2 - 8(x + y)y + 5(x + y)^2 + 2x \log(y^{2x} z_1^{-x})) (y^{2x} z_1^{-x})^{-2/x} \right. \\ & + 4e^4 \text{Ei} \left(-\frac{2 \log(y^{2x} z_1^{-x})}{x} - 4 \right) (-2y + 2(x + y) + \log(y^{2x} z_1^{-x})) \\ & \left. \left. \times (-2y + 3(x + y) + \log(y^{2x} z_1^{-x})) \right) \sum_{k=0}^n k \alpha_k(x + y)^{k-1} \right) + k_{n-2} \left(x + y, \frac{y^{2x}}{e^{2y} z_1^x} \right). \end{aligned}$$

Since the numerator of h_{n-2} must be a polynomial of degree $n - 2$ we have

$$\sum_{k=0}^n k \alpha_k(x + y)^{k-1} = 0, \quad \text{and} \quad a_0 \sum_{k=0}^n \alpha_k(x + y)^k = 0.$$

This implies that $\alpha_k = 0$ for $k = 1, \dots, n$ and $a_0 \alpha_0 = 0$.

If $\alpha_0 = 0$ then $h = 0$, in contradiction that h is a Darboux polynomial. Therefore $a_0 = 0$ and $K_1 = 0$. Thus $g = f \cdot \sigma(f)$ is a polynomial first integral. By Theorem 7 the unique polynomial first integrals are polynomials in the variable H_0 , we must have that f is a polynomial in the variable H_0 which is not possible because f is a Darboux polynomial of system (1) with nonzero cofactor. This concludes the proof of the theorem. \square

Theorem 9. *The unique rational first integrals of system (1) with $b \neq 0$ are rational functions in the variable H_0 .*

Proof. It follows directly from Corollary 5 and Theorems 7 and 8. \square

Now we proceed as in the proof of Theorem 8.

Proof of Theorem 1. It follows from Theorems 4, 7 and 8 and Proposition 3 that in order to have a first integral of Darboux type we must have q exponential factors $E_j = \exp(g_j/h_j(H_0))$ with cofactors L_j such that $\sum_{j=1}^q \mu_j L_j = 0$. Let $G = \sum_{j=1}^q \mu_j g_j/h_j(H_0)$, then $E = \exp(G)$ is an exponential factor of system (1) with cofactor $L = \sum_{j=1}^q \mu_j L_j$ (see Lemma 6) and G satisfies that $\mathcal{X}G = 0$, that is, G must be a rational first integral of system (1). By Theorem 9 it must be a rational function in the variable H_0 . \square

Proof of Corollary 2. When $b = 0$ it easy to check that $x^2 + p_x^2$ and $y^2 + p_y^2$ are two independent first integrals in involution for the Hamiltonian system (1). Note that both first integrals are given by functions of Darboux type.

When $b \neq 0$ Theorem 1 shows that there exists a unique independent first integral of Darboux type. So, the corollary is proved. \square

Acknowledgments

The first author was partially supported by the MICINN/FEDER grant MTM2008–03437, AGAUR grant 2009SGR-410 and ICREA Academia. The second author has been partially supported by FCT through the project PTDC/MAT/117106/2010 and through CAMGD, Lisbon.

References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series, 55 (Government Printing Office, Washington, DC, 1964).
- [2] E. Calzeta and C. E. Hasi, Chaotic Friedman–Robertson–Walker cosmology, *Class. Quantum Gravity* **10** (1993) 1825–1841.
- [3] C. Christopher, J. Llibre and J. V. Pereira, Multiplicity of invariant algebraic curves in polynomial vector fields, *Pacific J. Math.* **229** (2007) 63–117.
- [4] G. Darboux, Mémoire sur les équations différentielles algébriques du premier ordre et du premier degré (Mélanges), *Bull. Sci. Math. 2^{ème} série* **2** (1878) 60–96; 123–144; 151–200.
- [5] C. Dariescu and M. A. Dariescu, Vacuum electromagnetic modes in Friedman–Robertson–Walker universe, *Int. J. Mod. Phys. A* **18** (2003) 5725–5732.
- [6] F. Dumortier, J. Llibre and J. C. Artés, *Qualitative Theory of Planar Differential Systems*, Universitext (Springer-Verlag, New York, 2006).
- [7] K. Kleidis, A. Kuiroukidis, D. B. Papadopoulos and L. Vlahos, Gravitomagnetic instabilities in anisotropically expanding fluids, *Int. J. Mod. Phys. A* **23** (2008) 4467–4484.
- [8] J. Llibre and X. Zhang, Darboux theory of integrability for polynomial vector fields in \mathbb{R}^n taking into account the multiplicity at infinity, *Bull. Sci. Math.* **133** (2009) 765–778.
- [9] J. Llibre and X. Zhang, On the Darboux integrability of the polynomial differential systems, *Qualitative Theory and Dynamical Systems* **11** (2012) 129–144.