



Journal of Nonlinear Mathematical Physics

ISSN (Online): 1776-0852

ISSN (Print): 1402-9251

Journal Home Page: <https://www.atlantis-press.com/journals/jnmp>

A Large Time Asymptotics for Transparent Potentials for the Novikov–Veselov Equation at Positive Energy

A. V. Kazeykina, R. G. Novikov

To cite this article: A. V. Kazeykina, R. G. Novikov (2011) A Large Time Asymptotics for Transparent Potentials for the Novikov–Veselov Equation at Positive Energy, Journal of Nonlinear Mathematical Physics 18:3, 377–400, DOI:

<https://doi.org/10.1142/S1402925111001660>

To link to this article: <https://doi.org/10.1142/S1402925111001660>

Published online: 04 January 2021

Journal of Nonlinear Mathematical Physics, Vol. 18, No. 3 (2011) 377–400

© A. V. Kazeykina and R. G. Novikov

DOI: [10.1142/S1402925111001660](https://doi.org/10.1142/S1402925111001660)

A LARGE TIME ASYMPTOTICS FOR TRANSPARENT POTENTIALS FOR THE NOVIKOV–VESELOV EQUATION AT POSITIVE ENERGY

A. V. KAZEYKINA

*Centre des Mathématiques Appliquées
Ecole Polytechnique, Palaiseau, 91128, France*

and

*Lomonosov Moscow State University, GSP-1
Leninskie Gory, Moscow, 119991, Russia
kazeykina@cmap.polytechnique.fr*

R. G. NOVIKOV

*CNRS (UMR 7641), Centre des Mathématiques Appliquées
Ecole Polytechnique, Palaiseau, 91128, France*

and

*International Institute of Earthquake Prediction
Theory and Mathematical Geophysics, Russian Academy of Sciences
Profsoyuznaya str., 84/32, Moscow, 117997, Russia
novikov@cmap.polytechnique.fr*

Received 14 October 2010

Accepted 9 February 2011

In the present paper we begin studies on the large time asymptotic behavior for solutions of the Cauchy problem for the Novikov–Veselov equation (an analog of KdV in $2 + 1$ dimensions) at positive energy. In addition, we are focused on a family of reflectionless (transparent) potentials parameterized by a function of two variables. In particular, we show that there are no isolated soliton type waves in the large time asymptotics for these solutions in contrast with well-known large time asymptotics for solutions of the KdV equation with reflectionless initial data.

Keywords: Transparent potentials; Novikov–Veselov equation; KdV in $2 + 1$ dimensions; large time asymptotics.

1. Introduction

We consider the scattering problem for the two-dimensional Schrödinger equation

$$-\Delta\psi + v(x)\psi = E\psi, \quad x \in \mathbb{R}^2, \quad E = E_{\text{fixed}} > 0 \quad (1.1)$$

at a fixed positive energy, where

$$v(x) = \overline{v(x)}, \quad v \in L^\infty(\mathbb{R}^2),$$

$$|v(x)| < q(1 + |x|)^{-2-\varepsilon}, \quad \varepsilon > 0, \quad q > 0. \tag{1.2}$$

It is known that for any $k \in \mathbb{R}^2$, such that $k^2 = E$, there exists a unique continuous solution $\psi^+(x, k)$ of Eq. (1.1) with the following asymptotics

$$\psi^+(x, k) = e^{ikx} - i\pi\sqrt{2\pi}e^{-\frac{i\pi}{4}} f\left(k, |k|\frac{x}{|x|}\right) \frac{e^{i|k||x|}}{\sqrt{|k||x|}}$$

$$+ o\left(\frac{1}{\sqrt{|x|}}\right), \quad \text{as } |x| \rightarrow +\infty \left(\text{uniformly in } \frac{x}{|x|}\right) \tag{1.3}$$

for some *a priori* unknown function f .

This solution describes scattering of incident plane wave e^{ikx} on the potential $v(x)$. The function $f = f(k, l)$, $(k, l) \in M_E = \{k \in \mathbb{R}^2, l \in \mathbb{R}^2: k^2 = l^2 = E\}$, arising in (1.3), is the scattering amplitude for $v(x)$ in the framework of Eq. (1.1). Actually, formula (1.3) can be considered as one of the mathematical definitions of the scattering amplitude. For more information concerning the aforementioned facts see [2, 4, 10].

In the present paper we are focused on transparent (or invisible) potentials v for Eq. (1.1). We say that v is transparent if its scattering amplitude f is identically zero at fixed energy E , i.e. $f \equiv 0$ on M_E . We consider transparent potentials for Eq.(1.1) as analogs of reflectionless potentials for the one-dimensional Schrödinger equation at all positive energies; see, for example, [11, 15] as regards reflectionless potentials in dimension one.

In [4] it was shown that

- (1) There are no nonzero transparent potentials for Eq. (1.1), where

$$v(x) = \overline{v(x)}, \quad v \in L^\infty(\mathbb{R}^2), \quad |v(x)| < \alpha e^{\beta|x|}, \quad \alpha > 0, \quad \beta > 0. \tag{1.4}$$

- (2) There is a family of nonzero transparent potentials for Eq. (1.1), where

$$v(x) = \overline{v(x)}, \quad v \in \mathcal{S}(\mathbb{R}^2), \tag{1.5}$$

and \mathcal{S} denotes the Schwartz class.

In the present paper, in addition to the scattering problem for (1.1), we consider its isospectral deformation generated by the following (2 + 1)-dimensional analog of the KdV equation:

$$\partial_t v = 4 \operatorname{Re} (4\partial_z^3 v + \partial_z(vw) - E\partial_z w), \quad \partial_{\bar{z}} w = -3\partial_z v, \quad v = \bar{v},$$

$$v = v(x, t), \quad w = w(x, t), \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad t \in \mathbb{R}, \tag{1.6}$$

where

$$\partial_t = \frac{\partial}{\partial t}, \quad \partial_z = \frac{1}{2} \left(\frac{\partial}{\partial x_1} - i \frac{\partial}{\partial x_2} \right), \quad \partial_{\bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x_1} + i \frac{\partial}{\partial x_2} \right).$$

Equation (1.6) is contained implicitly in the paper of Manakov [7] as an equation possessing the following representation:

$$\frac{\partial(L - E)}{\partial t} = [L - E, A] + B(L - E), \tag{1.7}$$

(Manakov $L - A - B$ triple), where $L = -\Delta + v(x, t)$ or, in other words, L (at fixed t) is the Schrödinger operator of (1.1), A and B are suitable differential operators of the third and the zero order respectively. Equation (1.6) was written in an explicit form by Novikov and Veselov in [13, 14], where higher analogs of (1.6) were also constructed.

Note that both Kadomtsev–Petviashvili equations can be obtained from (1.6) by considering an appropriate limit $E \rightarrow \pm\infty$, see [16, 2].

In terms of scattering data the nonlinear equation (1.6), where $E = E_{\text{fixed}} > 0$,

$$\begin{aligned} v &\text{ is sufficiently regular and has sufficient decay as } |x| \rightarrow \infty, \\ w &\text{ is decaying as } |x| \rightarrow \infty, \end{aligned} \tag{1.8}$$

takes the form (2.1) and (2.2) (see Sec. 2) and, in particular,

$$\frac{\partial f(k, l, t)}{\partial t} = 2i[k_1^3 - 3k_1k_2^2 - l_1^3 + 3l_1l_2^2] f(k, l, t), \tag{1.9}$$

$k = (k_1, k_2) \in \mathbb{R}^2, l = (l_1, l_2) \in \mathbb{R}^2, k^2 = l^2 = E$, where $f(\cdot, \cdot, t)$ is the scattering amplitude for $v(\cdot, t)$. Equation (1.9) implies that the nonlinear evolution equation (1.6) under assumptions (1.8) preserves the transparency (or invisibility) property of $v(x, 0)$ in the framework of the scattering problem for (1.1).

In the present paper we begin studies on the large time asymptotic behavior for solutions of the Cauchy problem for (1.6) under assumptions (1.8). We give a large time estimate for the family of solutions of (1.6) with $E = E_{\text{fixed}} > 0$ given by (2.12)–(2.14) and parameterized by a function of two variables. All potentials of this family are transparent at fixed t and E . In addition, this family contains all solutions of (1.6) with $E = E_{\text{fixed}} > 0$ such that:

- $v(\cdot, 0) \in \mathcal{S}(\mathbb{R}^2)$, where \mathcal{S} denotes the Schwartz class, (1.10)

- $v(\cdot, 0)$ is transparent for (1.1), (1.11)

- $v(\cdot, 0)$ satisfies (1.2), where (1.12)

$$q < Q(E, \varepsilon) \quad \text{“small norm” condition,} \tag{1.13}$$

and Q is a special real function with the properties $Q(E, \varepsilon) > 0$ as $E > 0, \varepsilon > 0, Q(E, \varepsilon) \rightarrow +\infty$ for fixed $\varepsilon > 0$ as $E \rightarrow +\infty$,

- $v, w \in C^\infty(\mathbb{R}^2 \times \mathbb{R}), \partial_x^j v(x, t) = O(|x|^{-3}), w(x, t) = o(1), |x| \rightarrow \infty$, for $t \in \mathbb{R}, j \in (\mathbb{N} \cup 0)^2$.

The aforementioned family of solutions of (1.6) was considered for the first time in [4]. In the present work we prove the following estimate

$$|v(x, t)| \leq \frac{\text{const}(v) \ln(3 + |t|)}{1 + |t|}, \quad x \in \mathbb{R}^2, \quad t \in \mathbb{R} \tag{1.14}$$

for each v of this family.

Estimate (1.14) implies that there are no isolated soliton type waves in the large time asymptotics for $v(x, t)$, in contrast with large time asymptotics for solutions of the KdV equation with reflectionless initial data.

Apparently, it is not difficult to obtain the estimate (1.14), where the right-hand side is replaced by $\frac{\text{const}(v)}{1+|t|}$ and even to give precise expression for the leading term of the asymptotics of $v(x, t)$ as $t \rightarrow \infty$. However, already (1.14) in its present form implies the aforementioned absence of isolated soliton-type waves in the large-time asymptotics for $v(x, t)$.

Studies on the large time asymptotics for solutions of the Cauchy problem for the Kadomtsev–Petviashvili equations were fulfilled in [8, 5, 6].

Estimate (1.14) is proved in Secs. 3 and 4, using the stationary phase method (see, for example, [1, Chapter III]), techniques developed in [4] and [6] and an analysis of some cubic algebraic equation depending on a complex parameter.

2. Transparent Potentials and Inverse Scattering Transform

In order to study the large time behavior of the family of transparent potentials described in the previous section, we will use the inverse scattering transform for the two-dimensional Schrödinger equation (1.1). In this section we recall some of the results given in the literature in this connection and summarized in [4].

First, we give the complete definition of scattering data for (1.1). Let $k \in \mathbb{C}^2$, $k^2 = E$, $\text{Im } k \neq 0$ and $v(x)$ satisfy conditions (1.2) and (1.13). Then there exists a unique solution of (1.1) such that

$$\psi(x, k) = e^{ikx}(1 + o(1)), \quad |x| \rightarrow \infty.$$

It can be shown ([10]) that for $E \in \mathbb{R}$, $\text{Im } k \neq 0$ the function $\psi(x, k)$ can be expanded as

$$\psi(x, k) = e^{ikx} - \pi \operatorname{sgn}(\operatorname{Im } k_2, \bar{k}_1) e^{ikx} \left(\frac{a(k)}{-k_2 x_1 + k_1 x_2} + \frac{e^{-2i \operatorname{Re } kx} b(k)}{-\bar{k}_2 x_1 + \bar{k}_1 x_2} + o\left(\frac{1}{|x|}\right) \right),$$

$$|x| \rightarrow \infty.$$

For $E > 0$ the function $b(k)$ is considered to be scattering data for (1.1) in addition to the scattering amplitude $f(k, l)$ arising in (1.3). It was shown in [3, 9, 10] that at fixed positive energy $f(k, l)$ and $b(k)$ uniquely determine the potential $v(x)$ satisfying (1.2), (1.13) (while $f(k, l)$ alone is insufficient for this purpose).

If potential $v(x, t)$ satisfies the Novikov–Veselov equation (1.6) under assumptions (1.8), then dynamics of the scattering data is described by the following formulas

$$\frac{\partial b(k, t)}{\partial t} = 2i[k_1^3 + \bar{k}_1^3 - 3k_1 k_2^2 + 3\bar{k}_1 \bar{k}_2^2] b(k, t), \quad k \in \mathbb{C}^2, \operatorname{Im } k \neq 0, k^2 = E, \quad (2.1)$$

$$\frac{\partial f(k, l, t)}{\partial t} = 2i[k_1^3 - 3k_1 k_2^2 - l_1^3 + 3l_1 l_2^2] f(k, l, t), \quad k, l \in \mathbb{R}^2, k^2 = l^2 = E. \quad (2.2)$$

Equation (2.2) implies that the Novikov–Veselov equation preserves the property of transparency. It was also shown in [4] that this equation does not preserve, in general, the property of very fast decay of initial data. It can only be guaranteed that if $v(x, 0) \in \mathcal{S}(\mathbb{R}^2)$ and satisfies (1.2), (1.13), then for every t we have $v(x, t) = O(|x|^{-3})$.

In the present paper we are concerned with transparent potentials, i.e. in the further considerations we assume that $f(k, l, t) \equiv 0$. We will also put

$$E = 1 \tag{2.3}$$

without loss of generality (the case of an arbitrary fixed positive energy may be reduced to (2.3) by scaling transformation). Along with the function $\psi(x, k)$ we will consider the

function $\mu(x, k)$ related to $\psi(x, k)$ by the following expression

$$\psi(x, k) = e^{ikx} \mu(x, k). \tag{2.4}$$

It is also convenient in the two-dimensional scattering theory to introduce new notations

$$z = x_1 + ix_2, \quad \bar{z} = x_1 - ix_2, \quad \lambda = k_1 + ik_2.$$

Then

$$k_1 = \frac{1}{2} \left(\lambda + \frac{1}{\lambda} \right), \quad k_2 = \frac{i}{2} \left(\frac{1}{\lambda} - \lambda \right)$$

and we will consider that

$$\psi = \psi(z, \lambda, t), \quad \mu = \mu(z, \lambda, t), \quad b = b(\lambda, t).$$

In new notations the Schrödinger equation takes the form

$$L\psi = E\psi, \quad L = -4\partial_z\partial_{\bar{z}} + v(z, t), \quad z \in \mathbb{C},$$

Eq. (2.1) is written as

$$\frac{\partial b(\lambda, t)}{\partial t} = i \left(\lambda^3 + \frac{1}{\lambda^3} + \bar{\lambda}^3 + \frac{1}{\bar{\lambda}^3} \right) b(\lambda, t),$$

and (2.4) takes the form

$$\psi(z, \lambda, t) = e^{\frac{i}{2}(\lambda\bar{z} + z/\lambda)} \mu(z, \lambda, t).$$

We also note that in the present paper notation $f(z)$ does not imply that f is holomorphic on z , i.e. we omit the dependency on \bar{z} in the notations.

Let a transparent potential $v(x, t)$, satisfying (1.6) under assumptions (1.8), (2.3), satisfy also at $t = 0$ conditions (1.2), (1.13). Then the function $\mu(z, \lambda, t)$ has the following properties (see [4]):

- (1) $\mu(z, \lambda, t)$ is continuous on $\lambda \in \mathbb{C}$;
- (2) $\forall \lambda \in \mathbb{C}, |\lambda| \neq 1$, the function $\mu(z, \lambda, t)$ satisfies the equation

$$\frac{\partial \mu(z, \lambda, t)}{\partial \bar{\lambda}} = r(\lambda, z, t) \overline{\mu(z, \lambda, t)}, \tag{2.5}$$

where

$$r(\lambda, z, t) = \exp(iS(\lambda, z, t))r(\lambda), \quad r(\lambda) = \frac{\pi \operatorname{sgn}(\lambda\bar{\lambda} - 1)}{\bar{\lambda}} b(\lambda, 0), \tag{2.6}$$

$$S(\lambda, z, t) = F(\lambda, z) + tR(\lambda), \tag{2.7}$$

$$F(\lambda, z) = -\frac{1}{2} \left(\bar{\lambda}z + \lambda\bar{z} + \frac{z}{\lambda} + \frac{\bar{z}}{\bar{\lambda}} \right), \quad R(\lambda) = \lambda^3 + \bar{\lambda}^3 + \frac{1}{\lambda^3} + \frac{1}{\bar{\lambda}^3}; \tag{2.8}$$

- (3) $\mu(z, \lambda, t) \rightarrow 1$ as $\lambda \rightarrow 0, \infty$.

Properties 1–3 uniquely determine $\mu(z, \lambda, t)$ for all $\lambda \in \mathbb{C}$.

Under the same assumptions on the potential and if $v(x, 0) \in \mathcal{S}(\mathbb{R}^2)$, the function $b(\lambda, t)$ has the following properties: for every $t \in \mathbb{R}$

$$b(\cdot, t) \in \mathcal{S}(\mathbb{C}); \tag{2.9}$$

$$b(1/\bar{\lambda}, t) = b(\lambda, t), \quad b(-1/\bar{\lambda}, t) = \overline{b(\lambda, t)}; \tag{2.10}$$

$$\partial_\lambda^m \partial_{\bar{\lambda}}^n b(\lambda, t)|_{|\lambda|=1} = 0 \quad \text{for all } m, n \geq 0. \tag{2.11}$$

The reconstruction of the transparent potential $v(z, t)$ from these scattering data is based on the following scheme.

(1) Function $\mu(z, \lambda, t)$ is constructed as the solution of the following integral equation

$$\mu(z, \lambda, t) = 1 - \frac{1}{\pi} \iint_{\mathbb{C}} r(\zeta, z, t) \overline{\mu(z, \zeta, t)} \frac{d \operatorname{Re} \zeta d \operatorname{Im} \zeta}{\zeta - \lambda}. \tag{2.12}$$

which is uniquely solvable if the scattering data $b(\lambda, t)$ satisfy properties (2.9)–(2.11). Equation (2.12) is obtained from (2.5) by applying the Cauchy–Green formula

$$f(\lambda) = -\frac{1}{\pi} \iint_D (\partial_{\bar{\zeta}} f(\zeta)) \frac{d \operatorname{Re} \zeta d \operatorname{Im} \zeta}{\zeta - \lambda} + \frac{1}{2\pi i} \oint_{\partial D} f(\zeta) \frac{d\zeta}{\zeta - \lambda}.$$

(2) Expanding $\mu(z, \lambda, t)$ as $\lambda \rightarrow \infty$,

$$\mu(z, \lambda, t) = 1 + \frac{\mu_{-1}(z, t)}{\lambda} + o\left(\frac{1}{|\lambda|}\right), \tag{2.13}$$

we define $v(z, t)$ as

$$v(z, t) = 2i\partial_z \mu_{-1}(z, t). \tag{2.14}$$

(3) It can be shown ([3]) that

$$L\psi = \psi,$$

where

$$\begin{aligned} \psi(z, \lambda, t) &= e^{\frac{i}{2}(\lambda\bar{z}+z/\lambda)} \mu(z, \lambda, t), \quad L = -4\partial_z \partial_{\bar{z}} + v(z, t), \\ \overline{v(z, t)} &= v(z, t), \quad v(z, t) \text{ is transparent.} \end{aligned}$$

3. Estimate for the Linearized Case

Consider

$$\begin{aligned} I(t, z) &= \iint_{\mathbb{C}} f(\zeta) \exp(iS(\zeta, z, t)) d \operatorname{Re} \zeta d \operatorname{Im} \zeta, \\ J(t, z) &= -3 \iint_{\mathbb{C}} \bar{\zeta} f(\zeta) \exp(iS(\zeta, z, t)) d \operatorname{Re} \zeta d \operatorname{Im} \zeta, \end{aligned} \tag{3.1}$$

where $f(\zeta) \in L^1(\mathbb{C})$, S is defined by (2.7). If $v(z, t) = I(t, z)$, $w(z, t) = J(t, z)$, where

$$(|\zeta|^3 + |\zeta|^{-3})f(\zeta) \in L^1(\mathbb{C})$$

as a function of ζ , and, in addition,

$$\overline{f(\zeta)} = f(-\zeta) \quad \text{and/or} \quad \overline{f(\zeta)} = -|\zeta|^{-4} f\left(-\frac{1}{\bar{\zeta}}\right),$$

then v, w satisfy the linearized Novikov–Veselov equation (1.6) with $E = 1$. In addition,

$$\hat{v}(p, t) \equiv 0 \quad \text{for } |p| < 2, \quad t \in \mathbb{R},$$

where $\hat{v}(\cdot, t)$ is the Fourier transform of $v(\cdot, t)$, that is $v(\cdot, t)$ is transparent in the Born approximation at energy $E = 1$ for each $t \in \mathbb{R}$.

The goal of this section is to give, in particular, a uniform estimate of the large-time behavior of the integral $I(t, z)$ of (3.1) under the assumptions that

$$\begin{aligned} f &\in C^\infty(\mathbb{C}), \\ \partial_\lambda^m \partial_{\bar{\lambda}}^n f(\lambda) &= \begin{cases} O(|\lambda|^{-\infty}) & \text{as } |\lambda| \rightarrow \infty, \\ O(|\lambda|^\infty) & \text{as } |\lambda| \rightarrow 0, \end{cases} \\ \partial_\lambda^m \partial_{\bar{\lambda}}^n f(\lambda)|_{|\lambda|=1} &= 0 \end{aligned} \tag{3.2}$$

for all $m, n \geq 0$.

We can integrate $I(t, z)$ by parts in a similar way as in Lemma 2.1 in Chapter III of [1] using (3.2) and the fact that

$$\frac{\partial}{\partial \lambda} R(\lambda) \neq 0, \quad \frac{\partial}{\partial \bar{\lambda}} R(\lambda) \neq 0, \quad \text{for } |\lambda| \neq 1, \quad \lambda \neq 0,$$

where R is defined in (2.8). Thus we obtain

$$|I(t, z)| = O\left(\frac{1}{|t|^n}\right), \quad t \rightarrow \infty, \quad n \in \mathbb{N}, \tag{3.3}$$

uniformly on $z \in K$, where K is any compact set of the complex plane. Nevertheless, this is not sufficient to guarantee the absence of soliton-type waves in the large time asymptotics of the potential $v(z, t) = I(t, z)$. So our further reasoning will be devoted to obtaining an estimate like (3.3) for $n = 1$ uniformly on $z \in \mathbb{C}$.

For this purpose we introduce parameter $u = \frac{z}{t}$ and write the integral I in the following form

$$I(t, u) = \iint_{\mathbb{C}} f(\zeta) \exp(itS(u, \zeta)) d\operatorname{Re} \zeta d\operatorname{Im} \zeta, \tag{3.4}$$

where

$$S(u, \zeta) = -\frac{1}{2} \left(\bar{\zeta} u + \zeta \bar{u} + \frac{u}{\zeta} + \frac{\bar{u}}{\bar{\zeta}} \right) + \left(\zeta^3 + \bar{\zeta}^3 + \frac{1}{\zeta^3} + \frac{1}{\bar{\zeta}^3} \right). \tag{3.5}$$

We will start by studying the properties of the stationary points of the function $S(u, \zeta)$. These points satisfy the equation

$$S'_\zeta = -\frac{\bar{u}}{2} + \frac{u}{2\zeta^2} + 3\zeta^2 - \frac{3}{\zeta^4} = 0. \tag{3.6}$$

The degenerate stationary points obey additionally the equation

$$S''_{\zeta\zeta} = -\frac{u}{\zeta^3} + 6\zeta + \frac{12}{\zeta^5} = 0. \tag{3.7}$$

We denote $\xi = \zeta^2$ and

$$Q(u, \xi) = -\frac{\bar{u}}{2} + \frac{u}{2\xi} + 3\xi - \frac{3}{\xi^2}.$$

For each ξ , a root of the function $Q(u, \xi)$, there are two corresponding stationary points of $S(u, \zeta)$, $\zeta = \pm\sqrt{\xi}$.

The function $S'_\zeta(u, \zeta)$ can be represented in the following form

$$S'_\zeta(u, \zeta) = \frac{3}{\zeta^4}(\zeta^2 - \zeta_0^2(u))(\zeta^2 - \zeta_1^2(u))(\zeta^2 - \zeta_2^2(u)). \tag{3.8}$$

We will also use hereafter the following notations:

$$\mathcal{U} = \{u = 6(2e^{-i\varphi} + e^{2i\varphi}), \varphi \in [0, 2\pi)\}$$

and

$$\mathbb{U} = \{u = re^{i\varphi} : 0 \leq r \leq |6(2e^{-i\varphi} + e^{2i\varphi})|, \varphi \in [0, 2\pi)\},$$

the domain limited by the curve \mathcal{U} (see also Fig. 1).

Lemma 3.1. (1) *If $u = 18e^{\frac{2\pi ik}{3}}, k = 0, 1, 2$, then*

$$\zeta_0(u) = \zeta_1(u) = \zeta_2(u) = e^{-\frac{\pi ik}{3}}$$

and $S(u, \zeta)$ has two degenerate stationary points, corresponding to a third-order root of the function $Q(u, \xi)$, $\xi_1 = e^{-\frac{2\pi ik}{3}}$.

(2) *If $u \in \mathcal{U}$ (i.e. $u = 6(2e^{-i\varphi} + e^{2i\varphi})$) and $u \neq 18e^{\frac{2\pi ik}{3}}, k = 0, 1, 2$, then*

$$\zeta_0(u) = \zeta_1(u) = e^{i\varphi/2}, \quad \zeta_2(u) = e^{-i\varphi}.$$

Thus $S(u, \zeta)$ has two degenerate stationary points, corresponding to a second-order root of the function $Q(u, \xi)$, $\xi_1 = e^{i\varphi}$, and two non-degenerate stationary points corresponding to a first-order root, $\xi_2 = e^{-2i\varphi}$.

(3) *If $u \in \text{int}\mathbb{U}$, then*

$$\zeta_i(u) = e^{i\varphi_i}, \quad \text{and} \quad \zeta_i(u) \neq \zeta_j(u) \quad \text{for } i \neq j.$$

In this case the stationary points of $S(u, \zeta)$ are non-degenerate and correspond to the roots of the function $Q(u, \xi)$ with absolute values equal to 1.

(4) *If $u \in \mathbb{C} \setminus \mathbb{U}$, then*

$$\zeta_0(u) = (1 + \omega)e^{i\varphi/2}, \quad \zeta_1(u) = e^{-i\varphi}, \quad \zeta_2(u) = (1 + \omega)^{-1}e^{i\varphi/2}$$

for certain φ and $\omega > 0$.

In this case the stationary points of the function $S(u, \zeta)$ are non-degenerate, and correspond to the roots of the function $Q(u, \xi)$ that can be expressed as $\xi_0 = (1 + \tau)e^{i\varphi}$, $\xi_1 = e^{-2i\varphi}$, $\xi_2 = (1 + \tau)^{-1}e^{i\varphi}$, $(1 + \tau) = (1 + \omega)^2$.

Lemma 3.1 is proved in Sec. 5.

Formula (3.8) and Lemma 3.1 give a complete description of the stationary points of the function $S(u, \zeta)$.

In order to estimate the large-time behavior of the integral having the form

$$I(t, u, \lambda) = \iint_{\mathbb{C}} f(\zeta, \lambda) \exp(itS(u, \zeta)) d\operatorname{Re} \zeta d\operatorname{Im} \zeta \tag{3.9}$$

uniformly on $u, \lambda \in \mathbb{C}$, in the present and the following sections we will use the following general scheme.

- (1) Consider D_ε , the union of disks with a radius of ε and centers in singular points of function $f(\zeta, \lambda)$ and stationary points of $S(u, \zeta)$.
- (2) Represent $I(t, u, \lambda)$ as the sum of integrals over D_ε and $\mathbb{C} \setminus D_\varepsilon$:

$$\begin{aligned}
 I(t, u, \lambda) &= I_{\text{int}} + I_{\text{ext}}, \quad \text{where} \\
 I_{\text{int}} &= \iint_{D_\varepsilon} f(\zeta, \lambda) \exp(itS(u, \zeta)) d\operatorname{Re} \zeta d\operatorname{Im} \zeta, \\
 I_{\text{ext}} &= \iint_{\mathbb{C} \setminus D_\varepsilon} f(\zeta, \lambda) \exp(itS(u, \zeta)) d\operatorname{Re} \zeta d\operatorname{Im} \zeta.
 \end{aligned} \tag{3.10}$$

- (3) Find an estimate of the form

$$|I_{\text{int}}| = O(\varepsilon^\alpha), \quad \text{as } \varepsilon \rightarrow 0 \quad (\alpha \geq 1)$$

uniformly on u, λ, t .

- (4) Integrate I_{ext} by parts using Stokes formula

$$\begin{aligned}
 I_{\text{ext}} &= \frac{1}{2t} \int_{\partial D_\varepsilon} \frac{f(\zeta, \lambda) \exp(itS(u, \zeta))}{S'_\zeta(u, \zeta)} d\bar{\zeta} \\
 &\quad - \frac{1}{it} \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{f'_\zeta(\zeta, \lambda) \exp(itS(u, \zeta))}{S'_\zeta(u, \zeta)} d\operatorname{Re} \zeta d\operatorname{Im} \zeta \\
 &\quad - \frac{1}{it} \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{f(\zeta, \lambda) \exp(itS(u, \zeta)) S''_{\zeta\zeta}(u, \zeta)}{(S'_\zeta(u, \zeta))^2} d\operatorname{Re} \zeta d\operatorname{Im} \zeta \\
 &= -\frac{1}{t} (I_1 - I_2 - I_3).
 \end{aligned} \tag{3.11}$$

- (5) For each I_i find an estimate of the form

$$(a) |I_i| = O\left(\ln \frac{1}{\varepsilon}\right) \quad \text{or} \quad (b) |I_i| = O\left(\frac{1}{\varepsilon^\beta}\right), \quad \text{as } \varepsilon \rightarrow 0.$$

(6) In case (a) set $\varepsilon = \frac{1}{|t|}$ which yields the overall estimate

$$|I(t, u, \lambda)| = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty.$$

In case (b) set $\varepsilon = \frac{1}{|t|^k}$, where $k(\alpha + \beta) = 1$, which yields the overall estimate

$$|I(t, u, \lambda)| = O\left(\frac{1}{|t|^{\frac{\alpha}{\alpha+\beta}}}\right), \quad \text{as } t \rightarrow \infty.$$

Using this scheme we obtain, in particular, the following result

Lemma 3.2. *Under assumptions (3.2), (3.4),*

$$|I(t, u)| = O\left(\frac{\ln(3 + |t|)}{1 + |t|}\right) \quad \text{for } t \in \mathbb{R}$$

uniformly on $u \in \mathbb{C}$.

A detailed proof of Lemma 3.2 is given in Sec. 5.

4. Estimate for the Nonlinearized Case

In this section we prove estimate (1.14) for the solution $v(x, t)$ of the Cauchy problem for the Novikov–Veselov equation at positive energy with the initial data $v(x, 0)$ satisfying properties (1.10)–(1.13) or, more generally, for $v(x, t)$ constructed by means of (2.6)–(2.14).

We proceed from the formulas (2.13), (2.14) for the potential $v(z, t)$ and the integral equation (2.12) for $\mu(z, \lambda, t)$.

We write (2.12) as

$$\mu(z, \lambda, t) = 1 + (A_{z,t}\mu)(z, \lambda, t), \tag{4.1}$$

where

$$(A_{z,t}f)(\lambda) = \partial_{\bar{\lambda}}^{-1}(r(\lambda) \exp(itS(u, \lambda))\overline{f(\lambda)}) = -\frac{1}{\pi} \iint_{\mathbb{C}} \frac{r(\zeta) \exp(itS(u, \zeta))\overline{f(\zeta)}}{\zeta - \lambda} d\operatorname{Re} \zeta d\operatorname{Im} \zeta$$

and $S(u, \zeta)$ is defined by (3.5), $u = \frac{z}{t}$.

Equation (4.1) can be also written in the form

$$\mu(z, \lambda, t) = 1 + A_{z,t} \cdot 1 + (A_{z,t}^2\mu)(z, \lambda, t). \tag{4.2}$$

According to the theory of the generalized analytic functions (see [12]), Eqs. (4.1) and (4.2) have a unique solution for all z, t . This solution can be written as

$$\mu(z, \lambda, t) = (I - A_{z,t}^2)^{-1}(1 + A_{z,t} \cdot 1). \tag{4.3}$$

Equation (4.3) possesses a formal asymptotic expansion

$$\mu(z, \lambda, t) = (I + A_{z,t}^2 + A_{z,t}^4 + \dots)(1 + A_{z,t} \cdot 1). \tag{4.4}$$

From estimate (4.10) given below it follows that (4.4) uniformly converges for sufficiently large t . We will also write formula (4.4) in the form

$$\mu(z, \lambda, t) = 1 + A_{z,t} \cdot 1 + R, \tag{4.5}$$

where $R = (\sum_{k=1}^{\infty} A_{z,t}^{2k})(1 + A_{z,t} \cdot 1)$.

In addition to $A_{z,t}$ we introduce another integral operator $B_{z,t}$ defined as

$$B_{z,t} \cdot f = \iint_{\mathbb{C}} r(\zeta) \exp(itS(u, \zeta)) \overline{f(\zeta)} d\operatorname{Re} \zeta d\operatorname{Im} \zeta. \tag{4.6}$$

To study (4.4) we will need some estimates on the values of operators $A_{z,t}$ and $B_{z,t}$.

Lemma 4.1. *Under assumptions (2.9)–(2.11), the following estimates hold:*

(a)

$$|B_{z,t} \cdot 1| = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty \tag{4.7}$$

uniformly on $u \in \mathbb{C}$;

(b)

$$|B_{z,t} \cdot A_{z,t} \cdot 1| = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty \tag{4.8}$$

uniformly on $u \in \mathbb{C}$;

(c)

$$|(A_{z,t}^2 \cdot 1)(\lambda)| \leq \frac{\beta}{|t|^{1/14}}, \quad |t| \geq 1, \tag{4.9}$$

where β *is a constant independent of* u *and* λ ;

(d)

$$|(A_{z,t}^n \cdot 1)(\lambda)| \leq \frac{\beta^{n-1}}{|t|^{\lfloor n/2 \rfloor / 14}}, \quad |t| \geq 1; \tag{4.10}$$

(e)

$$|B_{z,t} \cdot A_{z,t}^{n-1} \cdot 1| \leq \frac{\beta^{n-1} \ln(3 + |t|)}{|t|^{1 + \lfloor (n-2)/2 \rfloor / 14}}, \quad |t| \geq 1. \tag{4.11}$$

Here $\lfloor s \rfloor$ denotes the integer part of $s \geq 0$.

Proof of Lemma 4.1. We proceed according to the scheme described in the previous section.

(a) This point follows from Lemma 3.2.

(b) As $(A_{z,t} \cdot 1)(\lambda) \in C(\mathbb{C})$, we take D_ε to be the union of disks of a radius ε centered in the stationary points of $S(u, \zeta)$. We note that $(A_{z,t} \cdot 1)(\lambda) = O(1)$ uniformly on

u (or, equivalently, on z) and λ . Thus the integral I_{int} (as in (3.10) with $f(\zeta, \lambda) = r(\zeta)\overline{(A_{z,t} \cdot 1)(\zeta)}$) can be estimated as

$$I_{\text{int}} = \iint_{D_\varepsilon} r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t} \cdot 1)(\zeta)} d\text{Re } \zeta d\text{Im } \zeta = O(\varepsilon^2).$$

Now let us estimate the integral I_{ext} (as in (3.10) with $f(\zeta, \lambda) = r(\zeta)\overline{(A_{z,t} \cdot 1)(\zeta)}$). For this purpose we apply the Stokes formula (as in (3.11)) taking into consideration that $\partial_{\bar{\lambda}}(A_{z,t} \cdot f)(\lambda) = r(\lambda) \exp(itS(u, \lambda))f(\lambda)$:

$$\begin{aligned} & \iint_{\mathbb{C} \setminus D_\varepsilon} r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t} \cdot 1)(\zeta)} d\text{Re } \zeta d\text{Im } \zeta \\ &= \int_{\partial D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t} \cdot 1)(\zeta)}}{2tS'_\zeta(u, \zeta)} d\bar{\zeta} \\ &\quad - \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{r'_\zeta(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t} \cdot 1)(\zeta)}}{itS'_\zeta(u, \zeta)} d\text{Re } \zeta d\text{Im } \zeta \\ &\quad + \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t} \cdot 1)(\zeta)} S''_{\zeta\zeta}(u, \zeta)}{it(S'_\zeta(u, \zeta))^2} d\text{Re } \zeta d\text{Im } \zeta \\ &\quad - \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{|r(\zeta)|^2 |\exp(itS(u, \zeta))|^2}{itS'_\zeta(u, \zeta)} d\text{Re } \zeta d\text{Im } \zeta. \end{aligned} \tag{4.12}$$

Now, proceeding as in the proof of Lemma 3.2, we obtain (4.8).

- (c) In this case we build D_ε as the union of disks with a radius of ε and centers in λ and stationary points of $S(u, \zeta)$. The integral I_{int} over D_ε (as in (3.10) with $f(\zeta, \lambda) = -\frac{1}{\pi} \frac{r(\zeta)\overline{(A_{z,t} \cdot 1)(\zeta)}}{\zeta - \lambda}$) behaves asymptotically as $O(\varepsilon)$. When estimating the integral I_{ext} over $\mathbb{C} \setminus D_\varepsilon$ we use (3.8), (3.11) and the following inequalities

$$|\zeta - \lambda| \geq \varepsilon, \quad |\zeta - \zeta_i| \geq \varepsilon$$

(ζ_i are stationary points of $S(u, \zeta)$) which hold for all $\zeta \in \mathbb{C} \setminus D_\varepsilon$. Thus we obtain that the asymptotical behavior of I_{ext} is at most $O(\frac{1}{|t|^\varepsilon})$. Then, as proposed by the scheme, we choose $\varepsilon = |t|^{-1/14}$ and obtain the required estimate.

- (d) This point is proved by induction. As in point (c) D_ε is the union of disks with a radius of ε and centers in λ and stationary points of $S(u, \zeta)$.

For the integral I_{int} we have

$$\begin{aligned} |I_{\text{int}}| &= \left| \frac{1}{\pi} \iint_{D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta))}{\zeta - \lambda} \overline{(A_{z,t}^{n-1} \cdot 1)(\zeta)} d\text{Re } \zeta d\text{Im } \zeta \right| \\ &\leq \frac{\beta^{n-2}}{|t|^{\lfloor (n-1)/2 \rfloor / 14}} \iint_{D_\varepsilon} \left| \frac{r(\zeta) \exp(itS(u, \zeta))}{\zeta - \lambda} \right| d\text{Re } \zeta d\text{Im } \zeta \leq \frac{\beta^{n-1} \varepsilon}{|t|^{\lfloor (n-2)/2 \rfloor / 14}}. \end{aligned} \tag{4.13}$$

To estimate I_{ext} we use the following representation

$$\begin{aligned}
 & \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t}^{n-1} \cdot 1)(\zeta)}}{\zeta - \lambda} d \operatorname{Re} \zeta d \operatorname{Im} \zeta \\
 &= \int_{\partial D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t}^{n-1} \cdot 1)(\zeta)}}{2t(\zeta - \lambda) S'_\zeta(u, \zeta)} d\bar{\zeta} \\
 &\quad - \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{r'_\zeta(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t}^{n-1} \cdot 1)(\zeta)}}{it(\zeta - \lambda) S'_\zeta(u, \zeta)} d \operatorname{Re} \zeta d \operatorname{Im} \zeta \\
 &\quad + \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t}^{n-1} \cdot 1)(\zeta)} S''_{\zeta\zeta}(u, \zeta)}{it(\zeta - \lambda) (S'_\zeta(u, \zeta))^2} d \operatorname{Re} \zeta d \operatorname{Im} \zeta \\
 &\quad - \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{|r(\zeta)|^2 |\exp(itS(u, \zeta))|^2 \overline{(A_{z,t}^{n-2} \cdot 1)(\zeta)}}{it(\zeta - \lambda) S'_\zeta(u, \zeta)} d \operatorname{Re} \zeta d \operatorname{Im} \zeta \\
 &\quad + \iint_{\mathbb{C} \setminus D_\varepsilon} \frac{r(\zeta) \exp(itS(u, \zeta)) \overline{(A_{z,t}^{n-1} \cdot 1)(\zeta)}}{it(\zeta - \lambda)^2 S'_\zeta(u, \zeta)} d \operatorname{Re} \zeta d \operatorname{Im} \zeta \\
 &= J_1 + J_2 + J_3 + J_4 + J_5.
 \end{aligned} \tag{4.14}$$

The integrals J_i can be estimated in the following way

$$|J_1| \leq \frac{\beta^{n-2}}{|t| \cdot |t|^{\lfloor (n-1)/2 \rfloor / 14}} \frac{1}{\varepsilon^7} \int_{\partial D_\varepsilon} |r(\zeta)| d\zeta \leq \frac{\beta^{n-1}}{|t| \cdot |t|^{\lfloor (n-1)/2 \rfloor / 14}} \frac{1}{\varepsilon^7}.$$

Similarly,

$$\begin{aligned}
 |J_2| &\leq \frac{\beta^{n-1}}{|t| \cdot |t|^{\lfloor (n-1)/2 \rfloor / 14}} \frac{1}{\varepsilon^7}, \\
 |J_3| &\leq \frac{\beta^{n-1}}{|t| \cdot |t|^{\lfloor (n-1)/2 \rfloor / 14}} \frac{1}{\varepsilon^{13}}, \\
 |J_4| &\leq \frac{\beta^{n-1}}{|t| \cdot |t|^{\lfloor (n-2)/2 \rfloor / 14}} \frac{1}{\varepsilon^7}, \\
 |J_5| &\leq \frac{\beta^{n-1}}{|t| \cdot |t|^{\lfloor (n-2)/2 \rfloor / 14}} \frac{1}{\varepsilon^8}.
 \end{aligned}$$

Thus,

$$|I_{\text{ext}}| \leq \frac{\beta^{n-1}}{|t| \cdot |t|^{\lfloor (n-2)/2 \rfloor / 14} \cdot \varepsilon^{13}}.$$

Now we set $\varepsilon = |t|^{-1/14}$ and obtain the overall estimate

$$|(A_{z,t}^n \cdot 1)(\lambda)| \leq \frac{\beta^{n-1}}{|t|^{1/14} |t|^{\lfloor (n-2)/2 \rfloor / 14}} = \frac{\beta^{n-1}}{t^{\lfloor n/2 \rfloor / 14}}.$$

(e) Proceeding from (d) this point is proved similarly to (b). □

Lemma 4.2. *Under the assumptions of Lemma 4.1, we have that:*

(a) $A_{z,t} \cdot 1 = \frac{a_1(z,t)}{\lambda} + O\left(\frac{1}{|\lambda|^2}\right)$ for $\lambda \rightarrow \infty$, where

$$|\partial_z a_1(z,t)| = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty \tag{4.15}$$

uniformly on $z \in \mathbb{C}$.

(b) $A_{z,t}^2 \cdot 1 = \frac{a_2(z,t)}{\lambda} + O\left(\frac{1}{|\lambda|^2}\right)$ for $\lambda \rightarrow \infty$, where

$$|\partial_z a_2(z,t)| = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty \tag{4.16}$$

uniformly on $z \in \mathbb{C}$.

(c) $R = \frac{q(z,t)}{\lambda} + O\left(\frac{1}{|\lambda|^2}\right)$ as $\lambda \rightarrow \infty$, where

$$|\partial_z q(z,t)| = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty \tag{4.17}$$

uniformly on $z \in \mathbb{C}$, where R is defined in (4.5).

Proof of Lemma 4.2. The asymptotics for $A_{z,t} \cdot 1$, $A_{z,t}^2 \cdot 1$ and R follow from the definitions of $A_{z,t}$ and R , formula (2.13) and properties (2.9), (2.10). The rest of the proof consists in the following.

(a) Estimate (4.15) follows from (4.7) and the formula

$$a_1(z,t) = \frac{1}{\pi} \iint_{\mathbb{C}} r(\zeta) \exp(itS(u,\zeta)) d\text{Re } \zeta d\text{Im } \zeta = B_{z,t} \cdot 1.$$

(b) Estimate (4.16) follows from (4.8) and the formula

$$a_2(z,t) = \frac{1}{\pi} \iint_{\mathbb{C}} r(\zeta) \exp(itS(u,\zeta)) \overline{(A_{z,t} \cdot 1)(\zeta)} d\text{Re } \zeta d\text{Im } \zeta = B_{z,t} \cdot A_{z,t} \cdot 1.$$

(c) We note that $q(z,t) = \sum_{k=2}^{\infty} a_k(z,t)$ where $a_k(z,t)$ is defined

$$(A_{z,t}^k \cdot 1)(\lambda) = \frac{a_k(z,t)}{\lambda} + O\left(\frac{1}{|\lambda|^2}\right).$$

Next,

$$a_k(z,t) = \frac{1}{\pi} \iint_{\mathbb{C}} r(\zeta) \exp(itS(u,\zeta)) \overline{(A_{z,t}^{k-1} \cdot 1)(\zeta)} d\text{Re } \zeta d\text{Im } \zeta.$$

Thus, proceeding as in point (e) of Lemma 4.1, we obtain that $a_2(z,t) + a_3(z,t) = O\left(\frac{\ln(|t|)}{|t|}\right)$ and the rest of the members form a geometric progression that converges to the sum of order $O\left(\frac{\ln(|t|)}{|t|^{1+1/14}}\right)$. □

Formulas (2.13), (2.14), (4.5) and Lemma 4.2 imply

Theorem 4.1. *Let $v(x, t)$ be a solution to the Cauchy problem for the Novikov–Veselov equation (1.6) with $E = 1$, constructed via (2.12)–(2.14) under assumptions (2.6)–(2.11). Then*

$$|v(x, t)| \leq \frac{\text{const}(v) \ln(3 + |t|)}{1 + |t|}, \quad x \in \mathbb{R}^2, \quad t \in \mathbb{R}.$$

5. Proofs of Lemmas 3.1 and 3.2

Proof of Lemma 3.1. Under the additional assumption that $\xi \neq 0$ the system of Eqs. (3.6) and (3.7) is equivalent to the following system

$$\begin{cases} \xi^3 - \frac{\bar{u}}{6}\xi^2 + \frac{u}{6}\xi - 1 = 0, \\ \xi^3 - \frac{u}{6}\xi + 2 = 0. \end{cases} \tag{5.1}$$

We claim that $\xi = \xi_1$ corresponds to a degenerate stationary point of (3.5), if

$$\begin{aligned} &\text{the polynomial } P(\xi) = \xi^3 - \frac{\bar{u}}{6}\xi^2 + \frac{u}{6}\xi - 1 \text{ can be represented} \\ &\text{in the form } P(\xi) = (\xi - \xi_1)^2(\xi - \xi_2). \end{aligned} \tag{5.2}$$

Indeed, if $\xi = \xi_1 \neq 0$ is a zero of the function $Q(u, \xi) = -\frac{\bar{u}}{2} + \frac{u}{2\xi} + 3\xi - \frac{3}{\xi^2} = 3P(\xi)/\xi^2$, then $Q(u, \xi)$ is holomorphic in a certain neighborhood of $\xi = \xi_1$ and can be expanded into the following Taylor series

$$Q(u, \xi) = c_1(\xi - \xi_1) + c_2(\xi - \xi_1)^2 + c_3(\xi - \xi_1)^3 + \dots$$

Thus S'_ζ can be represented as

$$S'_\zeta(u, \zeta) = c_1(\zeta^2 - \xi_1) + c_2(\zeta^2 - \xi_1)^2 + c_3(\zeta^2 - \xi_1)^3 + \dots$$

After differentiating with respect to ζ we obtain

$$S''_{\zeta\zeta}(u, \zeta) = 2c_1\zeta + 4c_2(\zeta^2 - \xi_1)\zeta + 6c_3(\zeta^2 - \xi_1)^2\zeta + \dots$$

The stationary point corresponding to $\xi = \xi_1$ can be degenerate if and only if $c_1 = 0$.

So for the polynomial $P(\xi)$ we get the following representation in the neighborhood of $\xi = \xi_1$

$$P(\xi) = c_2(\xi - \xi_1)^2\xi^2 + O(|\xi - \xi_1|^3).$$

As $P(\xi)$ is a third-order polynomial, it follows that $P(\xi) = (\xi - \xi_1)^2(\xi - \xi_2)$.

Expanding the expression for $P(\xi)$ and equating coefficients for the corresponding powers of ξ results in the following system

$$\begin{cases} 2\xi_1 + \xi_2 = \frac{\bar{u}}{6}, \\ 2\xi_1\xi_2 + \xi_1^2 = \frac{u}{6}, \\ \xi_1^2\xi_2 = 1. \end{cases} \tag{5.3}$$

Excluding u from the system yields

$$\begin{cases} 2\xi_1\xi_2 + \xi_1^2 = 2\bar{\xi}_1 + \bar{\xi}_2, \\ \xi_1^2\xi_2 = 1. \end{cases} \tag{5.4}$$

We represent ξ_1 in the form $\xi_1 = re^{i\varphi}$. Then from the second equation in (5.4) we get that $\xi_2 = r^{-2}e^{-2i\varphi}$. Substituting this into the first equation of (5.4) yields

$$(r^2 - r^{-2})e^{2i\varphi} = (r - r^{-1})e^{-i\varphi}. \tag{5.5}$$

For $r = 1$ Eq. (5.5) holds for all φ . In addition, if $r \neq 1, r > 0$, then Eq. (5.5) can be rewritten as

$$(r + r^{-1})e^{3i\varphi} = 2$$

and has no solutions for real φ .

Using now the second equation in (5.3) one can see that the set of u values, for which (5.4), (5.3) are solvable and (5.2) holds, is a curve on the complex plane described in the parametric form by

$$u = 6(2e^{-i\varphi} + e^{2i\varphi})$$

(see Fig. 1). This curve has three singular points corresponding to $\varphi = \frac{2\pi k}{3}, k = 0, 1, 2$. For these values of φ we have that $e^{i\varphi} = \xi_1 = \xi_2 = e^{-2i\varphi}$, and $P(\xi)$ can be represented in the




Fig. 1. The curve \mathcal{U} on the complex plane.

form $P(\xi) = (\xi - \xi_1)^3$. For $\varphi \neq \frac{2\pi k}{3}, k = 0, 1, 2$, we have that $P(\xi) = (\xi - \xi_1)^2(\xi - \xi_2)$ where $e^{i\varphi} = \xi_1 \neq \xi_2 = e^{-2i\varphi}$. The first two statements of the Lemma 3.1 are proved.

Let us now fix $\xi = e^{i\varphi}$ and find the set of u for which this ξ is a root of the polynomial $P(\xi)$.

One can see that ξ is the root of $P(\xi)$ if

$$\xi \frac{u}{6} - \xi^2 \frac{\bar{u}}{6} = 1 - \xi^3. \tag{5.6}$$

We now solve the homogeneous equation

$$e^{i\varphi} \frac{u}{6} - e^{2i\varphi} \frac{\bar{u}}{6} = 0 \tag{5.7}$$

with respect to u to find the plausible perturbations of u for which $\xi = e^{i\varphi}$ remains a root of $P(\xi)$.

From (5.7) we get $\frac{u}{\bar{u}} = e^{i\varphi}$, and thus $u = se^{i\varphi/2}$ for $s \in \mathbb{R}$. So for all u that belong to the line

$$u(\varphi, s) = 6(2e^{-i\varphi} + e^{2i\varphi}) + se^{i\varphi/2}, \quad s \in \mathbb{R} \tag{5.8}$$

one of the roots of $P(\xi)$ is equal to $e^{i\varphi}$.

Now we note that the tangent vector to \mathcal{U}

$$\frac{d}{d\varphi}(2e^{-i\varphi} + e^{2i\varphi}) = 2i(e^{2i\varphi} - e^{-i\varphi})$$

is collinear to the perturbation vector $se^{i\varphi/2}$ for all $\varphi \neq \frac{2\pi k}{3}, k = 0, 1, 2$. Thus $u(\varphi, s)$ given by (5.8) is the tangent line to \mathcal{U} passing through the point $6(2e^{-i\varphi} + e^{2i\varphi})$.

We note that for each $u \in \text{int } \mathbb{U}$ there exist two different tangents to the curve \mathcal{U} passing through this u . Indeed, note that the tangent lines to the curve \mathcal{U} passing through the points $u = 18e^{\frac{2\pi ik}{3}}, k = 0, 1, 2$, divide the domain $\text{int } \mathbb{U}$ into three parts

$$\begin{aligned} U_0 &= \left\{ u = re^{i\varphi}, 0 \leq r < 6(2e^{-i\varphi} + e^{2i\varphi}), \frac{-\pi}{3} \leq \varphi \leq \frac{\pi}{3} \right\}, \\ U_1 &= \left\{ u = re^{i\varphi}, 0 < r < 6(2e^{-i\varphi} + e^{2i\varphi}), \frac{\pi}{3} < \varphi \leq \pi \right\}, \\ U_2 &= \left\{ u = re^{i\varphi}, 0 < r < 6(2e^{-i\varphi} + e^{2i\varphi}), \pi < \varphi < \frac{5\pi}{3} \right\} \end{aligned}$$

(see Fig. 1). We first study U_0 . Let us consider K_{01} and K_{02} which are the sets of points of the following pencils of tangent lines:

$$\begin{aligned} K_{01} &= \left\{ u \in \mathbb{C} : u = 6(2e^{-i\varphi} + e^{2i\varphi}) + se^{i\varphi/2}, s \in \mathbb{R}, -\frac{2\pi}{3} \leq \varphi < 0 \right\}, \\ K_{02} &= \left\{ u \in \mathbb{C} : u = 6(2e^{-i\varphi} + e^{2i\varphi}) + se^{i\varphi/2}, s \in \mathbb{R}, 0 < \varphi \leq \frac{2\pi}{3} \right\}. \end{aligned}$$

It is easily seen that $U_0 \subset K_{01}, U_0 \subset K_{02}$, i.e. each point from U_0 is covered by a certain tangent line from both pencils K_{01} and K_{02} . It can be shown similarly that each point of $U_j, j = 1, 2$, is covered twice by the corresponding tangent lines.

Thus every point from $\text{int } \mathbb{U}$ is covered twice which means that for each u in the domain limited by the curve \mathcal{U} there exist two different roots of $P(\xi)$ equal to 1 in absolute value. As the product of the roots of the polynomial $P(\xi)$ is equal to 1, the third root of the polynomial is also equal to 1 in absolute value. We do not consider the values u from the boundary of \mathbb{U} which means that the above-mentioned roots correspond to non-degenerate stationary points. Thus the third statement of Lemma 3.1 is proved.

Now suppose $u \in \mathbb{C} \setminus \mathbb{U}$. We note that every such point u belongs to one and only one tangent line to the curve \mathcal{U} . That means that for every $u \in \mathbb{C} \setminus \mathbb{U}$ one of the roots of the polynomial $P(\xi), \xi_1$, is such that $|\xi_1| = 1$. As $u \notin \mathcal{U}, \xi_0 \neq \xi_1, \xi_2 \neq \xi_1$. Besides $|\xi_0| \neq 1$ because otherwise there would be two different tangent lines passing through the corresponding point u . That means that for every $u \in \mathbb{C} \setminus \mathbb{U}$ there exists a root of the polynomial $P(\xi)$, namely $\xi = \xi_0$, such that $|\xi_0| \neq 1$.

Considering Eq. (5.6) and its conjugate yields the following system of linear equations for u and \bar{u}

$$\begin{cases} au + b\bar{u} = c, \\ \bar{b}\bar{u} + a\bar{a}\bar{u} = \bar{c}, \end{cases}$$

where $a = \frac{\xi}{6}, b = -\frac{\xi^2}{6}, c = 1 - \xi^3$ for each $\xi \in \mathbb{C}, |\xi| \neq 1, \xi \neq 0$. Thus

$$u = \frac{c\bar{a} - \bar{c}b}{a\bar{a} - b\bar{b}} = 6 \frac{\bar{\xi} - \bar{\xi}\xi^3 + \xi^2 - \xi^2\bar{\xi}^3}{\xi\bar{\xi}(1 - \xi\bar{\xi})} = 6 \left(\frac{1}{\xi} + \bar{\xi} + \frac{\xi}{\bar{\xi}} \right)$$

for each $\xi \in \mathbb{C}, |\xi| \neq 1, \xi \neq 0$.

Now let us consider $\xi = \xi_0 = (1 + \tau)e^{i\varphi}, 0 < \tau < +\infty$. Then the corresponding value of the parameter u is

$$u = 6 \left(2e^{-i\varphi} + e^{2i\varphi} + \frac{\tau^2}{1 + \tau} e^{-i\varphi} \right).$$

In addition to $\xi = \xi_0$, for this value of the parameter u polynomial $P(\xi)$ has also a root $\xi = \xi_1 = e^{-2i\varphi}$. Indeed, if $u = 6(2e^{-i\varphi} + e^{2i\varphi})$, then $\xi_1 = e^{-2i\varphi}$ is a root of the polynomial $P(\xi)$, and the plausible perturbation of this u for which ξ_1 remains a root of $P(\xi)$ is equal to $se^{-i\varphi}$.

In addition, as the product of the roots of P is equal to 1, the third root is $\xi_2 = (1 + \tau)^{-1}e^{i\varphi}$. The fourth statement of Lemma 3.1 is proved. □

Proof of Lemma 3.2. In this case D_ε is the union of disks with a radius of ε centered in the stationary points of $S(u, \zeta)$. The integral I_{int} (as in (3.10) with $f(\zeta, \lambda) = f(\zeta)$ satisfying properties (3.2)) is estimated as

$$|I_{\text{int}}| = \left| \iint_{D_\varepsilon} f(\zeta) \exp(itS(u, \zeta)) d\text{Re } \zeta d\text{Im } \zeta \right| \leq \text{const} \left| \iint_{D_\varepsilon} d\text{Re } \zeta d\text{Im } \zeta \right| = O(\varepsilon^2).$$

The estimate for I_{ext} (as in (3.10) with $f(\zeta, \lambda) = f(\zeta)$ satisfying properties (3.2)) is proved separately for $u \in \mathbb{U}$ and $u \in \mathbb{C} \setminus \mathbb{U}$.

(I) $u \in \mathbb{U}$:

In this case all $\zeta \in D_\varepsilon$ lie in the ε -neighborhood of the unit circle. Consequently, from (2.6), (2.9) and (2.11) it follows that for any $N \in \mathbb{N}$ and any $\varepsilon_0 \in (0, 1/2]$ there exists $C = C(\varepsilon_0, N)$ such that

$$|f(\zeta)| \leq C\rho^N, \quad |f'_\zeta(\zeta)| \leq C\rho^N$$

for all $\zeta \in D_\rho$, $\rho \leq \varepsilon_0$.

The function $S'_\zeta(u, \zeta)$ can be estimated as

$$|S'_\zeta(u, \zeta)| \geq 3\frac{\varepsilon_0^6}{|\zeta|^4} \quad \text{for } \zeta \in \mathbb{C} \setminus D_{\varepsilon_0}, \quad \text{and}$$

$$|S'_\zeta(u, \zeta)| \geq 3\frac{\rho^6}{|\zeta|^4} \quad \text{for } \zeta \in \partial D_\rho, \quad \varepsilon \leq \rho \leq \varepsilon_0.$$

Taking $N = 5$ results in the following estimate for I_1

$$\begin{aligned} |I_1| &\leq \frac{1}{2} \int_{\partial D_\varepsilon} \frac{|f(\zeta)|}{|S'_\zeta(u, \zeta)|} d\bar{\zeta} \leq \text{const} \frac{\varepsilon^N}{\varepsilon^6} \int_{\partial D_\varepsilon} |\zeta|^4 d\bar{\zeta} \\ &\leq \text{const} \frac{\varepsilon^N \varepsilon}{\varepsilon^6} (1 + \varepsilon)^4 = O(1), \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

When estimating I_2 and I_3 we integrate separately over $D_{\varepsilon_0} \setminus D_\varepsilon$ and $\mathbb{C} \setminus D_{\varepsilon_0}$:

$$\begin{aligned} |I_2| &\leq \iint_{D_{\varepsilon_0} \setminus D_\varepsilon} \left| \frac{f'_\zeta(\zeta) \exp(itS(u, \zeta))}{S'_\zeta(u, \zeta)} \right| d\text{Re } \zeta d\text{Im } \zeta \\ &\quad + \iint_{\mathbb{C} \setminus D_{\varepsilon_0}} \left| \frac{f'_\zeta(\zeta) \exp(itS(u, \zeta))}{S'_\zeta(u, \zeta)} \right| d\text{Re } \zeta d\text{Im } \zeta \\ &\leq \text{const} \int_\varepsilon^{\varepsilon_0} \frac{\rho^N \rho}{\rho^6} d\rho + \text{const} \iint_{\mathbb{C} \setminus D_{\varepsilon_0}} |f(\zeta)| |\zeta^4| d\text{Re } \zeta d\text{Im } \zeta = O(1), \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

$$\begin{aligned} |I_3| &\leq \iint_{D_{\varepsilon_0} \setminus D_\varepsilon} \left| \frac{f(\zeta) \exp(itS(u, \zeta)) S''_{\zeta\zeta}(u, \zeta)}{(S'_\zeta(u, \zeta))^2} \right| d\text{Re } \zeta d\text{Im } \zeta \\ &\quad + \iint_{\mathbb{C} \setminus D_{\varepsilon_0}} \left| \frac{f(\zeta) \exp(itS(u, \zeta)) S''_{\zeta\zeta}(u, \zeta)}{(S'_\zeta(u, \zeta))^2} \right| d\text{Re } \zeta d\text{Im } \zeta \\ &\leq \text{const} \int_\varepsilon^{\varepsilon_0} \frac{\rho^N \rho}{\rho^{12}} d\rho + \text{const} \iint_{\mathbb{C} \setminus D_{\varepsilon_0}} |f(\zeta)| |\zeta^3| d\text{Re } \zeta d\text{Im } \zeta \stackrel{N \equiv 11}{=} O(1), \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Setting finally $\varepsilon = \frac{1}{|t|}$ yields

$$I(t, u) = O\left(\frac{1}{|t|}\right), \quad \text{as } t \rightarrow \infty$$

uniformly on $u \in \mathbb{U}$.

(II) $u \in \mathbb{C} \setminus \mathbb{U}$:

Let us divide the complex plane into six sets, each containing one and only one stationary point of $S(u, \zeta)$: $\mathbb{C} = \bigcup_{k=0}^2 (Z_k^+ \cup Z_k^-)$. We define the set Z_k^\pm as the set of points of the complex plane to which the stationary point $\pm\zeta_k$ is the closest:

$$\begin{aligned} Z_k^+ &= \{\zeta \in \mathbb{C} : |\zeta - \zeta_i| \geq |\zeta - \zeta_k|, |\zeta + \zeta_j| \geq |\zeta - \zeta_k|, i, j \in \{0, 1, 2\}\}, \\ Z_k^- &= \{\zeta \in \mathbb{C} : |\zeta - \zeta_i| \geq |\zeta + \zeta_k|, |\zeta + \zeta_j| \geq |\zeta + \zeta_k|, i, j \in \{0, 1, 2\}\}, \end{aligned}$$

where $k \in \{0, 1, 2\}$. We will estimate the integral over each Z_k^+ separately. The integrals over Z_k^- are treated similarly.

Let us first take $\zeta \in Z_1^+$. Using the definition of Z_1^+ and the property that all $\zeta \in D_\varepsilon \cap Z_1^+$ lie in the ε -neighborhood of the unit circle one can see that the scheme of reasoning for the case I is applicable.

Now let us consider $\zeta \in Z_0^+ \cup Z_2^+$.

(A) First, we will study the set of values of parameter ω for which $\zeta_2(u), -\zeta_2(u)$ lie outside the $2\varepsilon_0$ -neighborhood of zero, i.e. $\frac{1}{1+\omega} > 2\varepsilon_0$.

We will consider $\zeta \in Z_0^+$ (the case $\zeta \in Z_2^+$ is treated similarly). If $\zeta \in Z_0^+ \cap D_{\varepsilon_0}$ (for a certain ε_0), then it can be represented as

$$\zeta = (1 + \omega)e^{i\varphi/2} + \rho e^{i\theta}, \quad \rho \leq \varepsilon_0.$$

Let us estimate the ratio $\frac{f(\zeta)}{S'_\zeta(\zeta, u)}$ in D_{ε_0} . If ζ_0 belongs to the $2\varepsilon_0$ -neighborhood of ζ_1 , then all $\zeta \in Z_0^+ \cap D_{\varepsilon_0}$ belong to the $3\varepsilon_0$ -neighborhood of ζ_1 . The following estimates hold:

$$|\zeta + \zeta_1| \geq 2 - 3\varepsilon_0, \quad |\zeta + \zeta_0| > 2 - \varepsilon_0, \quad |\zeta + \zeta_2| \geq 1 + \varepsilon_0, \quad \zeta \in Z_0^+ \cap D_{\varepsilon_0}. \quad (5.9)$$

Further, we note that for any $N \in \mathbb{N}$ there exists a function $\tilde{f}(\zeta)$ satisfying properties (3.2), such that $|f(\zeta)| \leq |\zeta - \zeta_1|^N |\tilde{f}(\zeta)|$ for ζ belonging to the $3\varepsilon_0$ -neighborhood of ζ_1 . This and (5.9) imply that

$$\left| \frac{f(\zeta)}{S'_\zeta(\zeta, u)} \right| \leq \text{const} \frac{|\tilde{f}(\zeta)| |\zeta|^4}{|\zeta - \zeta_2| |\zeta - \zeta_0|}. \quad (5.10)$$

A similar reasoning holds for the case when ζ_0 belongs to the $2\varepsilon_0$ -neighborhood of $-\zeta_1$. Now if ζ_0 does not belong to the $2\varepsilon_0$ -neighborhood of ζ_1 and $-\zeta_1$, then $|\zeta - \zeta_1| \geq \varepsilon_0$ and $|\zeta + \zeta_1| \geq \varepsilon_0$ for all $\zeta \in Z_0^+ \cap D_{\varepsilon_0}$. Two last estimates of (5.9) hold and thus (5.10) holds with $\tilde{f}(\zeta) = f(\zeta)$.

The difference $|\zeta - \zeta_2|$ can be estimated

$$|\zeta - \zeta_2| \geq \frac{1}{2} |\zeta_0 - \zeta_2| = \frac{\omega(2 + \omega)}{2(1 + \omega)}, \quad \zeta \in Z_0^+. \quad (5.11)$$

In order to get rid of this member in the denominator, let us represent $\tilde{f}(\zeta)$ by the Taylor formula in the neighborhood of $(1 + \omega)e^{i\varphi/2}$:

$$\tilde{f}(\zeta) = \tilde{f}(e^{i\varphi/2} + \omega e^{i\varphi/2}) + \tilde{f}'(e^{i\varphi/2} + \omega e^{i\varphi/2} + s e^{i\theta})\rho, \quad s = \lambda\rho \text{ for some } \lambda \in [0, 1],$$

where ' denotes the derivative with respect to s and where λ depends, in particular, on ρ .

For an arbitrary value of ω the following estimates hold:

$$|\tilde{f}(e^{i\varphi/2} + \omega e^{i\varphi/2})| \leq \text{const}|\omega|^N, \tag{5.12}$$

$$|\tilde{f}'(e^{i\varphi/2} + \omega e^{i\varphi/2})| \leq \text{const}|\omega|^N. \tag{5.13}$$

This finally yields

$$\begin{aligned} \frac{|\tilde{f}(\zeta)||\zeta|^4}{|\zeta - \zeta_2||\zeta - \zeta_0|} &\leq \text{const} \left(\frac{|\tilde{f}(e^{i\varphi/2} + \omega e^{i\varphi/2})|}{\frac{\omega(2+\omega)}{2(1+\omega)}|\zeta - \zeta_0|} + \frac{|\tilde{f}'(e^{i\varphi/2} + \omega e^{i\varphi/2} + \lambda\rho e^{i\theta})|\rho}{\rho|\zeta - \zeta_0|} \right) \\ &\leq \frac{\text{const}}{|\zeta - \zeta_0|}, \quad \zeta \in Z_0^+ \cap D_{\varepsilon_0}. \end{aligned}$$

Now we are ready to estimate I_1 :

$$\int_{Z_0^+ \cap \partial D_\varepsilon} \left| \frac{f(\zeta) \exp(itS(u, \zeta))}{S'_\zeta(u, \zeta)} \right| d\bar{\zeta} \leq \text{const} \int_{Z_0^+ \cap \partial D_\varepsilon} \frac{d\bar{\zeta}}{|\zeta - \zeta_0|} = O(1) \quad \text{as } \varepsilon \rightarrow 0.$$

For I_2 we note that the estimate

$$\left| \frac{f'_\zeta(\zeta)}{S'_\zeta(u, \zeta)} \right| \leq \frac{\text{const}}{|\zeta - \zeta_0|}, \quad \zeta \in Z_0^+ \cap D_{\varepsilon_0},$$

can be obtained using the same reasoning as for the ratio $\frac{f(\zeta)}{S'_\zeta(u, \zeta)}$. Thus

$$\begin{aligned} &\iint_{Z_0^+ \setminus D_\varepsilon} \left| \frac{f'_\zeta(\zeta) \exp(itS(u, \zeta))}{S'_\zeta(u, \zeta)} \right| d\text{Re } \zeta d\text{Im } \zeta \\ &\leq \text{const} \left(\iint_{Z_0^+ \setminus D_{\varepsilon_0}} |f'_\zeta(\zeta)||\zeta|^4 d\text{Re } \zeta d\text{Im } \zeta + \iint_{Z_0^+ \cap (D_{\varepsilon_0} \setminus D_\varepsilon)} \frac{d\text{Re } \zeta d\text{Im } \zeta}{|\zeta - \zeta_0|} \right) \\ &= O(1) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

In order to estimate $\frac{f(\zeta)}{(S'_\zeta(u, \zeta))^2}$ in D_{ε_0} we take the members in the Taylor formula for $\tilde{f}(\zeta)$ up to the second order:

$$\tilde{f}(\zeta) = \tilde{f}(e^{i\varphi/2} + \omega e^{i\varphi/2}) + \tilde{f}'(e^{i\varphi/2} + \omega e^{i\varphi/2})\rho + \frac{1}{2}\tilde{f}''(e^{i\varphi/2} + \omega e^{i\varphi/2} + \lambda\rho e^{i\theta})\rho^2. \tag{5.14}$$

From formulas (5.10)–(5.14) it follows that

$$\begin{aligned} \left| \frac{f(\zeta)}{(S'_\zeta(u, \zeta))^2} \right| &\leq \frac{|\tilde{f}(\zeta)||\zeta|^8}{|\zeta - \zeta_2|^2|\zeta - \zeta_0|^2} \\ &\leq \text{const} \left(\frac{|\tilde{f}(e^{i\varphi/2} + \omega e^{i\varphi/2})|}{\left(\frac{\omega(2+\omega)}{2(1+\omega)}\right)^2|\zeta - \zeta_0|^2} + \frac{|\tilde{f}'(e^{i\varphi/2} + \omega e^{i\varphi/2})|\rho}{\left(\frac{\omega(2+\omega)}{2(1+\omega)}\right)^2|\zeta - \zeta_0|^2} \right. \\ &\quad \left. + \frac{|\tilde{f}''(e^{i\varphi/2} + \omega e^{i\varphi/2} + \lambda\rho e^{i\theta})|\rho^2}{2\rho^2|\zeta - \zeta_0|^2} \right) \\ &\leq \frac{\text{const}}{|\zeta - \zeta_0|^2}, \quad \zeta \in Z_0^+ \cap D_{\varepsilon_0}. \end{aligned}$$

Thus

$$\begin{aligned} &\iint_{Z_0^+ \cap (D_{\varepsilon_0} \setminus D_\varepsilon)} \left| \frac{f(\zeta) \exp(itS(u, \zeta)) S''_{\zeta\zeta}(u, \zeta)}{(S'_\zeta(u, \zeta))^2} \right| d\text{Re } \zeta d\text{Im } \zeta \\ &\leq \text{const} \int_\varepsilon^{\varepsilon_0} \frac{d\rho}{\rho} = \text{const} \ln \frac{\varepsilon_0}{\varepsilon}. \end{aligned}$$

Setting $\varepsilon = \frac{1}{|t|}$ yields $I_3 = O(\ln(|t|))$, as $t \rightarrow \infty$.

(B) Now let $\frac{1}{1+\omega} \leq 2\varepsilon_0$.

If $\zeta \in Z_0^+ \cap D_{\varepsilon_0}$, then the following estimates hold

$$|\zeta \pm \zeta_1| \geq \frac{1}{2\varepsilon_0} - 1 - \varepsilon_0, \quad |\zeta \pm \zeta_2| \geq \frac{1}{2\varepsilon_0} - 3\varepsilon_0, \quad |\zeta + \zeta_0| > 2 - \varepsilon_0.$$

Consequently,

$$|S'_\zeta(u, \zeta)| \geq \frac{\text{const}}{|\zeta|^4} |\zeta - \zeta_0|, \quad \zeta \in Z_0^+ \cap D_{\varepsilon_0}.$$

and the part of the integral I_{ext} over Z_0^+ for this case can be estimated, proceeding as in the previous section, as $O(\frac{\ln(|t|)}{|t|})$, $t \rightarrow \infty$.

If $\zeta \in Z_2^+ \cap D_{\varepsilon_0}$, then the following estimates hold

$$|\zeta \pm \zeta_1| \geq 1 - 3\varepsilon_0, \quad |\zeta \pm \zeta_0| \geq 1 - 3\varepsilon_0,$$

and thus

$$|S'_\zeta(u, \zeta)| \geq \frac{\text{const}}{|\zeta|^4} |\zeta - \zeta_2| |\zeta + \zeta_2|.$$

We can estimate $|\zeta + \zeta_2| \geq |\zeta_2| = \frac{1}{1+\omega}$. Now let us expand $f(\zeta)$ into Taylor formula in the neighborhood of $\frac{1}{1+\omega} e^{i\varphi/2}$:

$$f(\zeta) = f\left(\frac{1}{1+\omega} e^{i\varphi/2}\right) + f'\left(\frac{1}{1+\omega} e^{i\varphi/2} + \lambda\rho e^{i\theta}\right)\rho, \quad \lambda \in [0, 1].$$

For an arbitrary value of $\omega > 0$ (satisfying $\frac{1}{1+\omega} \leq 2\varepsilon_0$) the following estimate holds:

$$\left| f\left(\frac{1}{1+\omega}e^{i\varphi/2}\right) \right| \leq \text{const} \left| \frac{1}{1+\omega} \right|^N.$$

This yields

$$\left| \frac{f(\zeta)}{S'_\zeta(u, \zeta)} \right| \leq \text{const} \left(\frac{|f(\frac{1}{1+\omega}e^{i\varphi/2})|}{\frac{1}{1+\omega}|\zeta - \zeta_2|} + \frac{|f'(\frac{1}{1+\omega}e^{i\varphi/2} + \lambda\rho e^{i\theta})|\rho}{\rho|\zeta - \zeta_2|} \right) \leq \frac{\text{const}}{|\zeta - \zeta_2|}.$$

In the same manner

$$\begin{aligned} \left| \frac{f(\zeta)}{(S'_\zeta(u, \zeta))^2} \right| &\leq \frac{|f(\zeta)||\zeta|^8}{|\zeta + \zeta_2|^2|\zeta - \zeta_2|^2} \leq \text{const} \left(\frac{|f(\frac{1}{1+\omega}e^{i\varphi/2})|}{\left(\frac{1}{1+\omega}\right)^2|\zeta - \zeta_2|^2} \right. \\ &\quad \left. + \frac{|f'(\frac{1}{1+\omega}e^{i\varphi/2})|\rho}{\left(\frac{1}{1+\omega}\right)^2|\zeta - \zeta_2|^2} + \frac{|f''(\frac{1}{1+\omega}e^{i\varphi/2} + \lambda\rho e^{i\theta})|\rho^2}{2\rho^2|\zeta - \zeta_2|^2} \right) \leq \frac{\text{const}}{|\zeta - \zeta_2|^2}. \end{aligned}$$

Following further the reasoning from the case (A) we obtain that

$$I_{\text{ext}} = O\left(\frac{\ln(|t|)}{|t|}\right), \quad \text{as } t \rightarrow \infty$$

uniformly on $u \in \mathbb{C}$. □

References

- [1] M. V. Fedoryuk, *Asymptotics: Integrals and Series*, Mathematical Reference Library (Nauka, Moscow, 1987) [in Russian].
- [2] P. G. Grinevich, Scattering transformation at fixed non-zero energy for the two-dimensional Schrödinger operator with potential decaying at infinity, *Russian Math. Surveys* **55**(6) (2000) 1015–1083.
- [3] P. G. Grinevich and S. V. Manakov, The inverse scattering problem for the two-dimensional Schrödinger operator, the $\bar{\partial}$ -method and non-linear equations, *Funkt. Anal. i Pril.* **20**(2) (1986) 14–24 [translation in *Funkt. Anal. Appl.* **20** (1986) 94–103].
- [4] P. G. Grinevich and R. G. Novikov, Transparent potentials at fixed energy in dimension two. Fixed energy dispersion relations for the fast decaying potentials, *Commun. Math. Phys.* **174** (1995) 409–446.
- [5] N. Hayashi, P. I. Naumkin and J.-C. Saut, Asymptotics for large time of global solutions to the generalized Kadomtsev–Petviashvili equation, *Commun. Math. Phys.* **201**(3) (1999) 577–590.
- [6] O. M. Kiselev, Asymptotics of a solution of the Kadomtsev–Petviashvili-2 equation, *Tr. Inst. Mat. Mekh.* **7**(1) (2001) 105–134 [translation in *Proc. Inst. Math. Mech.* **suppl.1** (2001) S107–S139].
- [7] S. V. Manakov, The inverse scattering method and two-dimensional evolution equations, *Uspekhi Mat. Nauk.* **31**(5) (1976) 245–246 [in Russian].
- [8] S. V. Manakov, P. M. Santini and L. A. Takhtadzhyan, An asymptotic behavior of the solutions of the Kadomtsev–Petviashvili equations, *Phys. Lett. A.* **75** (1980) 451–454.
- [9] R. G. Novikov, Reconstruction of a two-dimensional Schrödinger operator from the scattering amplitude at fixed energy, *Funkt. Anal. i Pril.* **20**(3) (1986) 90–91 [translation in *Funkt. Anal. Appl.* **20** (1986) 246–248].

- [10] R. G. Novikov, The inverse scattering problem on a fixed energy level for the two-dimensional Schrödinger operator, *J. Funct. Anal.* **103** (1992) 409–463.
- [11] H. Segur, The Korteweg–de Vries equation and waves. I. Solutions of the equation, *J. Fluid Mech.* **59** (1973) 721–736.
- [12] I. N. Vekua, *Generalized Analytic Functions* (Pergamon Press, Oxford, 1962).
- [13] A. P. Veselov and S. P. Novikov, Finite-zone, two-dimensional, potential Schrödinger operators. Explicit formula and evolutions equations, *Dokl. Akad. Nauk SSSR.* **279** (1984) 20–24 [translation in *Sov. Math. Dokl.* **30** (1984) 588–591].
- [14] A. P. Veselov and S. P. Novikov, Finite-zone, two-dimensional Schrödinger operators. Potential operators, *Dokl. Akad. Nauk SSSR.* **279** (1984) 784–788 [translation in *Sov. Math. Dokl.* **30** (1984) 705–708].
- [15] V. E. Zakharov and S. V. Manakov, Asymptotic behaviour of non-linear wave systems integrated by the inverse scattering method. *J. Exp. Theor. Phys.* **71** (1976) 203–215 [translation in *Sov. Phys. J. Exp. Theor. Phys.* **44**(1) (1976) 106–112].
- [16] V. E. Zakharov and E. I. Shulman, *Integrability of Nonlinear Systems and Perturbation Theory*, What is Integrability? (Springer–Verlag, Berlin, 1991), pp. 185–250.