















- [19] F. S. Wahyuni *et al.*, “A new ring-reduced tetraprenyltoluquinone and a prenylated xanthone from *Garcinia cowa*,” *Aust. J. Chem.*, vol. 57, no. 3, pp. 223–226, 2004, doi: 10.1071/CH03175.
- [20] F. A. Tanha, “Antioxidant and Anti-microbial Activity Assessment of Methanol Extract of *Garcinia cowa* Leaves,” pp. 1–90, 2013.
- [21] S. E. Burdall, A. M. Hanby, M. R. J. Lansdown, and V. Speirs, “Breast cancer cell lines: Friend or foe?,” *Breast Cancer Res.*, vol. 5, no. 2, pp. 89–95, 2003, doi: 10.1186/bcr577.
- [22] S. W. Lim, H. S. Loh, K. N. Ting, T. D. Bradshaw, and Z. N. Allaudin, “Reduction of MTT to purple formazan by vitamin E isomers in the absence of cells,” *Trop. Life Sci. Res.*, vol. 26, no. 1, pp. 111–120, 2015.
- [23] A. Bahuguna, I. Khan, V. K. Bajpai, and S. C. Kang, “MTT assay to evaluate the cytotoxic potential of a drug,” *Bangladesh J. Pharmacol.*, vol. 12, no. 2, pp. 115–118, 2017, doi: 10.3329/bjp.v12i2.30892.
- [24] Y. Liu and D. Peterson, “Mechanism of cellular 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) reduction,” *J. ...*, vol. 69, no. 2, pp. 581–93, 1997, [Online]. Available: <http://www.ncbi.nlm.nih.gov/pubmed/9231715> %5Cn<http://onlinelibrary.wiley.com/doi/10.1046/j.1471-4159.1997.69020581.x/full>.
- [25] Y. L. Franco, T. R. Vaidya, and S. Ait-Oudhia, “Anticancer and cardio-protective effects of liposomal doxorubicin in the treatment of breast cancer,” *Breast Cancer Targets Ther.*, vol. 10, pp. 131–141, 2018, doi: 10.2147/BCTT.S170239.
- [26] R. Mustarichie, D. Runadi, and D. Ramdhani, “The antioxidant activity and phytochemical screening of ethanol extract, fractions of water, ethyl acetate, and n-hexane from mistletoe tea (*Scurrula atropurpurea* BL. dans),” *Asian J. Pharm. Clin. Res.*, vol. 10, no. 2, pp. 343–347, 2017, doi: 10.22159/ajpcr.2016.v10i2.15724.