



Precision Rehabilitation for Healthy Longevity: A MetaMotionRL-Based Framework Integrating Traditional Chinese Medicine Exercise

Zhenguo Luo¹, Xiang Liu^{1*}, Qianyun Fu¹, Dunhua Zou³, Anhui Tao²

¹ Urban Vocational College of Sichuan, Chengdu, China

² Sichuan Xingke Urban Transportation Advanced Technical School, Chengdu, China

³ Chengdu Xindu District Guicheng Shizudao Health Management Co., Ltd., Chengdu, China

liuxiangjkd@qq.com

Abstract. Narrowing the gap between lifespan and healthspan requires objective, scalable rehabilitation tools. While Traditional Chinese Medicine (TCM) exercises improve balance, their deployment is constrained by subjective evaluations and lack of quantitative metrics. This study presents a "Precision Rehabilitation" framework integrating TCM exercises with research-grade MetaMotionRL (MMRL) inertial sensors. We operationalize qualitative TCM concepts ("Rooting", "Flow") into quantitative biomechanical metrics, specifically Root Mean Square (RMS) acceleration and Jerk analysis, establishing a closed-loop biofeedback system. Sensor deployment at the Mingmen point (GV4) approximates center-of-mass sway. Case-driven evidence in geriatric balance disorders demonstrates that high-precision sensing detects subtle stability gains overlooked by conventional clinical scales. Furthermore, we outline a Triple-Helix governance model to facilitate community-based deployment, bridging ancient wellness practices with digital health technology for healthy longevity.

keywords: Precision Rehabilitation, Healthy Longevity, Traditional Chinese Medicine Exercise, Inertial Measurement Unit (IMU), Sustainable Healthcare Management, Lifespan Innovation, Case-Driven Evidence.

1. Introduction

1.1 The Global Imperative: Redefining Longevity in the Era of Lifespan Innovation

Modern demographics show a widening gap between lifespan and healthspan, with postural instability and falls representing major threats to elderly functional independence [6, 11]. Although Traditional Chinese Medicine (TCM) exercises like Taijiquan and Baduanjin effectively enhance neuromuscular control [1, 4, 13], their clinical integration faces an "experience trap." Standard evaluations rely on subjective observations of concepts like "rooting" or "internal power," causing metric ambiguity and limiting scalable, longitudinal monitoring [8, 12]. Consumer wearables lack the 16-bit resolution and high sampling rates required to capture low-velocity, micro-postural oscillations, creating an urgent need for research-grade sensing frameworks.

1.2 The Clinical and Cultural Efficacy of Traditional Chinese Medicine

To bridge qualitative TCM practice with modern biomechanics, this study proposes a data-driven precision rehabilitation framework using the MetaMotionRL (MMRL) 9-axis inertial measurement unit (IMU). By mapping qualitative movement concepts into a structured biomechanical ontology, the system establishes an objective biofeedback loop (Fig. 1) that supports individualized exercise adaptation and scalable health governance.

1.3 Problem Statement: The “Experience Trap” in TCM Rehabilitation

Despite clinical efficacy, integrating TCM rehabilitation into modern healthcare is constrained by an "experience trap"—a management bottleneck where evaluation relies on qualitative, expert-dependent judgment [8, 12]. This presents three key challenges:

- **Metric Ambiguity:** Foundational constructs like "rooting" (Xia Pan Wen Gu) lack standardized biomedical definitions.
- **Episodic Monitoring:** Assessment is restricted to brief clinical encounters rather than continuous home or community tracking.
- **Scalability Barriers:** Expertise remains concentrated among few specialists, reinforcing healthcare inequalities.

Furthermore, consumer-grade wearables lack the 16-bit resolution and multi-axis sensitivity required to detect subtle postural instabilities, underscoring the need for research-grade sensing.

1.4 Strategic Entry: A Data-Driven Framework Using MetaMotionRL

To address these challenges, this study introduces a high-precision, data-driven framework using the MetaMotionRL (MMRL) 9-axis inertial sensor. By translating qualitative TCM knowledge into a structured biomechanical ontology, the framework establishes a real-time, objective biofeedback model (Fig. 1). This bridges traditional intuition with standardized digital health, transitioning experience-driven care into scalable, data-informed precision rehabilitation for healthy longevity.




Figure 1. Schematic Representation of the Closed-Loop Precision Rehabilitation Framework. The diagram illustrates the integrated biofeedback loop, transitioning from raw data acquisition via the MetaMotionRL (MMRL) platform to signal processing and TCM-informed clinical decision support. It highlights the flow from physiological sensing to actionable rehabilitation feedback.

2. Interdisciplinary Theoretical Framework

To substantiate the proposed data-driven rehabilitation strategy, this section presents an interdisciplinary theoretical framework that bridges Traditional Chinese Medicine with contemporary biomechanics and systems-oriented perspectives.

2.1 Biomechanical Ontology

We map qualitative TCM principles ("Xing Shen He Yi") to measurable biomechanical variables. "Rooting" (stability) corresponds to Acceleration Root Mean Square (RMS) and sway area; "Flow" (movement continuity) is quantified via Jerk Analysis (temporal derivative of acceleration); and "Centering" (postural alignment) is operationalized using Euler angles (pitch, roll, yaw).

2.2 Biofeedback and Health Economics

The framework operates as a closed-loop system: MMRL captures kinematic data, processing algorithms extract stability features, and real-time corrective feedback is provided. Digitizing these profiles mitigates Baumol's cost disease in labor-intensive rehabilitation, allowing a single specialist to remotely supervise multiple individuals and reducing long-term fall-related healthcare expenditures [6, 11].

2.3 Sustainable Management and Health Economic Implications

Beyond its technical merits, the broader significance of the proposed framework lies in its potential for sustainable, large-scale application. Rehabilitation services are inherently labor intensive and, from a health economics perspective, particularly vulnerable to Baumol's cost disease, wherein productivity growth fails to keep pace with rising labor costs. In societies experiencing rapid population aging, exclusive dependence on one-to-one, expert-driven TCM rehabilitation models may therefore prove economically unsustainable [6,11].

Digital scaling offers a plausible means of mitigating these pressures. Through the use of digitally generated rehabilitation profiles, a single specialist can, in principle, supervise multiple individuals simultaneously, effectively amplifying expert capacity. In addition, the availability of precise, real-time feedback may reduce exercise-related injuries, thereby lowering downstream healthcare expenditures. The portability of MetaMotionRL further enables a shift away from high-cost clinical environments toward community-based or home-based rehabilitation settings.

From a health economics standpoint, the integration of high-precision sensing technologies is hypothesized to contribute to gains in quality-adjusted life

years among older adults. Taken together, this interdisciplinary framework outlines a realistic and sustainable pathway for advancing lifespan innovation.

3. Technical System Design

3.1 Hardware and Mingmen Deployment

The research-grade MetaMotionRL (MMRL) platform utilizes Bosch BMI160/BMM150 sensors featuring 16-bit resolution and uncompressed 100 Hz raw data streams. The IMU is anchored at the Mingmen point (GV4; L3–L4 spinous processes) via a medical-grade band (Fig. 2). GV4 approximates the human Center of Mass (CoM), enabling sensitive detection of global anterior–posterior (AP) and medial–lateral (ML) postural sway.

3.2 Anatomical Deployment and the Mingmen-Centric Strategy

The precision required for medical-grade TCM sports rehabilitation relies on high-fidelity sensing hardware. This study utilizes the MetaMotionRL (MMRL; MbitentLab) platform, a research-grade nine-axis inertial measurement unit (IMU) featuring Bosch Sensortec sensors (BMI160/BMM150) with 16-bit resolution and uncompressed 100 Hz raw data streams [4]. This high sensitivity captures low-velocity physiological micro-tremors without proprietary smoothing distortions [13].

To capture global postural stability rather than isolated limb movements, the MMRL sensor is anchored via a medical-grade elastic band at the Mingmen point (GV4; L3–L4 spinous processes; Fig. 2). Within TCM theory, Mingmen is the central axis for constitutional energy and balance; biomechanically, it closely approximates the external projection of the human Center of Mass (CoM). Positioned at this anatomical pivot, the configuration provides high-sensitivity detection of anterior–posterior (AP) and medial–lateral (ML) sway, offering a quantitative analogue to TCM "lower-body rooting."




Figure 2. Anatomical Deployment of the IMU Sensor at the Mingmen Point (GV4). Positioning the MMRL at the L3–L4 spinous processes ensures proximity to the human center of mass (CoM), capturing global postural stability and AP/ML oscillations.

3.2 Data Flow and Digital Rehabilitation Metrics

Raw IMU streams undergo Kalman-filtering preprocessing and feature extraction before being mapped into three core digital metrics:

Jerk Analysis: Quantification of Movement Flow. In TCM practice, the quality and continuity of movement—frequently termed "flow" (Yuan Huo)—are vital for proper exercise execution. We quantified this characteristic using jerk analysis, where jerk is defined as the temporal derivative of acceleration:

$$Jerk_{Total} = \sqrt{\left(\frac{da_x}{dt}\right)^2 + \left(\frac{da_y}{dt}\right)^2 + \left(\frac{da_z}{dt}\right)^2}$$

Elevated jerk values indicate frequent corrective adjustments or involuntary micro-tremors, signaling compromised movement continuity, whereas lower jerk profiles reflect harmonious, fluid motion [9].

Sway Area: Stability Quantification. Postural stability ("rooting") is evaluated by calculating the CoM sway confidence ellipse derived from MMRL quaternion sensor fusion:

$$SwayArea = \pi \cdot a \cdot b$$

where a and b represent the semi-major and semi-minor axes of the sway trajectory [1, 5]. Smaller sway areas denote superior static equilibrium in geriatric populations.

Spectral Shift Analysis - Fatigue Monitoring. To prevent overexertion, acceleration signals are processed using Fast Fourier Transformation (FFT). Muscle fatigue is identified via spectral compression, manifested as a downward shift in the median signal frequency. Continuous monitoring enables real-time alerts to adjust rest periods, maintaining intervention safety and long-term compliance.

4. Case-Driven Analysis and Digital Prescriptions

4.1 Empirical Case Insights

We evaluated the framework across age-related balance disorders and geriatric frailty [2, 5]:

- Case 1 (Joint Protection): A 68-year-old male practicing Ma Bu (horse stance) experienced knee pain due to unnoticed medial collapse. Roll angle (Φ) monitoring at GV4 detected asymmetric pelvic tilt. Setting a real-time $\Phi > 5^\circ$ threshold triggered haptic alerts, correcting joint alignment immediately.
- Case 2 (Overcoming Plateaus): A 72-year-old female post-hip fracture plateaued on standard clinical scales (e.g., Berg Balance Scale) at week 8 (Fig. 3). Longitudinal MMRL tracking revealed that her Sway Area continued to decline despite flatlining clinical scores, validating ongoing internal balance recovery and guiding dynamic prescription updates.

4.2 Digital Prescription Evolution

Sensor profiles create an adaptive feedback loop. Discrepancies (e.g., elevated jerk) inform TCM diagnostic logic (e.g., Qi deficiency or stability deficits), prompting recursive algorithm updates that adjust exercise duration, intensity, or rest intervals [10].




Figure 3. Longitudinal Trends in Postural Stability: Clinical Scales vs. Biomechanical Metrics. Data derived from Case Study 2, illustrating the "Plateau Effect." While traditional clinical scales (e.g., Berg Balance Scale) show minimal change after week 8, high-precision metrics like Sway Area and Jerk continue to track subtle internal improvements in stability.

5. Sustainable Governance and Community Pathways

5.1 Community Scalability and Triple-Helix Model

The MMRL’s 8MB NAND Flash "Logging Mode" enables asynchronous local recording, bypassing Bluetooth instability in elderly care and supporting a "Practice at Home, Supervise Remotely" workflow [6, 11, 15]. Implementation is governed by a Triple-Helix Model: the Sports Department defines kinematic standards, the TCM Department provides diagnostic logic, and the Technology Hub maintains algorithm infrastructure [13].

5.2 Social Impact and Cross-Cultural Translation

By objective QALY optimization and real-time fall prevention, the framework lowers long-term care costs [7]. As piloted between Polus International College (China) and Payap University (Thailand), converting qualitative "internal flow" into standardized Jerk metrics creates a universal digital protocol for cross-cultural traditional medicine integration.

6. Conclusion

This study establishes an interdisciplinary Precision TCM Rehabilitation framework by operationalizing traditional movement concepts into objective biomechanical metrics via the MetaMotionRL platform. Placing sensors at the Mingmen point provides high-fidelity tracking of sway and smoothness, detecting subtle recovery trends missed by conventional scales. Integrated within a Triple-Helix governance model, this system offers a scalable, data-informed pathway for healthy longevity. Future work will incorporate deep learning models for automated syndrome differentiation across larger clinical cohorts.

Acknowledgments. This study was supported by the Research Project Outcomes of Sichuan Education, Science, Culture and Public Health Trade Union (Grant No. SJKWWKT2026-002)

Disclosure of Interests. The authors have no competing interests to declare that are relevant to the content of this article.

References

1. Cui, Z., Xiong, J., Li, Z., & Yang, C. (2024). Tai chi improves balance performance in healthy older adults: A systematic review and meta-analysis. *Frontiers in Public Health*, 12, Article 1443168. <https://doi.org/10.3389/fpubh.2024.1443168>
2. Dong, R., Hamzah, M. S. G. B., Awang, M. M. B., Qiu, J., & Chen, S. (2026). Effects of Tai Chi on balance and fall prevention in healthy older adults: A randomized controlled meta-analysis. *Frontiers in Public Health*, 13, Article 1638006. <https://doi.org/10.3389/fpubh.2025.1638006>
3. Ercan Yildiz, S., Fidan, O., Gulsen, C., Colak, E., & Genc, G. A. (2024). Effect of dual-task training on balance in older adults: A systematic review and meta-analysis. *Archives of Gerontology and Geriatrics*, 121, Article 105368. <https://doi.org/10.1016/j.archger.2024.105368>
4. Li, L., Guo, S., Ding, B., & Zhang, J. (2024). Effectiveness of Tai Chi exercise on balance, falls, and motor function in older adults: A meta-analysis. *Frontiers in Medicine*, 11, Article 1486746. <https://doi.org/10.3389/fmed.2024.1486746>
5. Lin, J., Ning, S., Lyu, S., Gao, H., Shao, X., & Tan, Z. (2024). The effects of different types of Tai Chi exercises on preventing falls in older adults: A systematic review and network meta-analysis. *Aging Clinical and Experimental Research*, 36, Article 65. <https://doi.org/10.1007/s40520-023-02674-7>

6. Sherrington, C., Fairhall, N. J., Wallbank, G. K., Tiedemann, A., Michaleff, Z. A., Howard, K., Clemson, L., Hopewell, S., & Lord, S. R. (2019). Exercise for preventing falls in older people living in the community. *Cochrane Database of Systematic Reviews*, 1(1), Article CD012424. <https://doi.org/10.1002/14651858.CD012424.pub2>
7. Wang, D., Wang, P., Lan, K., Zhang, Y., & Pan, Y. J. (2020). Effectiveness of tai chi exercise on overall quality of life and its physical and psychological components among older adults: A systematic review and meta-analysis. *Brazilian Journal of Medical and Biological Research*, 53(11), Article e101196. <https://doi.org/10.1590/1414-431X202010196>
8. Wang, Y., Zhang, Y., Liu, J., & Wang, H. (2023). Effects of Baduanjin exercise on the physical function of middle-aged and elderly people: A systematic review and meta-analysis of randomized controlled trials. *BMC Complementary Medicine and Therapies*, 23(1), Article 38. <https://doi.org/10.1186/s12906-023-03866-4>
9. Xie, J., Guo, J., Wang, B., et al. (2024). Comparing the effectiveness of five traditional Chinese exercises in improving balance function in older adults: A systematic review and Bayesian network meta-analysis. *PeerJ*, 12, Article e18512. <https://doi.org/10.7717/peerj.18512>
10. Yang, Y., Li, E., Gong, Z., Tualaulelei, M., Zhao, Z., & Zhang, Z. (2025). Optimal exercise parameters of Baduanjin for balance in older adults: A systematic review and meta-analysis. *Frontiers in Public Health*, 13, Article 1541170. <https://doi.org/10.3389/fpubh.2025.1541170>
11. Yu, H., Zhong, J., Li, M., & Chen, S. (2025). Effects of exercise intervention on falls and balance function in older adults: A systematic review and meta-analysis. *PeerJ*, 13, Article e20190. <https://doi.org/10.7717/peerj.20190>
12. Zhang, X., & Liu, Q. (2022). Baduanjin exercise for balance function in community dwelling older adults. *BMC Complementary Medicine and Therapies*, 22(1), Article 276. <https://doi.org/10.1186/s12906-022-03764-1>
13. Zhang, Y., Liu, Q., & Wang, X. (2025). Effects of Tai Chi and Baduanjin on muscle mass, muscle function, and activities of daily living in individuals with sarcopenia: A systematic review and meta-analysis. *Frontiers in Public Health*, 13, Article 1636857. <https://doi.org/10.3389/fpubh.2025.1636857>

Open Access This chapter is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

