

Fuzzy logics with truth hedges revisited

Francesc Esteva, Lluís Godo, Carles Noguera

IIIA, Artificial Intelligence Research Institute
 CSIC, Spanish National Research Council
 Campus de la Univ. Autònoma de Barcelona s/n
 08193 Bellaterra, Catalonia, Spain
 {esteva,godo,cnoguera}@iiia.csic.es

Abstract

In this paper we build upon previous works of Hájek and Vychodil on the axiomatization of truth-stressing and depressing hedges as expansions of BL logic by new unary connectives. They show that their logics are chain-complete, but standard completeness is only proved for the expansions over Gödel logic. We propose weaker axiomatizations that have as main advantages the preservation of standard completeness properties of the original logic and the fact that any subdiagonal (resp. superdiagonal) non-decreasing function on $[0, 1]$ preserving 0 and 1 is a sound interpretation of the truth stresser (resp. depresser) connectives.

Keywords: Truth hedges, Mathematical Fuzzy Logic, Standard completeness, T-norm based logics.

1. Introduction

In this paper we deal with interpretation of linguistic hedges in the t-norm based fuzzy logics. Typical examples of linguistic hedges in the sense of Zadeh [1] are “very true”, “quite true”, “more or less true”, “slightly true”, etc. They are represented in fuzzy logic systems in broad sense as functions from the set of truth values (typically the real unit interval) into itself, that modify the meaning of a proposition by applying over the membership function of the fuzzy set underlying the proposition.

In order to cope with these linguistic hedges in the setting of mathematical fuzzy logic, Hájek proposes in a series of papers [2, 3, 4] to understand them as truth functions of new unary connectives called truth-stressing or truth depressing hedges, depending on whether they reinforce or weaken the meaning of the proposition they apply over. The intuitive interpretation of a truth-stressing (resp. depressing) hedge like *very true* (resp. *slightly true*) on a chain of truth-values is a subdiagonal (resp. superdiagonal) non-decreasing function preserving 0 and 1. The class of such functions will be called *hedge functions* from now on.

This paper builds upon previous works, mainly those by Hájek [4] and Vychodil [5], on the axiomatization of truth-stressing (resp. depressing) hedges as

expansions of BL logic (and of some of their prominent extensions, like Łukasiewicz or Gödel logics) by a new unary connective *vt*, for *very true*, and another one *st*, for *slightly true*, respectively. The logics they define are shown to be algebraizable and to enjoy completeness with respect to the classes of chains of their corresponding varieties, however not any BL-chain expanded with such functions (called hedges from now on) are models of them, or in other words, belong to the corresponding varieties. Moreover, the defined logics are not proved to enjoy in general standard completeness, except for the case of logics over Gödel logic. One of the main reasons for both problems is the presence in the axiomatizations of the well-known modal axiom K for the *vt* connective,

$$vt(\varphi \rightarrow \psi) \rightarrow (vt \varphi \rightarrow vt \psi),$$

which puts quite a lot of constraints on the hedges to be models of these logics with no natural algebraic interpretation.

Particular classes of truth-stressers have been also addressed in the literature. For instance, the well-known globalization operator Δ (introduced independently by Monteiro in the context of intuitionistic logic [6] and by Baaz in the context of Gödel-Dummett logics [7]) is a limit case of a truth stresser since, over a chain, it sends 1 to 1 and all the other elements to 0, and the intuitive interpretation would be *definitely true*.

In this paper we propose weaker axiomatizations for both the truth-stressing and depressing connectives not imposing any constraint on hedges other than the ones we have mentioned above, and for which we can prove standard completeness.

The paper is structured as follows. In the next section we provide the necessary logical and algebraic preliminaries that will be used in the rest of the paper. In Section 3 we propose a general axiomatization for truth stressers while Section 4 focuses on truth depressers. Related work is discussed in Section 5 and we conclude with some final remarks in Section 6.

2. Preliminaries

In this section we gather from [8, 9] some necessary results we will use in sections 3 and 4.

Let L be a finitary logic in a language \mathcal{L} . We say that L is a *Rasiowa-implicative logic* (c.f. [10]) if there is a binary (either primitive or definable by a formula) connective \rightarrow of its language such that:

- (R) $\vdash_L \varphi \rightarrow \varphi$
- (MP) $\varphi, \varphi \rightarrow \psi \vdash_L \psi$
- (T) $\varphi \rightarrow \psi, \psi \rightarrow \chi \vdash_L \varphi \rightarrow \chi$
- (Cng) $\varphi \rightarrow \psi, \psi \rightarrow \varphi \vdash_L c(\chi_1, \dots, \chi_i, \varphi, \dots, \chi_n)$
 $\rightarrow c(\chi_1, \dots, \chi_i, \psi, \dots, \chi_n)$
for each n -ary $c \in \mathcal{L}$ and each $i < n$.
- (W) $\varphi \vdash_L \psi \rightarrow \varphi$

As proved in [8], every Rasiowa implicative logic is algebraizable in the sense of Blok and Pigozzi [11] and, if it is finitary, its equivalent algebraic semantics, the class of L -algebras, is a quasivariety; call it \mathbb{L} . Every L -algebra satisfies $x \rightarrow x = y \rightarrow y$ for any x, y , and hence the language can be expanded by a definable constant $\bar{1} = p \rightarrow p$. Then, the algebraizability gives the following strong completeness theorem:

For every set $\Gamma \cup \{\varphi\}$ of formulae, $\Gamma \vdash_L \varphi$ iff for every $\mathbf{A} \in \mathbb{L}$ and every \mathbf{A} -evaluation e , $e(\varphi) = \bar{1}^{\mathbf{A}}$, whenever $e[\Gamma] \subseteq \{\bar{1}^{\mathbf{A}}\}$.

Every L -algebra \mathbf{A} is naturally endowed with a pre-order relation by setting for every $a, b \in A$: $a \leq^{\mathbf{A}} b$ iff $a \rightarrow^{\mathbf{A}} b = \bar{1}^{\mathbf{A}}$. \mathbf{A} is called an *L -chain* if $\leq^{\mathbf{A}}$ is a total order¹. L is called a *semilinear logic* iff it is strongly complete with respect to the semantics given by L -chains.

Many systems informally referred to as *fuzzy logics* in the literature are actually finitary Rasiowa-implicative semilinear logics. Well known examples are the three main fuzzy logics G , Π and \mathbb{L} (see e.g. [12]), BL (the logic of all continuous t-norms, see [12, 13]) and MTL (the logic of all left-continuous t-norms, see [14, 15]). A big class of fuzzy logics which are finitary, Rasiowa-implicative and semilinear, and contain the mentioned prominent examples, are the so-called *core fuzzy logics*: the axiomatic expansions of MTL satisfying (Cng) for any possible new connective.

Finally, we need to recall a couple of definitions and results about disjunction connectives, for they provide a useful characterization of semilinearity.

A (primitive or definable) binary connective \vee is called a *disjunction* in L whenever it satisfies:

- (PD) $\varphi \vdash_L \varphi \vee \psi$ and $\psi \vdash_L \varphi \vee \psi$
- (PCP) If $\Gamma, \varphi \vdash_L \chi$ and $\Gamma, \psi \vdash_L \chi$, then $\Gamma, \varphi \vee \psi \vdash_L \chi$

Given a disjunction \vee and an a finitary inference rule $(R) : \Gamma \vdash \varphi$ (axioms are taken as rules with $\Gamma = \emptyset$), we define the \vee -form of (R) , denoted as (R^\vee) , as the rule $\Gamma \vee p \vdash \varphi \vee p$, where p is an arbitrary propositional variable not appearing in $\Gamma \cup \{\varphi\}$.

¹For the sake of a lighter notation, from now on we will drop the super-indexes in the algebra operations whenever no confusion is possible.

Proposition 1. [9] *Let L_1 be a logic with a disjunction \vee and let L_2 be an expansion of L_1 by a set of finitary rules \mathcal{C} . Then, \vee is a disjunction in L_2 iff (R^\vee) holds in L_2 for each $(R) \in \mathcal{C}$. In particular, \vee is a disjunction in any axiomatic expansion of L_1 .*

Proposition 2. [9] *Let L be a finitary Rasiowa-implicative logic with a binary connective \vee satisfying (PD). Consider the following two properties:*

- (P \vee) $\vdash_L (\varphi \rightarrow \psi) \vee (\psi \rightarrow \varphi)$
- (MP \vee) $\varphi \rightarrow \psi, \varphi \vee \psi \vdash_L \psi$ and $\varphi \rightarrow \psi, \psi \vee \varphi \vdash_L \psi$

The following are equivalent:

- (i) \vee is a disjunction and satisfies (P \vee),
- (ii) L is semilinear and satisfies (MP \vee).

As mentioned before, core fuzzy logics are semilinear, hence they are strongly complete with respect to the class of their chains. However, this completeness may be sometimes refined to special subclasses of chains. We will use the following notions of completeness with respect to a given class of chains.

Definition 3 ($\mathbb{K}C$, $FS\mathbb{K}C$, $S\mathbb{K}C$). *Let L be a core fuzzy logic and let \mathbb{K} be a class of L -chains. We say that L has the (finitely) strong \mathbb{K} -completeness property, (F) $S\mathbb{K}C$ for short, when for every (finite) set of formulae T and every formula φ it holds that $T \vdash_L \varphi$ iff $e(\varphi) = \bar{1}^{\mathbf{A}}$ for each \mathbf{A} -evaluation such that $e[T] \subseteq \{\bar{1}^{\mathbf{A}}\}$ for every L -algebra $\mathbf{A} \in \mathbb{K}$. We say that L has the \mathbb{K} -completeness property, $\mathbb{K}C$ for short, when the equivalence is true for $T = \emptyset$.*

Of course, the $S\mathbb{K}C$ implies the $FS\mathbb{K}C$, and the $FS\mathbb{K}C$ implies the $\mathbb{K}C$. When \mathbb{K} is the class of all chains whose support is the real unit interval $[0, 1]$ we will denote it as \mathcal{R} , call its elements as *real* or *standard* chains, and we will speak about *real* or *standard* completeness properties.

3. Truth stressers: a general axiomatization

In order to solve the problems with the axiomatization of truth stressers and depressers proposed by Hájek and Vychodil mentioned in the Introduction section, in what follows we will make use of available results described in Section 2 to obtain a very simple and general axiomatizations with very intuitive properties and nice completeness results. To begin with let us consider the case of truth stressers.

Let L be a core fuzzy logic, and consider L_s the expansion of L with a new unary connective s (for *stresser*) defined by the following additional axioms:

- (VTL1) $s\varphi \rightarrow \varphi$,
- (VTL2) $s\bar{1}$,

and the following additional inference rule:

- (MON) from $(\varphi \rightarrow \psi) \vee \chi$ infer $(s\varphi \rightarrow s\psi) \vee \chi$.

If we denote by \vdash_{L_s} the notion of deduction defined as usual from the above axioms and rules, one can easily show the following provabilities.

Lemma 4. *In L_s the following deductions are valid:*

- (i) $\vdash_{L_s} \neg s\bar{0}$
- (ii) $\varphi \rightarrow \psi \vdash_{L_s} s\varphi \rightarrow s\psi$
- (iii) $\psi \vdash_{L_s} s\psi$
- (iv) $s\varphi, \varphi \rightarrow \psi \vdash_{L_s} s\psi$

Proof. (i) It follows directly from (VTL1) taking $\varphi = \bar{0}$.
(ii) It follows directly from (MON) taking $\chi = \bar{0}$.
(iii) It follows directly from (ii) taking $\varphi = \bar{1}$ and using (VTL2).
(iv) Very easy using (ii) and *modus ponens*. \square

Notice that (iv) is a kind of *stronger version of the modus ponens rule*: if φ implies ψ , and φ is very true, then one can derive that ψ is very true as well.

On the other hand, (ii) shows that (Cng) is satisfied for the new unary connective too. Therefore, the logic L_s is Rasiowa-implicative and its equivalent algebraic semantics is the class of L_s -algebras. An algebra $\mathcal{A} = \langle A, \wedge, \vee, \&, \rightarrow, s, \bar{0}, \bar{1} \rangle$ of type $\langle 2, 2, 2, 2, 1, 0, 0 \rangle$ is an L_s -algebra if it is an L-algebra expanded by a unary operator $s : A \rightarrow A$ (truth-stressing hedge) that satisfies, for all $x, y, z \in A$,

- (1) $s(\bar{1}) = \bar{1}$,
- (2) $s(x) \leq x$,
- (3) if $(x \rightarrow y) \vee z = \bar{1}$ then $(s(x) \rightarrow s(y)) \vee z = \bar{1}$.

It is clear that the class of L_s -algebras forms a quasivariety, call it \mathbb{L}_s . Notice that if $\langle A, \wedge, \vee, \&, \rightarrow, \bar{0}, \bar{1} \rangle$ is a totally ordered L-algebra and $s : A \rightarrow A$ is any non-decreasing mapping such that $s(\bar{1}) = \bar{1}$ and $s(a) \leq a$ for any $a \in A$, then the expanded structure $\langle A, \wedge, \vee, \&, \rightarrow, s, \bar{0}, \bar{1} \rangle$ is an L_s -chain². In other words, in L_s -chains the quasiequation (3) turns out to be equivalently expressed by this simplified form: if $x \rightarrow y = \bar{1}$ then $s(x) \rightarrow s(y) = \bar{1}$, and this condition simply expresses that s is non-decreasing.

Moreover, since the rule (MON) is closed under \vee -forms, by Proposition 1 we know that \vee keeps being a disjunction in the expanded logic. On the other hand, since (P_\vee) was already valid in L, by Proposition 2 we obtain that L_s is also semilinear and hence it is complete with respect to the semantics of all L_s -chains.

Theorem 1. *L_s is complete with respect to the class of all L_s -chains.*

Corollary 5. *In L_s the following deductions are valid:*

²Observe that these three simple conditions required for s would be not enough to define an L_s -chain in case L_s would have been defined with the additional axiom K as in [4].

- (v) $\vdash_{L_s} s(\varphi \vee \psi) \leftrightarrow s\varphi \vee s\psi$
- (vi) $\vdash_{L_s} s(\varphi \wedge \psi) \leftrightarrow s\varphi \wedge s\psi$

Proof. Both properties are easily checked on L_s -chains. \square

One might wonder whether the corresponding equation for the monotonicity of s (i.e. $s(x \wedge y) = s(x) \wedge s(y)$ or, equivalently, $s(x \vee y) = s(x) \vee s(y)$) may substitute the quasiequation (3) in the definition of L_s -algebras. In other words, does the quasivariety \mathbb{L}_s coincide with the variety \mathbb{V} of expansions of L-algebras satisfying the equations (1), (2) and monotonicity of s ? The answer is negative as shown by the following example.

Example 1. *If L is not classical propositional logic, then there must be an L-chain \mathbf{A} with at least three elements on its domain. Take any $a \in A \setminus \{\bar{0}^A, \bar{1}^A\}$ and consider the direct product algebra $\mathbf{A} \times \mathbf{A}$ and expand it with a unary operator s by putting $s(a, b) = \langle a \wedge b, a \wedge b \rangle$. An easy computation shows that, although s satisfies (1), (2) and monotonicity, the rule (MON) is not sound. Indeed, we have $\langle \bar{1}, \bar{1} \rangle \rightarrow \langle \bar{1}, a \rangle \vee \langle a, \bar{1} \rangle = \langle \bar{1}, a \rangle \vee \langle a, \bar{1} \rangle = \langle \bar{1}, \bar{1} \rangle$, while $(s(\bar{1}, \bar{1}) \rightarrow s(\bar{1}, a)) \vee \langle a, \bar{1} \rangle = (\langle \bar{1}, \bar{1} \rangle \rightarrow \langle a, a \rangle) \vee \langle a, \bar{1} \rangle = \langle a, a \rangle \vee \langle a, \bar{1} \rangle = \langle a, \bar{1} \rangle \neq \langle \bar{1}, \bar{1} \rangle$.*

Thus, \mathbb{V} and \mathbb{L}_s coincide over chains but they are different. While L_s is semilinear due to the rule (MON), the logic associated to \mathbb{V} is not. This also shows that in the presentation of L_s (MON) cannot be substituted by the simpler rule: from $\varphi \rightarrow \psi$ infer $s\varphi \rightarrow s\psi$ (which, as we have just seen, is sound in L_s -chains but not for all L_s -algebras).

Similarly, inspired by the well-known presentation of logics with Δ , one might also ask whether (MON) could be substituted by the globalization rule for s : from φ infer $s\varphi$. The answer is again negative.

Example 2. *Consider the finite MTL-chain \mathbf{C} defined over the domain $C = \{0, 1, 2, 3, 4, 5\}$ with the natural order and the following monoidal operation:*

$\&$	0	1	2	3	4	5
0	0	0	0	0	0	0
1	0	1	1	1	1	1
2	0	1	1	1	2	2
3	0	1	1	1	2	3
4	0	1	2	2	4	4
5	0	1	2	3	4	5

Take the MTL-filter $F = \{4, 5\}$ and the following unary operation s :

x	$s(x)$
0	0
1	1
2	1
3	3
4	4
5	5

It is clear that s is subdiagonal, sends the top element to itself and it is non-decreasing. Moreover for every $x \in F$, $s(x) \in F$, i.e. it is sound w.r.t. the globalization rule for s . However, it is not sound w.r.t. the rule (MON). Indeed, $3 \rightarrow 2 = 4 \in F$, while $s(3) \rightarrow s(2) = 3 \rightarrow 1 = 3 \notin F$.

We consider now the issue of completeness of L_s with respect to the distinguished semantics of L_s -chains. One can prove that if L has the finite strong standard completeness property (FSRC), then L_s has it as well. As usual, this can be done by showing that the class L_s -chains is partially embeddable into the subclass of standard L_s -chains.

Theorem 2 (Finite strong standard completeness). *If L is a finite strong standard complete core fuzzy logic, then the logic L_s is finite strong standard complete as well.*

Proof. Assume that L has the FSRC. Take any L_s -chain $\mathbf{A} = \langle A, \wedge, \vee, \&, \rightarrow, s, \bar{0}, \bar{1} \rangle$ and let B be a finite partial subalgebra of \mathbf{A} . We have to show that there exists a standard L_s -chain $\langle [0, 1], \wedge, \vee, *, \Rightarrow, s', 0, 1 \rangle$ and a mapping $f : B \rightarrow [0, 1]$ preserving the existing operations. By assumption, the s -free reduct of \mathbf{A} is partially embeddable into a standard L -chain $\langle [0, 1], \wedge, \vee, *, \Rightarrow, 0, 1 \rangle$. Denote this embedding by f and consider any non-decreasing and subdiagonal function $s' : [0, 1] \rightarrow [0, 1]$ satisfying $s'(f(x)) = f(s(x))$ for every $x \in B$ such that $s(x) \in B$. There are obviously many such functions s' interpolating the set of points $P = \{ \langle f(x), f(s(x)) \rangle \mid x, s(x) \in B \}$, for instance a linear interpolant. Another interpolant can be defined as follows: let $0 = z_1 < \dots < z_n < 1$ be the set of elements of $[0, 1]$ such that $\langle z_i, \cdot \rangle \in P$ and define $s'(1) = 1$ and, for all $z \in [0, 1]$,

$$s'(z) = f(s(x_i)), \text{ if } z_i \leq z < z_{i+1}$$

where $x_i \in B$ is such that $z_i = f(x_i)$. In any case s' makes $\langle [0, 1], \wedge, \vee, *, \Rightarrow, s', 0, 1 \rangle$ an L_s -chain and f a partial embedding of L_s -chains. \square

Actually, this theorem can be generalized to arbitrary classes of L -chains and their s -expansions, proved in a completely analogous way, and yielding a more general result.

Corollary 6. *Let L be a core fuzzy logic, \mathbb{K} a class of L -chains, and \mathbb{K}_s the class of the L_s -chains whose s -free reducts are in \mathbb{K} . If L has the FSRC.*

Theorem 3 (Strong standard completeness). *If L is a strong standard complete core fuzzy logic, then the logic L_s is strong standard complete as well.*

Proof. Assume that L has the SRC. We have to show that any countable L_s -chain can be embedded into a standard L_s -chain. Let \mathbf{A} be a countable L_s -chain. By the assumption, the s -free reduct of \mathbf{A} is embeddable into a standard L -chain $\mathbf{B} =$

$\langle [0, 1], \wedge, \vee, *, \Rightarrow, 0, 1 \rangle$. Denote this embedding by f and define $s' : B \rightarrow B$ in the following way: for each $z \in [0, 1]$, $s'(z) = \sup\{f(s(x)) \mid x \in A, f(x) \leq z\}$. So defined, s' is a non-decreasing and subdiagonal function such that $s'(f(x)) = f(s(x))$ for any $x \in A$ and hence \mathbf{B} expanded with s' is a standard L_s -chain where \mathbf{A} is embedded. \square

Observe that the proof of the previous theorem can be repeated whenever the linear order of the chains is complete. Therefore we obtain the following corollary.

Corollary 7. *Let L be a core fuzzy logic, \mathbb{K} a class of completely ordered L -chains, and \mathbb{K}_s the class of the L_s -chains whose s -reducts are in \mathbb{K} . If L has the SKC, then L_s has the SK $_s$ C.*

4. The case of truth depressors

Very similarly to the case of truth stressers, we can proceed to define an axiomatization for the case of truth depressors just by replacing axioms (VTL1) and (VTL2) with dual versions (STL1) and (STL2) (for *slightly true*). Namely, given a core fuzzy logic L , we define L_d as the expansion of L with a new unary connective d , the following additional axioms

$$\begin{aligned} \text{(STL1)} \quad & \varphi \rightarrow d\varphi, \\ \text{(STL2)} \quad & \neg d\bar{0}, \end{aligned}$$

and the following additional inference rule

$$\text{(MON)} \text{ from } (\varphi \rightarrow \psi) \vee \chi \text{ infer } (d\varphi \rightarrow d\psi) \vee \chi.$$

Being a kind of dual version of L_s , many properties are proved in a completely analogous way:

Lemma 8. *In L_d the following deductions are valid:*

- (i) $\vdash_{L_d} d\bar{1}$
- (ii) $\varphi \rightarrow \psi \vdash_{L_d} d\varphi \rightarrow d\psi$
- (iii) $\neg\varphi \vdash_{L_d} \neg d\varphi$
- (iv) $\vdash_{L_d} \neg d\varphi \rightarrow \neg\varphi$
- (v) $d\varphi, \varphi \rightarrow \psi \vdash_{L_m} d\psi$

Proof. (i) It follows directly from (STL1) taking $\varphi = \bar{0}$.

(ii) It follows directly from (MON) taking $\chi = \bar{0}$.

(iii) It follows from (ii) for $\psi = \bar{0}$ and (STL2).

(iv) It follows directly from (STL1) using the fact that $(\varphi \rightarrow \psi) \rightarrow (\neg\psi \rightarrow \neg\varphi)$ is derivable in MTL.

(v) Very easy using (ii) and *modus ponens*. \square

It is interesting to remark that (v) provides a kind of *weaker* or modified version of *modus ponens* with the truth-depressor: if φ implies ψ , and φ is *slightly true*, then one can derive that ψ is *slightly true* as well.

Again, (ii) shows that the (Cng) condition is satisfied for the new unary connective too. Therefore, the logic L_d is Rasiowa-implicative and its

equivalent algebraic semantics is the class of L_d -algebras. An algebra $\mathcal{A} = \langle A, \wedge, \vee, \&, \rightarrow, s, \bar{0}, 1 \rangle$ of type $\langle 2, 2, 2, 2, 1, 0, 0 \rangle$ is an L_d -algebra if it is an L -algebra expanded by a unary operator $d : A \rightarrow A$ (truth-depressing hedge) that satisfies, for all $x, y, z \in A$,

- (1') $d(0) = 0$,
- (2') $x \leq d(x)$,
- (3') if $(x \rightarrow y) \vee z = \bar{1}$ then $(d(x) \rightarrow d(y)) \vee z = \bar{1}$.

Analogously to the case of truth-stressers, every L -chain \mathbf{A} can be expanded to an L_d -chain by adding an arbitrary non-decreasing mapping $d : A \rightarrow A$ such that $d(\bar{0}) = \bar{0}$ and $x \leq d(x)$ for every $x \in A$.

Also, since the lattice disjunction keeps satisfying the (PCP) in the expanded logic, L_d is semilinear and hence it is complete with respect to the semantics of all L_d -chains. As a straightforward consequence, we have for d an analogous result to Corollary 5 for s

Lemma 9. *In L_d the following deductions are valid:*

- (v) $\vdash_{L_d} d(\varphi \vee \psi) \leftrightarrow d\varphi \vee d\psi$
- (vi) $\vdash_{L_d} d(\varphi \wedge \psi) \leftrightarrow d\varphi \wedge d\psi$

If we modify Example 1 by taking $s(a, b) = \langle a \vee b, a \vee b \rangle$, it shows that in the context of truth depressers the rule (MON) cannot be substituted by simple monotonicity. Similarly, Example 2 can be modified by taking the function d :

x	$d(x)$
0	0
1	1
2	2
3	4
4	4
5	5

and then it shows that in the presentation of L_d (MON) cannot be substituted by the following rule: from $\neg\varphi$ infer $\neg d\varphi$. Indeed, d is superdiagonal, sends the bottom element to itself and it is non-decreasing, it satisfies $\neg d(x) \in F$ whenever $\neg x \in F$; however, $3 \rightarrow 2 = 4 \in F$, while $d(3) \rightarrow d(2) = 4 \rightarrow 2 = 3 \notin F$.

Finally, analogous proofs of those of Corollaries 6 and 7 allows us to obtain this theorem about the preservation of completeness properties.

Theorem 4 (Completeness properties). *Let L be a core fuzzy logic, \mathbb{K} a class of L -chains and \mathbb{K}_d the class of L_d -chains whose d -free reducts are in \mathbb{K} . Then:*

- (i) *If L has the FSKC, then L_d has the FSK $_d$ C.*
- (ii) *If L has the SKC and all the chains in \mathbb{K} are completely ordered, then L_d has the SK $_d$ C.*

5. Related work

As mentioned in the introduction, there are two main references when talking about the formalisation of truth-stressing hedges within the framework of mathematical fuzzy logic. The first one is Hájek's paper [4] where he axiomatizes over BL a logic for the hedge *very true*. The second one is the paper by Vychodil [5] where he extends Hájek's analysis to truth-depressing hedges. In the rest of this section we overview these and some related logics and compare them with our proposal.

5.1. On Hájek's logic of "very true"

Hájek defines the logic BL_{vt} as the expansion of BL with a new connective vt and the following axioms

- (VE1) $vt\varphi \rightarrow \varphi$,
- (VE2) $vt(\varphi \rightarrow \psi) \rightarrow (vt\varphi \rightarrow vt\psi)$,
- (VE3) $vt(\varphi \vee \psi) \rightarrow (vt\varphi \vee vt\psi)$.

and the following necessitation inference rule:

- (NEC) necessitation for vt : from φ infer $vt\varphi$

If we define the corresponding logic L_{vt} with this kind of truth stresser over a core fuzzy logic L instead of BL (with the same axioms and rules), it turns out, by simple inspection, that L_{vt} can be obtained from the logic L_s by replacing the (MON) inference rule by the axiom (VE2) and the necessitation rule for vt (NEC). In general, this makes L_{vt} a strictly stronger logic than L_s . An exception is when L is Gödel logic, since in that case G_{vt} and G_s are equivalent. This is due to the fact that in G_s the axiom (VE2) is derivable.

Hájek proves completeness of BL_{vt} with respect to the class of linearly ordered BL_{vt} -algebras, which extends to any axiomatic extension of BL, but the issue of standard completeness is left as an open problem, except for the case of Gödel logic.

A relevant further study of logics with truth stressers can be found in the paper by Ciabattoni et al. [16], that makes significant contributions in various respects. They basically consider extensions of MTL as base logics to be expanded with a unary connective (called *modality* in [16]), they consider three possible additional axioms to be added to Hájek axiomatics, and they develop proof systems for the arising new logics and study their algebraic and completeness properties. Particularly relevant for our purposes, they consider for a given logic L that is an extension of MTL the following logics:

$$\begin{aligned} L\text{-KT}^r &= L + (\text{VE1}) + (\text{VE2}) + (\text{VE3}) + (\text{NEC}) \\ L\text{-S4}^r &= L\text{-KT}^r + (\text{VE4}) \quad vt\varphi \rightarrow vt(vt\varphi) \end{aligned}$$

Axiom (VE4), together with axiom (VE1) forces the truth-stressing hedges to be closed over their image, i.e. that $vt\varphi$ has to be equivalent to $vt(vt\varphi)$.

Notice that Hájek's BL_{vt} logic is nothing but the logic $BL\text{-KT}^r$. Moreover, in their paper they prove

standard completeness of the L-S4^r logics for different choices for L: namely MTL, SMTL, C_nMTL, IMTL and C_nIMTL.

Other papers dealing with particular types of truth stressers are [2] where truth-stressers are closure operators while in [3, 17] where truth-stressers are further required to map any element to the greatest idempotent (w.r.t. to the monoidal operation) below it. Finally, observe that adding the axiom $vt\varphi \vee \neg vt\varphi$ to L-KT^r, vt turns to be equivalent to the well-know Baaz-Monteiro projection connective Δ .

5.2. Vychodil's logic of "slightly true"

Vychodil first introduces in [5] a logic combining both a truth-streaser and a truth depresser. Indeed the logic BL_{vt,st} is defined as an expansion of Hájek's BL_{vt} logic with a new unary connective "slightly true" denoted by st and with the following additional axioms

- (ST1) $\varphi \rightarrow st\varphi$,
- (ST2) $st\varphi \rightarrow \neg vt\neg\varphi$,
- (ST3) $vt(\varphi \rightarrow \psi) \rightarrow (st\varphi \rightarrow st\psi)$

This logic is proved to be complete with respect to the class of all linearly-ordered BL_{vt,st}-algebras (defined in the obvious way). Note that axioms (ST1) and (ST2) put into relation both connectives vt and st . Vychodil also proposes two slightly different axiomatizations (systems I and II) for the truth depressing hedge *slightly true* alone. They are defined again as expansions of BL with the unary connective st . Namely, system the (I) has the following set of additional axioms:

- (DH1) $\varphi \rightarrow st\varphi$,
- (DH2) $\neg st(\bar{0})$,
- (DH3) $st(\varphi \rightarrow \psi) \rightarrow (st\varphi \rightarrow st\psi)$

while the system (II) consists of the axioms (DH1), (DH2) and

- (DH4) $(\varphi \rightarrow \psi) \rightarrow (st\varphi \rightarrow st\psi)$

Both systems also have the following rule of inference:

- (RN_{st}) from $\neg\varphi$ infer $\neg st\varphi$.

Chain-completeness for both systems is proved, but again the issue of standard completeness is left open.

Notice that axioms (DH1) and (DH2) correspond exactly to (STL1) and (STL2) of our logic L_d, and that the inference rule (RN_{st}) is derivable from the rule (MON) using axiom (STL2). So, again, the main difference between Vychodil's logics and our proposal is the presence of the K-like axioms (DH3) and (DH4), which do not appear in the logics L_d.

6. Conclusions

In this paper we have been concerned in providing new axiomatic systems L_s and L_d for the truth hedges *very true* and *slightly true*, by weakening the proposed systems by Hájek and Vychodil, and defined over any core fuzzy logic L. The main advantage of the proposed systems with respect to the previously ones is that we can show standard completeness, i.e. completeness with respect to the class of standard L-chains expanded by arbitrary (stressing and depressing) hedges. The price paid in this process is that the class of corresponding algebras cannot be shown in general to be a variety any longer, but only a quasi-variety. Actually it remains as an open problem to prove or disprove whether they form in fact a variety in the general case (for instance, for L being Gödel logic it is indeed a variety). An easy solution to get a variety would be to make use of the Baaz-Monteiro Δ operator. In case this operator is definable in L, as it happens e.g. when L is the n -valued Łukasiewicz logic L_n (where $\Delta\varphi$ is defined as $\varphi \& .n. \&\varphi$) or when L is an extension of SBL_~, the expansion of SBL with an involutive negation \sim (where $\Delta\varphi$ is defined as $\neg\sim\varphi$), then the (MON) inference rule in L_s can equivalently be replaced by the axiom

$$(MON_{\Delta}) \Delta(\varphi \rightarrow \psi) \rightarrow (s\varphi \rightarrow s\psi)$$

and analogously for the d connective in L_d. If Δ is not definable in L, then we can expand the base logic L with the operator Δ together with its usual set of axioms and necessitation rule (see e.g. [12]) and consider the logics L_{Δ,s} and L_{Δ,d} instead, where the (MON) inference is again replaced by the above (MON_Δ) axioms.

Finally, to study logics combining both kinds of hedges along the line proposed in this paper is also a matter of future research.

Acknowledgments The authors acknowledge partial support of the Spanish projects FFI2008-03126-E/FILO, TIN2009-14704-C03-03 and TIN2010-20967-C04-01.

References

- [1] Lotfi A. Zadeh. A fuzzy set theoretic interpretation of linguistic hedges. *Journal of Cybernetics*, 2(3):4–34, 1972.
- [2] Petr Hájek and Dagmar Harmancová. A hedge for Gödel fuzzy logic. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 8(4):495–498, 2000.
- [3] Petr Hájek. Some hedges for continuous t-norms logics. *Neural Network World*, 12(2):159–164, 2002.
- [4] Petr Hájek. On very true. *Fuzzy Sets and Systems*, 124(3):329–333, 2001.

- [5] Vilém Vychodil. Truth-depressing hedges and BL-logic. *Fuzzy Sets and Systems*, 157(15):2074–2090, 2006.
- [6] António A. Monteiro. Algebrés de Heyting symétriques. *Portugaliae Mathematica*, 39(1–4):1–239, 1980.
- [7] Matthias Baaz. Infinite-valued Gödel logic with 0-1-projections and relativisations. In Petr Hájek, editor, *Gödel’96: Logical Foundations of Mathematics, Computer Science, and Physics*, volume 6 of *Lecture Notes in Logic*, pages 23–33. Springer-Verlag, Brno, 1996.
- [8] Petr Cintula and Carles Noguera. Implicational (semilinear) logics I: Basic notions and hierarchy. *Archive for Mathematical Logic*, 49(4):417–446, 2010.
- [9] Petr Cintula and Carles Noguera. A general framework for fuzzy logics. In Petr Cintula, Petr Hájek, and Carles Noguera, editors, *Handbook of Mathematical Fuzzy Logic*. College Publications, 201x. To appear.
- [10] Helena Rasiowa. *An Algebraic Approach to Non-Classical Logics*. North-Holland, Amsterdam, 1974.
- [11] Willem J. Blok and Don L. Pigozzi. *Algebraizable Logics*, volume 396 of *Memoirs of the American Mathematical Society*. American Mathematical Society, Providence, RI, 1989.
- [12] Petr Hájek. *Metamathematics of Fuzzy Logic*, volume 4 of *Trends in Logic*. Kluwer, Dordrecht, 1998.
- [13] Roberto Cignoli, Francesc Esteva, Lluís Godo, and Antoni Torrens. Basic fuzzy logic is the logic of continuous t-norms and their residua. *Soft Computing*, 4(2):106–112, 2000.
- [14] Francesc Esteva and Lluís Godo. Monoidal t-norm based logic: Towards a logic for left-continuous t-norms. *Fuzzy Sets and Systems*, 124(3):271–288, 2001.
- [15] Sándor Jenei and Franco Montagna. A proof of standard completeness for Esteva and Godo’s logic MTL. *Studia Logica*, 70(2):183–192, 2002.
- [16] Agata Ciabattoni, George Metcalfe, and Franco Montagna. Algebraic and proof-theoretic characterizations of truth stressers for MTL and its extensions. *Fuzzy Sets and Systems*, 161(3):369–389, 2010.
- [17] Franco Montagna. Storage operators and multiplicative quantifiers in many-valued logics. *Journal of Logic and Computation*, 14(2):299–322, 2004.