

# Main diagenesis of middle-lower Ordovician of Yubei Area in Tarim Basin

Zhao Hanting<sup>1,a</sup>, Fan Tailiang<sup>1,b</sup>

1,Key Laboratory for Marine Reservoir Evolution and Hydrocarbon Accumulation Mechanism,

Ministry of Education of China, China University of Geosciences, Beijing 100083, China

<sup>a</sup>ibwinibwin@hotmail.com, <sup>b</sup>tlfan@cugb.edu.cn

Fund Program: 2011ZX05005-002-010HZ The sequence structure restriction to diagenesis of reservoir in Tarim basin


**Abstract:**According to analysis of cores, rock sections, casting sections, logging data of Ordovician carbonate reservoirs in the Yubei area, reservoir rock in the study area include 4 types: mud micrite limestone, grain limestone, biological limestone and dolomitic limestone. In combination with core photos and imaging logging analysis, reservoir can be divided into pores and cracks. Diagenesis mainly includes compaction, pressolution, cementation, filling, recrystallization, dolomitization, dissolution. To sum up, the main constructive diagenesis include recrystallization, dolomitization, dissolution, otherwise the main destructive diagenesis include compaction, pressolution, cementation, filling.

**Key words** Yubei area; Tarim; Ordovician; Diagenesis;

## 1. Introduction

Yubei area is located in the eastern margin slope of Tarim basin, whose northern border is Bachu uplift, Mazhatage fault zone, eastern border extends to Tangguzibasi sag, and southern border Yecheng—Hetian sag of south-western depression. The Ordovician strata distribution is complex, by means of Yubei No.1 NE fault zone, it can be divided into three small units, namely the western Yubei platform area, NE structural belt and Yudong fault depression zone (Figure 1). NE and nearly EW fracture and local structures developed very well in the area. Well Yubei 1 lies on NW fault zone, which shows "Y" shape. Multiple fault anticline traps developed in the top of the Ordovician Yingshan formation, which developed in middle-late Caledonian and late Hercynian<sup>[1-2]</sup>.

In recent years, along with the deepening of exploration, there are more researches to discuss the characteristics and genetic analysis of the Ordovician reservoir in Tarim basin, putting forward a variety of the main controlling factors of diagenesis<sup>[3-5]</sup>. On the basis of a large number of core observation and thin section identification, using of the geochemical data, author probes into the Ordovician diagenesis of this area, so as to provide the basis for exploration and development.



**Fig.1 Tectonic location of Yubei area in Tarim Basin**

## 2. Main diagenesis


There are various types of diagenesis. It includes cementation, dissolution, compaction and pressolution, dolomitization, filling and recrystallization<sup>[6-10]</sup>.

### 2.1 Cementation

The Ordovician carbonate reservoir has experienced three periods of cementation, respectively, the submarine cementation, atmospheric fresh-water cementation and burial cementation in Yubei area.

#### a. submarine cementation.

It occurred in the sea bottom diagenetic environment, and cement is fine columnar, fibrous, radiation fibrous, vertical grain growth (oolitic, arenic) wall, isochoric thickness rim and micritic set. In Yingshan calcarenite formation lots of micritic set, fiber, horse tooth shape and isochoric thickness rim cement can be microscopically observed (FIG 2-1); The distribution of cement was dense, and some grains were seriously damaged because of later dissolution, but almost have been preserved owing to the strong submarine cementation.



Well YD4, 5403.12m, O<sub>1-2y</sub>, sparry (algae) calcarenite    Well YB8, 6794.47m, O<sub>3-l</sub>, sparry (oolitic) calcarenite    Well YB6A, 6360.26m, O<sub>2-yj</sub>, sparry calcarenite


**Fig.2-1 submarine cementation characteristics**

#### b. atmospheric fresh-water cementation

It included atmospheric fresh water vadose zone and hyporheic zone two environments. In vadose zone, there are two types of cement. One is composed of fine grained calcite with crescent or pendulous shape cement, with geopetal structure or vadose silt phenomena. The diameter of vadose zone cement was general < 0.10 mm, covered on the fibre shape cement or directly covered on the grains; hyporheic zone cement were syntaxial and isometric texture and common around acanthosis clastic particles (FIG. 2-2).

#### c. burial cementation


It can be seen in the remaining intergranular pore space which have been filled by the first and



Well YB1,561.5m, $O_{1-2}y$ , sparry (algae) calcarenite    Well YD4,5403.88m, $O_{1-2}y$ , sparry (algae) calcarenite    Well YB8,6966.15m, $O_{1-2}y$ , sparry algae calcarenite

**Fig.2-2 atmospheric fresh-water cementation characteristics**

second phase cement, its content is less. Such as in Yubei 2 well with algal arene and maerl granular limestone, two generations of granular sparry calcite cement were visible, otherwise the third phase of cement were visible in the maerl granular limestone. In the early diagenetic period, firstly filling two phases of granular sparry calcite between maerls. In the middle diagenetic period, the second stage sparry calcite is completely dissolution; Otherwise, after the first phase sparry calcite incomplete dissolution, the third phase the single crystal calcite filled (FIG. 2-3).



YB2 井,5929.35m, $O_{1-2}y$ , sparry maerl limestone


YB2 井,5929.17m, $O_{1-2}y$ , sparry calcarenite

**Fig.2-3 multiple cementation characteristics of well Yubei 2**

## 2.2 Dissolution

The ordovician carbonate reservoir has experienced three periods of dissolution, respectively, penecontemporaneous atmospheric dissolution, burial dissolution and supergene karst dissolution, especially the supergene karst dissolution and burial dissolution was the most of importance. a.penecontemporaneous atmospheric dissolution.

It mainly happened just from deposition to shallow burial stage, forming selective dissolution pore, such as intragranular pore, mold pore and intergranular dissolved pore. The identification marks in the study area are: in the sparry oolitic and calcarenite, fibre shape calcite cement were partly or completely dissolution, and the residual fiber calcite and the next phase granular calcite show unconformity contact. (FIG. 2-4).



Well Pishanbei2,6898.94m, $O_3l$ , sparry granular-clast limestone

Well YB4,5900.88m, $O_3l$ , sparry oolitic limestone


Well YB4 井,5838.6m, $O_3l$ , sparry oolitic limestone

**Fig.2-4 atmospheric dissolution characteristics**

## b. Burial dissolution




Burial dissolution is not selective which developed inter-grain pores, intergranular dissolved pore, soluble seam along suture lines, etc. its identification marks are(FIG. 2-5): (1) harbor shaped edge;(2) within the dolomite intracrystalline hole and siliceous dissolved pore, filling black bituminous and black carbonaceous bitumen also visible in edge;(3) bitumen also visible in solution pores around the suture line;



**Fig.2-5 burial dissolution characteristics**

### c.Supergene karst dissolution


Its identification marks (FIG. 2-6) : (1) pyrites occurred ferritization;(2) iron oxide visible within the sutures and microfracture;(3) shale along the suture line corrosion.



**Fig.2-6 Supergene karst dissolution characteristics**

### 2.3 Compaction and pressolution

Compaction and pressolution is the most basic diagenesis in Yubei area, one of its main performance is a particle dense filled compaction, characterized by particle flattening stretched, fragmentations. Another is chemical compaction effect existing in the form of sutures (chart I - VIII, figure 2-7).



**Fig.2-7 Compaction and pressolution characteristics**

## 2.4 Dolomitization


dolomitization occurs mainly on the top of yingshan formation in the fault zone and the bottom of yingshan formation-penglaiba formation in the whole Yubei region . It can be divided into pene-sedimentary dolomitization and burial dolomitization.

### (1) penecontemporaneous dolomitization


Penecontemporaneous dolomitization mainly forms the layer micritic dolomite, arene mud - powder crystal dolomite with better crystal shape, mainly from idiomorphism, hypidiotopic crystal (FIG. 2-8).

### (2) burial dolomitization

The burial dolomitization is the most common dolomitization in Yubei region. The grain is relatively bulky, microtek - coarse grain;Crystal shape is mainly idiomorphism, hypidiotopic, with a inlay contact between grains , apparent wavy extinction, fog heart and bright edges, metasomatism and burial, recrystallization characteristics.The burial dolomitization is common on the top of yingshan formation, the bottom of yingshan formation-penglaiba formation in fault zone; very rare on Lianglitage formation-Yijianfang formation (FIG. 2-9).



**Fig.2-8 penecontemporaneous dolomitization characteristics in Yubei area**



**Fig.2-9 burial dolomitization characteristics in Yubei area**

## 2.5 Filling


Filling is also a kind of very common destructive diagenesis in YuBei region . Cave and cleft are filled by calcite, shale, dolomite, siliceous, asphaltenes, organic matter and so on ,and the most common filling is dolomite.Overall, Cave and cleft filling in YuBei region showed the following characteristics : Calcite and a little of shale is mainly holes fillings in Lianglitage formation and upper Yingshan formation at the platform area; the upper Yingshan formation in the fault belt is filled by calcite, shale mainly, and filled by organic matter (oil), asphaltenes, siliceous and dolomite partly; semi-calcite filling , asphaltene and siliceous are mainly holes fillings in the bottom member of Yingshan formation- Penglaiba formation.Thus, the caves in Lianglitage formation, Yijianfang




formation, and the upper Yingshan formation which is not in the fault belt are basically full-filled, and a little of holes are half-filled, In upper Yinshan formation in the fault belt and the bottom member of Yingshan formation- Penglaiba formation, there are many caves and cleft was not filled, both of them are isolated with poor connectivity (FIG. 2-10).

## 2.6 Recrystallization

Recrystallization in Yubei areas are also more common constructive diagenesis. In the bottom of yingshan formation-penglaiba formation, it is common, showing the mud - powder crystal recrystallization formed fine microtek dolomite, partially recrystallization into coarse grain dolomite, and mineral structure be thicker, throat also be more smooth and flat and intercrystalline pore increasing, providing more conducive to later dissolution channels, forming more intergranular pore and intergranular dissolved pore. Recrystallization cannot form a large storage space, but it can improve the original microscopic pore structure of rock. According to 47 casting microvoid structure in Yubei area, dolomite intercrystal pore, intergranular dissolved pore rate was 4% (FIG. 2-11).



**Fig.2-10 Filling characteristics in Yubei area**



**Fig.2-11 Recrystallization characteristics in Yubei area**

## Conclusion

According to the study above, main diagenesis of Yubei area in Tarim basin is divided into cementation, dissolution, compaction and pressolution, dolomitization, filling and recrystallization. The constructive diagenesis includes dissolution, dolomitization and recrystallization; while the destructive diagenesis includes cementation, compaction and pressolution and filling.

## Reference

- [1] Zhang Zhongpei, Liu Shilin, Yang Ziyu, et al. Tectonic evolution and its petroleum geological significances of the Maigaiti Slope, Tarim Basin [J]. Oil & Gas Geology, 2011, 32(6) : 909-918.
- [2] Qiao Guilin, Qian Yixiong, Cao Zicheng, et al. Reservoir characteristics and karst model of Ordovician

Yingshan Formation in Yubei area,Tarim Basin[J]. Petroleum Geology & Experiment,2014,36(4):416-428.

[3] Yang Haijun, Han Jianfa, Sun Chonghao, et al. A development model and petroleum exploration of Karst reservoirs of Ordovician Yingshan Formation in the northern slope of Tazhong palaeouplift[J].Acta Petrolei Sinica, 2011, 32(2): 199-205.

[4] Han Jianfa, Mei Lianfu, Yang Haijun, et al. The study of hydrocarbon origin, transport and accumulation in Tazhong area,Tarim Basin [J]. Natural Gas Geoscience, 2007, 18 (3):426-435.

[5] Wang Zhaoming, Zhao Kuanzhi, Wu Guanghui, et al. Characteristics and main controlling factors of the Upper Ordovician reef-bank reservoir development in the Tazhong I slope-break zone[J]. Oil & Gas Geology, 2007, 28(6): 797-801.

[6] Wang Zhaoming, Wang Qinghua, Wang Yuan. Conditions and controlling factors of hydrocarbon accumulation of Hetianhe Gas Field, Tarim Basin[J]. Marine Origin Petroleum Geology, 2000,5(1/2): 124-132.

[7] Lizhong,Huang Sijing,Liu Jiaqing,et al. Buried diagenesis, structurally controlled thermal fluid process and their effect on ordovician carbonatereservoirs in Tahe, Tarim Basin[J]. Acta Sedimentology sinica,2010,28(5):969-979.

[8] Luo Chunshu, Yang Haijun, Li Jianghai, et al. Characteristics of high quality Ordovician reservoirs and controlling effects of faults in the Tazhong area, Tarim Basin[J]. Petroleum Exploration and Development, 2011, 38(6): 716-724.

[9] Wang Zhenyu,Yang Liuming,Mafeng,et al. Origin of dolomite of Lower Ordovician Yingshan Formation in Tazhong area[J]. Lithologic Reservoirs,2012,24(2):21-23.

[10] Zhang Nai, Zhao Zongju, Xiao Zhongyao, et al. Characteristics of hydrocarbon fluid in the Ordovician vein calcite of Tazhong No.1 slope-break zone[J]. Natural Gas Geoscience, 2010, 21(3):389-396.