

Application of Chebyshev Polynomial in Annual Electricity Consumption

Lai-bin GAO^{*} College of Information and Technology, Jilin Agricultural University Changchun, China 130118 wanggao911@163.com

Abstract—based on statistical data of National Statistical Bureau of China, and giving the least-squares fitting of Chebyshev polynomial, the data of electricity consumption in per capita from 1983 to 2013 is analyzed by GM (1,1) method and Chebyshev polynomial least squares method respectively. The results showed that Chebyshev polynomial least squares fitting method is excellent, per capita energy consumption from 2014 to 2018 is predicted by this method.

Keywords-GM (1,1); least squares; Chebyshev polynomial

In the scientific research and production practice, the data obtained from observation, prediction and calculation should be analyzed and processed, and then the conclusions concerning the problems can be found out. The polynomial least square method is an effective method, because the accuracy and reliability of the least square method relies on the condition number of the coefficient matrix of its normal equation. Chebyshev polynomial is adopted as basis function, so as to make the small condition number of coefficient matrix of the normal equation of the least square method, and then the accuracy and reliability of the least square method are improved obviously. As China's increasingly affluent urban and rural living and modern, per capita electricity consumption increasing every year, energy conservation has become important part of China's energy conservation. Therefore, that the electricity consumption per capita is predicted accurately is very important for socioeconomic development.

I. BASIC THEORY

A. GM (1, 1) Prediction Model [1][2]

Let non-negative sequence $X^{(0)}$ of the original data sequence as follow

$$X^{(0)} = \left\{ X^{(0)}(1), X^{(0)}(2), \cdots, X^{(0)}(n) \right\}$$

Where

Where

$$X^{(0)}(k) \ge 0, k = 1, 2, \cdots, n$$

The sequence $X^{(1)}$ obtained is as follow

$$X^{(1)} = \left\{ X^{(1)}(1), X^{(1)}(2), \cdots, X^{(1)}(n) \right\}$$

Bo WANG College of Information and Technology, Jilin Agricultural University Changchun, China 130118 zhanghq_@126.com

Where

$$X^{(1)}(k) = \sum_{i=1}^{k} X^{(0)}(i) , k = 1, 2, \dots, n.$$

 $Z^{(1)}$ Denotes a sequence close to the mean

$$Z^{(1)} = \left\{ Z^{(1)}(2), Z^{(1)}(3), \cdots, Z^{(1)}(n) \right\},\$$

Where

$$Z^{(1)}(k) = 0.5X^{(1)}(k) + 0.5X^{(1)}(k-1)$$

k = 2,3,...,n.

Named $x^{(0)}(k) + az^{(1)}(k) = b$ the GM (1, 1) model, Where a, b is solution of the model parameters?

$$Y_n = (x^{(0)}(2), x^{(0)}(3), \dots, x^{(0)}(n))^T$$
.

$$B = \begin{bmatrix} -\frac{1}{2}Z^{(1)}(2) & 1\\ -\frac{1}{2}Z^{(1)}(3) & 1\\ \vdots & \vdots\\ -\frac{1}{2}Z^{(1)}(n) & 1 \end{bmatrix}$$

Algorithm for parameters with a minimum two columns

a

$$\hat{a} = \begin{pmatrix} a \\ b \end{pmatrix} = (B^T B)^{-1} B^T Y_n$$

294



Named $\frac{dx^{(1)}}{dt} + ax^{(1)} = b$ the albino equation of the gray differential equation $x^{(0)}(k) + az^{(1)}(k) = b$

Albino equation or that the time response function

$$\hat{x}^{(1)}(t) = \left(x^{(1)}(0) - \frac{b}{a}\right)e^{-at} + \frac{b}{a}.$$

GM (1, 1) gray differential equation $x^{(0)}(k) + az^{(1)}(k) = b$ For the time response

$$\hat{x}^{(1)}(k+1) = \left(x^{(1)}(0) - \frac{b}{a}\right)e^{-ak} + \frac{b}{a} \\ k = 1, 2, \dots, n$$

Take

$$x^{(1)}(0) = x^{(0)}(1)$$

$$x^{(1)}(k+1) = \left(x^{(0)}(1) - \frac{b}{a} \right) e^{-ak} + \frac{b}{a} \quad k = 1, 2, \cdots, n$$

Restore value

$$\hat{x}^{(0)}(k+1) = \hat{x}^{(1)}(k+1) - \hat{x}^{(1)}(k) \quad k = 1, 2, \dots, n$$

 B. The Linear Least Square of Chebyshev Polynomial [3] The expression of the Chebyshev polynomial is that:

$$I_k(t) = \cos(k \arccos t) \quad k \ge 0$$

In the interval [-1,1], the Chebyshev polynomial can be obtained by the following recurrence relations :

$$T_{k+1}(t) = 2xT_{k}(t) - T_{k-1}(t) \quad k = 1, 2, \cdots$$
$$T_{0}(t) = 1$$
$$T_{1}(t) = t$$

By transformation, the Cebyshev polynomial can be obtained in the interval [a,b]:

$$t = \frac{2x - (a+b)}{b-a}, \quad x \in [a,b], \quad t \in [-1,1]$$

$$\widetilde{T}_k(x) = T_k(t) = T_k(\frac{2x - (a+b)}{b-a})$$

Giving a set of data $\{(x_i, y_i)\}_{i=1}^m$,

$$a = x_1 < x_2 < \dots < x_m = b$$

Let $\widetilde{T}_k(x)$, $k = 0, 1, \dots, n$ as the basis function, c_0, c_1, \dots, c_n is minimum value point of

$$\varphi(x) = c_0 + c_1 \widetilde{T}_1(x) + \cdots + c_n \widetilde{T}_n(x) ,$$

$$E_2(c_0, c_1, \cdots + c_n) = \sum_{i=1}^m |y_i - \varphi(x_i)|^2$$

$$= \sum_{i=1}^m \left[\sum_{j=0}^n c_j \widetilde{T}_j(x_i) - y_i \right]^2$$

Let $\widetilde{T}_j = [\widetilde{T}_j(x_1), \cdots \widetilde{T}_j(x_m)]^T$ $j = 0, 1, \cdots, n$,

 $y = [y_1, \dots, y_m]^T$, $C = [c_0, \dots, c_n]^T$. therefore its normal equation of the least square is

$$GC = b$$
,

Where $G = [(\tilde{T}_j, \tilde{T}_k)]$, $(\tilde{T}_j, \tilde{T}_k)$ denotes the inner product of \tilde{T}_j and \tilde{T}_k ; $b = [(y, \tilde{T}_0), \dots, (y, \tilde{T}_n)]^T$. The condition number of the coefficient matrix of this normal equation is very tiny, so the fitting accuracy and practicability is high.

II. THE PREDICTION OF ELETRICITY CONSUMPTION PER CAPITA

A. Fitting Function

The data of electricity consumption per capita from 1983 to 2013 is analyzed and electricity consumption per capita from 2014 to 2018 is predicted by the least square fitting method. Data from 1983 to 2013 is as Table 1.

By using the Matlab mathematic software, the fitting result of GM(1, 1) as follows:

$$p(x) = 200.7845 \exp(0.10931x) - 187.3845$$

The fitting result of Cebyshev polynomial least square is that

Let
$$a = 1983$$
 $b = 2013$
 $t = \frac{2x - (a+b)}{b-a}$, then

 $\phi(t) = 181.0926 + 231.9297t + 81.7784 \times (2t^2 - 1)$



Figure1 The actual fitting curve energy consumption per capita from 1983 to 2013

It can be seen from Fig.1 that the effect of GM(1, 1) is bad, but the fitting result of the Cebyshev polynomial least square is more accurate.

B. Evaluation on the Fitting Results

1) Evaluation on unbiased variance^[5].

The formula is used to evaluate the unbiased variance is:

$$MSE = \frac{1}{N} \sum_{i=1}^{N} e_i^2 = \frac{1}{N} \sum_{i=1}^{N} (y_i - \hat{y}_i)^2$$

Where \hat{y}_i is the fitting evaluation value of x_i . When using the unbiased variance to evaluate, the usually followed principle is the smaller the mean square error, the better the method used. Of GM(1, 1), *MSE* MSE=569.8124, but the linear least square of the Chebyshev polynomial, *MSE* = 315.8785.

2) Determination of the coefficient[4]

The formula of the determination coefficient is as follows:

$$R^{2} = 1 - \frac{\sum (y_{i} - \hat{y}_{i})^{2}}{\sum (y_{i} - \overline{y})^{2}}$$

In the formula, \overline{y} is the average value of y_i . When using the determination coefficient R^2 to evaluate, usually we follow the principle that R^2 closes to 1 is the best method. GM(1, 1) is $R^2 = 0.9879$, but the linear least square Chebyshev polynomial is $R^2 = 0.9948$.

TABLE1 THE DATA OF ELECTRICITY CONSUMPTION PER CAPITA FROM 1983 TO 2013

Year	Yield	Year	Yield
1983	13.4	1999	108.6
1984	15.3	2000	115.0
1985	21.2	2001	126.5
1986	23.2	2002	138.3
1987	26.4	2003	159.7
1988	31.2	2004	184.0
1989	35.3	2005	221.3
1990	42.2	2006	255.6
1991	47.2	2007	308.3
1992	54.9	2008	331.9
1993	62.5	2009	336.0
1994	72.7	2010	383.1
1995	83.5	2011	418.1
1996	87.7	2012	460.4
1997	98.6	2013	515.0
1998	104.2		

III. CONCLUSIONS AND DISCUSSIONS

Firstly, when analyzing the data of scientific research, the linear least square of the Chebyshev polynomial is better method.

Secondly, by establishing the Chebyshev polynomial least square model of the data of energy consumption per capita from 1983 to 2013, electricity consumption in annual per capita from 2014 to 2018 can be predicted, which is 568.9249kwh, 625.1516kwh, 685.4718kwh, 750.0408kwh, 819.0140kwh, respectively.

Thirdly, with the world energy shortage, but also with low carbon and improvement of consciousness environmental protection and energy conservation, people naturally explore to seek new energy, especially renewable energy. In recent years, photovoltaic power generation is a kind of energy, which can provide power to the people's life and economic growth. Photovoltaic power generation has gradually been associated with industry, agriculture, service industry project and people's life

REFERENCES

- L.Fu, gray system theory and its application [M], Science and Technology Literature Press, 1991
- [2] J.L. Deng, the basic method of gray system [M], Hua zhong University Press, 1992: P131-135.
- [3] C. S. LIN. Numerical Analysis [M]. Beijing: Science Press .2007:P174-178.
- [4] M. T. HEATH. Scientific Computing [M].Beijing: Tsinghua University Press .2005:P59-61



- [5] S. MAO CHENG YM. Tutorial of Probability and mathematical statistics [M]. Beijing: Higher Education Press .2004:418-419.
- [6] C. F. DING. Application of Combination forecasting method in China's grain output prediction [J]. Research of Agricultural Modernization, 2007 (3):10-103.