

REFERENCES

- [1] A. Listkowski, M. Pietraszkiewicz, G. Accorsi, J. Mohanraj, "Solid state photoluminescence of novel lanthanide complexes based on 4-benzoylpyrazolone Schiff base," *Synth. Met.* Vol. 160, pp. 2377-2380, November 2010.
- [2] C.R. Bhattacharjee, G. Das, P. Goswami, P. Mondal, S.K. Prasad, D.S.S. Rao, "Novel photoluminescent lanthanidomesogens forming bilayer smectic phase derived from blue light emitting liquid crystalline, one ring O-donor Schiff-base ligands," *Polyhedron*, Vol. 30, pp.1040-1047, April 2011.
- [3] A. Kagkalar, V. Bekiari, E. Stathatos, G.S. Papaefstathiou, C.P. Raptopoulou, T.F. Zafropoulos, et al, "Photoluminescence and electroluminescence by gallium(III) complexes of N-salicylidene-o-aminophenol and its derivatives," *J Lumin.*, Vol. 129, pp. 578-583, May 2009.
- [4] M. Altaf, H. Stoeckli-Evans, "Nickel(II) and copper(II) complexes of 2-(2-pyridyl)benzimidazole: synthesis and structural characterization," *Transition Met. Chem.* Vol. 34, pp.613-620, June 2009.
- [5] W.G. Quirino, C. Legnani, M. Cremona, P.P. Lima, S.A. Junior, O.L. Malta, "White OLED using β -diketones rare earth binuclear complex as emitting layer," *Thin Solid Films*, Vol. 494, pp.23-27, January 2006.
- [6] R. Reyes, M. Cremona, E.E.S. Teotonio, H.F. Brito, O.L. Malta, "Voltage color tunable OLED with (Sm,Eu)- β -diketonate complex blend," *Chem. Phys. Lett.* Vol. 396, pp.54-58, September 2004.
- [7] F.F. Chen, Z.Q. Bian, C.H. Huang, "Progresses in electroluminescence based on europium(III) complexes," *J. Rare Earth.* Vol. 27, pp.345-355, June 2009.
- [8] H.G. Liu, S. Park, K. Jang, W.S. Zhang, H.J. Seo, Y.I. Lee, "Different photoluminescent properties of binary and ternary europium chelates doped in PMMA," *Mater. Chem. Phys.* Vol. 82, pp. 84-92, September 2003.
- [9] Y.L. Zhao, D.J. Zhou, G.Q. Yao, C.H. Huang, "Strong Luminescent Terbium(III)-Pyrazolone Complexes and Their Langmuir-Blodgett Films," *Langmuir* Vol. 13, pp. 4060-4065, July 1997.
- [10] Z.G. Yu, B. Liu, Z.H. Jiang, "Characterization by Spectra and Identification by HPLC-MS of Schiff Base Derived from 1-Phenyl-3-methyl-4-naphthylaceto-4,5-dihydropyrazol-5-one (PMNAP) and 2-Aminobenzimidazole (2-AB) and Its Transitional Metal Complexes," *Chinese J. Org. Chem.* Vol. 29, pp. 1217-1222, August 2009.
- [11] J.Z. Li, H.Q. Zhang, H.X. Li, P.Z. Che, T.C. Wang, "1-(4-Chlorophenyl)-4-(2-furyl)-3-(2-furyl)-1H-pyrazol-5-ol," *Acta Cryst E*, Vol. 63, pp.1289-1290, March 2007.
- [12] W.Z. Lv, B Liu, S.H. Wei, Y. Gao, "Synthesis and characteristics of 2-aminobenzimidazole under microwave irradiation," *Chem. Reagents* Vol. 28, pp. 751-752, December 2006.
- [13] S.P. Mital, R.V. Singh, J.P. Tandon, "Imine derivatives of lanthanons," *Curr. Sci.* Vol. 49, pp.130-132, April 1980.
- [14] R.N. Jadeja, J.R. Shah, E. Suresh, P. Paul, "Synthesis and structural characterization of some Schiff bases derived from 4-[(arylimino)ethyl]-3-methyl-1-(4'-methylphenyl)-2-pyrazolin-5-one and spectroscopic studies of their Cu(II) complexes," *Polyhedron* Vol. 23, pp. 2465-2474, October 2004.