









(Sato, 1992)[14]. The Sr isotopic data obtained by the present study support the conclusion proposed by Blum et al. (1992)[1] that the depositional age of sedimentary target materials is close to 170Ma (Jurassic). Mixing calculations based on the model proposed by Ho and Chen (1996)[2] for various amounts and combinations of target rocks indicate that the best fit for tektites from the Dalat area is a mixture of 39% shale, 29% sandstone, 32% greywacke, 1% quartzite.

## 5. References

- [1] J. D. Blum , D. A. Papanastassiou, C. Koeberl, and G. J. Wasserburg, "Nd and Sr isotopic study of Australasian tektites: New constraints on the provenance and age of target material", *Geochim. Cosmochim. Acta*, 56, pp. 483-492, 1992.
- [2] K. S. Ho, and J. C. Chen, "Geochemistry and origin of tektites from the penglei area, Hainan province, southern China." *J. of Southeast Asian Earth Sciences* Vol. 13, No. 1, pp. 61-72, 1996.
- [3] A. Montanari, and C. Koeberl, "Impact Stratigraphy: The Italian Record." Springer, Heideberg, 2000.
- [4] L. Folco, B. P. Glass, M. D'Orazio, P. Rochette, "A common volatilization trend in Transantarctic Mountain and Australasian microtektites: Implications for their formation model and parent crater location." *Earth and Planetary Science Letters* 293, pp.135-139, 2010.
- [5] F. Moynier, P. Beck, F. Jourdan, Q. Z. Yin, U. Reimold, C. Koeberl, "Isotopic fractionation of zinc in tektites." *Earth and Planetary Science Letters* 277, pp. 482-489, 2009.
- [6] B. P. Glass, "Tektites and microtektites: key facts and inferences." *Tectonophysics* 171, pp. 393-404, 1990.
- [7] F. Moynier, C. Koeberl, P. Beck, F. Jourdan, P. Telouk, "Isotopic fractionation of Cu in tektites." *Geochimica et Cosmochimica Acta* 74, pp. 799-807, 2010.
- [8] Y. T. Lee, J. C. Chen, K. S. Ho and W. S. Juang, "Geochemical studies of tektites from East Asia." *Geochem. Jour.* 38, pp. 1-17, 2004.
- [9] B. P. Glass, H. Huber and C. Koeberl "Geochemistry of Cenozoic microtektites and clinopyroxene-bearing spherules." *Geochim. Cosmochim. Acta* 68, pp. 3971-4006, 2004.
- [10] B. P. Glass and J. E. Pizzuto, "Geographical variation in Australasian microtektite concentrations: Implications concerning the location and size of the source crater." *J. Geophys. Res.* 99, pp. 19075-19081, 1994.
- [11] C. Koeberl, "Geochemistry and origin of Muong Nong-type tektites." *Geochim. Cosmochim. Acta* 56, pp. 1033-1064, 1992.
- [12] C. Koeberl, "Geochemistry and origin of Muong Nong-type tektites." *Geochim. Cosmochim. Acta* 56, pp. 1033-1064, 1992.
- [13] G. Schmidt, L. Zhou, J. T. Wasson, "Iridium anomaly associated with the Australian tektite-producing impact: Masses of the impactor and of the Australian tektites." *Geochim. Cosmochim. Acta* 57, pp. 4851-4859, 1993.
- [14] T. Sato, "Regional geology and stratigraphy : Southeast Asia and Japan." In *The Jurassic of the Circum-Pacific* (Edited by Westermann G. E. G.), pp. 194-213. Cambridge University Press, 1992.