

Academy of Sciences for the Envisat ASAR image data. Besides, the authors also would like to thank National Natural Science Foundation of China, No. 41201335 and No.41001201 for the support.

REFERENCES

- [1] X.H.Shen, Q.Q.Liu, and C.Jiang, "The research discussed on effect increasing yield of soy bean with different ridges cultivation," *Journal of Agriculture*, vol.1, no.6, pp.8-11, 2011.
- [2] W.Li, J.L.Xiao, and H.Tan, "Study on crop net photosynthetic rate and yield under different ridge cultivation," *Science and technology for China rural prosperity*, vol.14, no.11, pp.24-26, 2010.
- [3] S.H.Shi, B.H.Liu, and Q.Fu, "Research on effect of different direction of ridge in increasing yield of maize," *Heilongjiang Agricultural Sciences*, vol.34, no.5, pp.23-25, 2011.
- [4] H.Liu, H.D.Guo, "The extraction of texture information from radar image and its value in geo-science analysis-using a SIR-B image in huangyangzhen area(Gansu province) as a example," *Remote Sensing of Environment China*, vol.10, no.2, pp.107-113, 1995.
- [5] S.Berberoglu, P.J.Curran, and C.D.Lloyd, "Texture classification of Mediterranean land cover," *International Journal of Applied Earth Observation and Geo-information*, vol.9, no.3, pp.322-334, 2007.
- [6] J.Q.Zhang, Q.She, and L.Pan, "Change detection of residential area by remote sensing image based on LBP/C texture," *Geomatics and Information Science of Wuhan University*, vol.33, no.1, pp.7-11, 2008.
- [7] C.Y.Zhu, C.Z.Lan, and G.W.Qin, "Habitation area extraction by the method of intensity separation from SAR texture image," *Jouranal of Image and Graphics*, vol.8, no.6, pp.616-619, 2003.
- [8] A.Baraldia, F.Parmiggiani, "An investigation of the texture characteristics associated with Gray Level Co-occurrence Matrix statistical parameters," *IEEE Transactions on Geoscience and Remote Sensing*, vol.32, no.2, pp.293-304, 1995.
- [9] L.F.Liu, Y.H.Chen, and J.Li, "Texture analysis methods used in remote sensing images," *Remote Sensing Technology and Application*, vol.18, no.6, pp.441-447, 2003.
- [10] F.Wu, C.Wang, and H.Zhang, "Residential areas extraction in high resolution SAR image based on texture features," *Remote Sensing Technology and Application*, vol.20, no.1, pp.148-152, 2005.
- [11] R.M.Haralick, K.Shamugam, and I.Dinstein, "Texture features for image classification," *IEEE Transaction on systems, man, and cybernetics*, vol.SMC-3, no.6, pp.610-621, 1973.
- [12] J.G.Zhang, T.N.Tan, "Brief review of invariant texture analysis methods," *Pattern Recognition*, vol.35, pp.735-747, 2002.
- [13] S.Zhang, "Wetland information extraction combined with texture features," *Remote Sensing Information*, vol.0, no.3, pp.30-34, 2010.
- [14] D.F.Zhang, "MATLAB digital image processing," *Mechanical Industry Press*, pp.210-216, 224-228, 2009.
- [15] F.Y.Shin, S.X.Cheng, "Automatic seeded region growing for color image segmentation," *Image and Vision Computing*, vol.23, no.10, pp.877-886, 2005.
- [16] Q.Y.Yu, "IRGS: Image segmentation using edge penalties and region growing," *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol.30, no.12, pp.2126-2139, 2008.